


Families of Curves with Prescribed Singularities



Thomas Keilen

Vom Fachbereich Mathematik
der Universität Kaiserslautern
zur Verleihung des akademischen Grades
Doktor der Naturwissenschaften
(Doctor rerum naturalium, Dr. rer. nat.)
genehmigte Dissertation.

1. Gutachter: Prof. Dr. G.-M. Greuel
2. Gutachter: Prof. Dr. E. Shustin

Vollzug der Promotion: 30.10.2001

To my parents
Hildegard and
Eduard Keilen

The picture on the titlepage shows a singular curve on an affine chart of the Hirzebruch surface $\mathbb{F}_0 = \mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$ embedded into $\mathbb{P}_{\mathbb{C}}^3$ as a quadric.

Preface

The study of families of curves with prescribed singularities has a long tradition. Its foundations were laid by Plücker, Severi, Segre, and Zariski at the beginning of the 20th century. Leading to interesting results with applications in singularity theory and in the topology of complex algebraic curves and surfaces it has attained the continuous attraction of algebraic geometers since then.

Throughout this thesis we examine the varieties $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ of irreducible reduced curves in a fixed linear system $|D|$ on a smooth projective surface Σ over \mathbb{C} having precisely r singular points of types $\mathcal{S}_1, \dots, \mathcal{S}_r$ – for a more precise definition we refer to Chapter I. We are mainly interested in the following three questions:

- (a) Is $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ non-empty?
- (b) Is $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ \mathbb{T} -smooth, that is smooth of the expected dimension?
- (c) Is $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ irreducible?

That the dimension be the expected one means that the dimension of $|D|$ drops for each imposed singularity type \mathcal{S}_i exactly by the number of conditions imposed by \mathcal{S}_i – e. g. a node imposes one condition, a cusp two.

The simplest possible case of nodal plane curves was more or less completely answered by Severi in the early 20th century. He showed that $V_{|dH|}^{\text{irr}}(rA_1)$, where H is a line in $\mathbb{P}_\mathbb{C}^2$, is non-empty if and only if

$$0 \leq r \leq \frac{(d-1) \cdot (d-2)}{2}.$$

Moreover, he showed that $V_{|dH|}^{\text{irr}}(rA_1)$ is \mathbb{T} -smooth whenever it is non-empty, and he claimed that the variety is always irreducible. Harris proved this claim, which had become known as the Severi Conjecture by then, in 1985 (cf. [Har85b]). Considering more complicated singularities we may no longer expect such complete answers. Hirano provides in [Hir92] a series of examples of irreducible cuspidal plane curves of degree $d = 2 \cdot 3^k$, $k \in \mathbb{N}$, imposing more than $\frac{d(d-3)}{2}$ conditions on $|dH|$ – that means in particular, we may hardly expect to be able to realize all smaller quantities of cusps on an irreducible curve of degree d . Moreover, we see that $V_{|dH|}^{\text{irr}}(rA_2)$ does not necessarily have the expected dimension – examples of this behaviour were already known to Segre (cf. [Seg29]). In 1974 Jonathan Wahl (cf. [Wah74b]) showed that $V_{|104 \cdot H|}^{\text{irr}}(3636 \cdot A_1, 900 \cdot A_2)$ is non-reduced and hence singular. However, its reduction is smooth. The first example, where also the reduction is singular, is due to Luengo. In [Lue87a] he shows that the plane curve C given by $x^9 + z(xz^3 + y^4)^2$ has a single singular point of simple type A_{35} and that

$V_{|9 \cdot H|}^{\text{irr}}(A_{35})$ is non-smooth, but reduced at C . Thus also the smoothness will fail in general. And finally, already Zariski (cf. [Zar35]) knew that $V_{|6 \cdot H|}^{\text{irr}}(6 \cdot A_2)$ consists of two connected components.

The best we may expect thus is to find numerical conditions, depending on the divisor D and certain invariants of the singularities, which imply either of the properties in question. In order to see that the conditions are of the right kind - we then call them *asymptotically proper* -, they should not be too far from necessary conditions respectively they should be nearly fulfilled for series of counterexamples. Let us make this last statement a bit more precise. Suppose that $D = dL$ for some fixed divisor L and $d \geq 0$. We are looking for conditions of the kind

$$\sum_{i=1}^r \alpha(\mathcal{S}_i) < p(d),$$

where α is some invariant of topological respectively analytical singularity types and $p \in \mathbb{R}[x]$ is some polynomial, neither depending on d nor on the \mathcal{S}_i . We say that the condition is *asymptotically proper*, if there is a necessary condition with the same invariants and a polynomial of the same degree. If instead we find an infinite series of examples not having the desired property, where, however, the above inequality is reversed for the same invariants and some other polynomial of the same degree, we say that at least for the involved subclass of singularity types, the condition is *asymptotically proper*. (See also [Los98] Section 4.1.)

While the study of nodal and cuspidal curves has a long tradition, the consideration of families of more complicated singularities needed a suitable description of the tangent space of the family at a point, giving a concrete meaning to “the number of conditions imposed by a singularity type”, that is to the expected dimension of the family. Greuel and Karras in [GrK89] in the analytical case, Greuel and Lossen in [GrL96] in the topological case identify the tangent spaces basically with the global sections of the ideal sheaves of certain zero-dimensional schemes associated to the singularity types (see Definition I.2.6 and Remark I.2.14). This approach - in combination with a Viro gluing type method in the existence case - allows to reduce the existence, T-smoothness and irreducibility problem to the vanishing of certain cohomology groups. Various efforts in this direction culminate in asymptotically proper conditions for the existence (cf. [GLS98c]) and conditions for the T-smoothness and irreducibility, which are better than any previously known ones (cf. [GLS00]). Due to known examples the conditions for the T-smoothness are even asymptotically proper for simple singularities and ordinary multiple points. For an overview on the state of the art in the case of the plane curves we refer to [GrS99].

While the previous investigations mainly considered curves in $\mathbb{P}_{\mathbb{C}}^2$, we study curves on arbitrary smooth projective surfaces and derive results on various

families of surfaces, in many cases the first results in that direction known on these surfaces at all.

In Chapter III we study the question of the *existence* of curves and the main condition (cf. Corollary III.2.5), which we derive in the case of topological singularity types, is of the form

$$\sum_{i=1}^r \delta(\mathcal{S}_i) \leq \alpha D^2 + \beta D \cdot K + \gamma,$$

and for analytical singularity types it is of the form

$$\sum_{i=1}^r \mu(\mathcal{S}_i) \leq \alpha D^2 + \beta D \cdot K + \gamma,$$

for some fixed divisor K and some absolute constants α , β and γ , where $\delta(\mathcal{S}_i)$ is the delta-invariant of \mathcal{S}_i and $\mu(\mathcal{S}_i)$ is its Milnor number. The corresponding necessary condition

$$\sum_{i=1}^r \mu(\mathcal{S}_i) \leq 2 \cdot \sum_{i=1}^r \delta(\mathcal{S}_i) \leq D^2 + D \cdot K_\Sigma + 2$$

is of the same asymptotical behaviour.

Also for the main condition

$$\left(\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1) + \frac{D \cdot K_\Sigma}{2} \right)^2 < \left(r + \frac{K_\Sigma^2}{4} \right)^2 \cdot D^2,$$

which we get in Chapter IV for the *T-smoothness* (cf. Theorem IV.1.1) we cannot expect to do better in general – here for topological singularity types $\tau^*(\mathcal{S}_i) = \tau^{\text{es}}(\mathcal{S}_i)$ is the codimension of the equisingular stratum of \mathcal{S}_i in its semiuniversal deformation, and for analytical types $\tau^*(\mathcal{S}_i) = \tau(\mathcal{S}_i)$, the Tjurina number of \mathcal{S}_i . In the case of simple singularities on plane curves the mentioned examples in [GLS00] show that the conditions are asymptotically proper. However, already for ordinary multiple points this is no longer the case, as is shown there as well. Still, our results apply to surfaces which have not been considered before, and even in the well studied case of nodal curves on quintics in $\mathbb{P}_\mathbb{C}^3$ we get the same sharp results as Chiantini and Serresi (cf. [ChS97]).

We do not know whether our main condition (cf. Theorems V.2.1-V.2.4)

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2$$

in the study of the *irreducibility* in Chapter V shows a similar properness, since neither sufficient conditions of this form nor counterexamples with the same asymptotical behaviour are known – here, γ is some constant, possibly depending on D . And we do not even expect them to be of the right kind. However, the results which we get are for most of the considered surfaces the only

known ones, and even in the well studied plane case they are asymptotically of the same quality than the best previously known conditions.

All our results somehow rely on the vanishing or non-vanishing of certain cohomology groups. The results of Chapter IV and of Chapter V could be formulated completely as a vanishing theorem; and although we derive the existence of curves with prescribed topological or analytical singularities by gluing these into a suitable given curve, we derive the existence of that curve (cf. Theorem III.1.2) again with the aid of a vanishing theorem. The latter is a generalisation of a vanishing theorem of Geng Xu (cf. [Xu95]) and thus of the Kawamata–Viehweg Vanishing Theorem. Chapter II is devoted to its proof.

In Chapter I we introduce the notions used throughout the thesis and state several important facts which are well known. A compact and profound description of the introduced objects and cited results can be found in the thesis of Christoph Lossen [Los98], and for the convenience of the reader we usually refer to this thesis as well as to the original sources.

Apart from the introduction, each chapter of the thesis consists of one or two sections describing the main results possibly followed by a section containing essential technical details of the proofs. At the end of each chapter we examine the derived conditions on several classes of surfaces:

- (a) $\mathbb{P}_{\mathbb{C}}^2$,
- (b) geometrically ruled surfaces,
- (c) product-surfaces,
- (d) surfaces in $\mathbb{P}_{\mathbb{C}}^3$, and
- (e) K3-surfaces.

We have chosen these partly due to their important role in the classification of surfaces, and partly since they all have advantages in their own which make it possible to keep control on the numerical conditions. In the appendix we gather a number of facts which we suppose are well known, but for which we nevertheless could not find a suitable reference. In particular, we give an overview of the properties of the studied surfaces, which we need for the examinations.

Throughout a fixed chapter the theorems, lemmata, definitions and remarks are numbered by sections, e. g. within Section 1 of Chapter V the Theorem 1.1 is followed by Corollary 1.2, while in Section 2 we start again with Theorem 2.1. The same applies to the equations. There is, however, one exception from this rule. In the example sections we reproduce certain statements (equations, theorems, ...) in special cases, and instead of using new numbers we rather keep the old ones adding the letter of the corresponding subsection, e. g. in the example section of Chapter V we have Theorem 2.1a with equation (2.2a) in Subsection 4.a, which is just the appropriate form of Theorem 2.1

in Chapter V. Whenever we refer to a statement within the same chapter, we just cite it by its number. If we, however, refer to it in some other chapter, we add the number of the chapter, e. g. we refer to Theorem 1.1 of Chapter IV within Chapter IV as Theorem 1.1, while in the preface we would cite it as Theorem IV.1.1.

Chapter II and Chapter III are a joint work with Ilya Tyomkin (Tel Aviv University) and have been accepted for publication in the Transactions of the American Mathematical Society. Chapter IV is a slight modification of results already published by Gert-Martin Greuel (Universität Kaiserslautern), Christoph Lossen (Universität Kaiserslautern) and Eugenii Shustin (Tel Aviv University) in [GLS97], and a slightly stronger version may be found in [GLS02]. Finally, Chapter V extends an approach via Bogomolov instability of vector bundles used by Gert-Martin Greuel, Christoph Lossen and Eugenii Shustin in the plane case (cf. [GLS98b]).

Acknowledgements

I would like to express my thanks to my supervisor, Gert-Martin Greuel, for supporting me in many ways during the last years, and I would also like to thank Christoph Lossen, Eugenii Shustin and Ilya Tyomkin for their cooperation while working on this thesis. I owe a special thank to Bernd Kreuzler for many fruitful conversations widening my understanding of algebraic geometry. Finally, I have the pleasure to thank my parents, Hildegard and Eduard Keilen, for their love and care which accompanied me all my life, and Hannah Markwig for her love and for her patience.

Contents

Preface	I
Chapter I. Introduction	1
1. General Assumptions and Notations	1
2. Singularity Schemes	2
Chapter II. A Vanishing Theorem	13
1. The Vanishing Theorem	13
2. Generalisation of a Lemma of Geng Xu	16
3. Examples	19
3.a. <i>The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$</i>	20
3.b. <i>Geometrically Ruled Surfaces</i>	20
3.c. <i>Products of Curves</i>	21
3.d. <i>Products of Elliptic Curves</i>	21
3.e. <i>Surfaces in $\mathbb{P}_{\mathbb{C}}^3$</i>	22
3.f. <i>K3-Surfaces</i>	22
4. Yet Another Vanishing Theorem	23
5. Examples	33
Chapter III. Existence	37
1. Existence Theorem for Ordinary Fat Point Schemes	39
2. Existence Theorem for General Singularity Schemes	44
3. Examples	54
3.a. <i>The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$</i>	54
3.b. <i>Geometrically Ruled Surfaces</i>	55
3.c. <i>Products of Curves</i>	56
3.d. <i>Products of Elliptic Curves</i>	58
3.e. <i>Surfaces in $\mathbb{P}_{\mathbb{C}}^3$</i>	58
3.f. <i>K3-Surfaces</i>	59
Chapter IV. T-Smoothness	61
1. T-Smoothness	63
2. Examples	69

2.a. <i>The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$</i>	70
2.b. <i>Geometrically Ruled Surfaces</i>	72
2.c. <i>Products of Curves</i>	76
2.d. <i>Products of Elliptic Curves</i>	77
2.e. <i>Surfaces in $\mathbb{P}_{\mathbb{C}}^3$</i>	78
2.f. <i>K3-Surfaces</i>	79
Chapter V. Irreducibility	81
1. $V^{\text{irr,reg}}$ is irreducible	83
2. Conditions for the irreducibility of V^{irr}	85
3. The Main Technical Lemmata	88
4. Examples	110
4.a. <i>The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$</i>	110
4.b. <i>Geometrically Ruled Surfaces</i>	111
4.c. <i>Products of Curves</i>	114
4.d. <i>Products of Elliptic Curves</i>	116
4.e. <i>Complete Intersection Surfaces</i>	116
4.f. <i>Surfaces in $\mathbb{P}_{\mathbb{C}}^3$</i>	117
4.g. <i>Generic K3-Surfaces</i>	118
Appendix	121
A. Very General Position	121
B. Curves of Self-Intersection Zero	127
C. Some Facts used for the Proof of the Lemma of Geng Xu	134
D. Some Facts on Divisors on Curves	137
E. Some Facts on Divisors on Surfaces	138
F. The Hilbert Scheme Hilb_{Σ}^n	140
G. Examples of Surfaces	142
G.a. <i>Geometrically Ruled Surfaces</i>	142
G.b. <i>Products of Curves</i>	150
G.c. <i>Products of Elliptic Curves</i>	154
G.d. <i>Surfaces in $\mathbb{P}_{\mathbb{C}}^3$ and Complete Intersections</i>	160
G.e. <i>K3-Surfaces</i>	162
Bibliography	163
Index	169

CHAPTER I

Introduction

1. General Assumptions and Notations

Throughout this thesis Σ will denote a smooth projective surface over \mathbb{C} .

Given distinct points $z_1, \dots, z_r \in \Sigma$, we denote by $\pi: \text{Bl}_{\underline{z}}(\Sigma) = \tilde{\Sigma} \rightarrow \Sigma$ the blow up of Σ in $\underline{z} = (z_1, \dots, z_r)$, and the exceptional divisors π^*z_i will be denoted by E_i . We shall write $\tilde{C} = \text{Bl}_{\underline{z}}(C)$ for the strict transform of a curve $C \subset \Sigma$.

For any smooth projective surface Σ we will denote by $\text{Div}(\Sigma)$ the group of divisors on Σ and by K_{Σ} its canonical divisor. If D is any divisor on Σ , $\mathcal{O}_{\Sigma}(D)$ shall be a corresponding invertible sheaf. A curve $C \subset \Sigma$ will be an effective (non-zero) divisor, that is a one-dimensional locally principal scheme, not necessarily reduced; however, when we talk of an “irreducible curve” this shall include that the curve is reduced. $|D| = |D|_l = \mathbb{P}(H^0(\Sigma, \mathcal{O}_{\Sigma}(D)))$ denotes the system of curves linearly equivalent to D , while we use the notation $|D|_a$ for the system of curves algebraically equivalent to D (cf. [Har77] Ex. V.1.7), that is the reduction of the connected component of Hilb_{Σ} , the Hilbert scheme of all curves on Σ , containing any curve algebraically equivalent to D (cf. [Mum66] Chapter 15).¹ We write $\text{Pic}(\Sigma)$ for the Picard group of Σ , that is $\text{Div}(\Sigma)$ modulo linear equivalence (denoted by \sim_l), $\text{NS}(\Sigma)$ for the Néron–Severi group, that is $\text{Div}(\Sigma)$ modulo algebraic equivalence (denoted by \sim_a), and $\text{Num}(\Sigma)$ for $\text{Div}(\Sigma)$ modulo numerical equivalence (denoted by \sim_n). Given a reduced curve $C \subset \Sigma$ we write $p_a(C)$ for its arithmetical genus and $g(C)$ for the geometrical one.

Given any scheme X and any coherent sheaf \mathcal{F} on X , we will often write $H^v(\mathcal{F})$ instead of $H^v(X, \mathcal{F})$ when no ambiguity can arise. Moreover, if $\mathcal{F} = \mathcal{O}_X(D)$ is the invertible sheaf corresponding to a divisor D , we will usually use the notation $H^v(X, D)$ instead of $H^v(X, \mathcal{O}_X(D))$. Similarly when considering tensor products over the structure sheaf of some scheme X we may sometimes just write \otimes instead of $\otimes_{\mathcal{O}_X}$.

Given any subscheme $X \subset Y$ of a scheme Y , we denote by $\mathcal{I}_X = \mathcal{I}_{X/Y}$ the *ideal sheaf* of X in \mathcal{O}_Y , and for $z \in Y$ we denote by $\mathcal{O}_{Y,z}$ the local ring of Y at z and by $\mathfrak{m}_{Y,z}$ its maximal ideal, while $\hat{\mathcal{O}}_{Y,z}$ denotes the $\mathfrak{m}_{Y,z}$ -adic completion of $\mathcal{O}_{Y,z}$. If X is zero-dimensional we denote by $\#X$ the number of points in its *support* $\text{supp}(X)$, by $\text{deg}(X) = \sum_{z \in Y} \dim_{\mathbb{C}}(\mathcal{O}_{Y,z}/\mathcal{I}_{X/Y,z})$ its *degree*, and by $\text{mult}(X, z) = \max \{m \in \mathbb{N} \mid \mathcal{I}_{X/Y,z} \subseteq \mathfrak{m}_{Y,z}^m\}$ its *multiplicity at z* .

¹Note that indeed the reduction of the Hilbert scheme gives the Hilbert scheme $\text{Hilb}_{\Sigma}^{\text{red}}$ of curves on Σ over reduced base spaces.

If $X \subset \Sigma$ is a zero-dimensional scheme on Σ and $D \in \text{Div}(\Sigma)$, we denote by $|\mathcal{J}_{X/\Sigma}(D)|_1 = \mathbb{P}\left(\mathrm{H}^0(\mathcal{J}_{X/\Sigma}(D))\right)$ the linear system of curves C in $|D|_1$ with $X \subset C$.

If $L \subset \Sigma$ is any reduced curve and $X \subset \Sigma$ a zero-dimensional scheme, we define the *residue scheme* $X : L \subset \Sigma$ of X by the ideal sheaf $\mathcal{J}_{X:L/\Sigma} = \mathcal{J}_{X/\Sigma} : \mathcal{J}_{L/\Sigma}$ with stalks

$$\mathcal{J}_{X:L/\Sigma,z} = \mathcal{J}_{X/\Sigma,z} : \mathcal{J}_{L/\Sigma,z},$$

where “:” denotes the ideal quotient. This leads to the definition of the *trace scheme* $X \cap L \subset L$ of X via the ideal sheaf $\mathcal{J}_{X \cap L/L}$ given by the exact sequence

$$0 \longrightarrow \mathcal{J}_{X:L/\Sigma}(-L) \xrightarrow{\cdot L} \mathcal{J}_{X/\Sigma} \longrightarrow \mathcal{J}_{X \cap L/L} \longrightarrow 0.$$

Let Y be a Zariski topological space (cf. [Har77] Ex. II.3.17). We say a subset $U \subseteq Y$ is *very general* if it is an at most countable intersection of open dense subsets of Y . Some statement is said to hold for points $z_1, \dots, z_r \in Y$ (or $\underline{z} \in Y^r$) *in very general position* if there is a suitable very general subset $U \subseteq Y^r$, contained in the complement of the closed subvariety $\bigcup_{i \neq j} \{z_i = z_j\}$ of Y^r , such that the statement holds for all $\underline{z} \in U$.

2. Singularity Schemes

This thesis is a generalisation of results described in [Los98], and the zero-dimensional schemes with which we are concerned are examined very carefully in [Los98] Chapter 2. We therefore restrict ourselves to a very short description and refer to [Los98] for more details.

2.1 Definition

- (a) Let $f \in \mathcal{O}_{\Sigma,z} \subset \mathbb{C}\{x, y\}$ be given. The germ $(C, z) \subset (\Sigma, z)$ defined by a representative $C = \{z' \mid f(z') = 0\}$ is called a *plane curve singularity*. The \mathbb{C} -algebra $\mathcal{O}_{C,z} := \mathcal{O}_{\Sigma,z}/(f)$ is called the *local ring of* (C, z) .
- (b) Two plane curve singularities (C, z) and (C', z') are said to be *topologically equivalent* if there exists a homeomorphism $\Phi : (\Sigma, z) \rightarrow (\Sigma, z')$ such that $\Phi(C) = C'$.² We write $(C, z) \sim_t (C', z')$. An equivalence class \mathcal{S} of this equivalence relation is called a *topological singularity type*.
- (c) Two plane curve singularities (C, z) and (C', z') are said to be *analytically equivalent* (or *contact equivalent*) if there exists an isomorphism $\Phi^\# : \widehat{\mathcal{O}}_{C',z'} \rightarrow \widehat{\mathcal{O}}_{C,z}$ of the complete local rings. We write $(C, z) \sim_c (C', z')$. An equivalence class \mathcal{S} of this equivalence relation is called an *analytical singularity type*.
- (d) Let $f \in \mathcal{O}_{\Sigma,z}$ and $f' \in \mathcal{O}_{\Sigma,z'}$. We say f and f' are *topologically* respectively *analytically equivalent* if the singularities defined by f and f' are so. We then write $f \sim_t f'$ respectively $f \sim_c f'$.

²This means of course that the equality holds for suitable representatives.

2.2 Remark

We will need a number of well known invariants of topological respectively analytical singularity types, which we are just going to list here. Note that every invariant of the topological type of a singularity is of course also an invariant of its analytical type. Let (C, z) be a reduced plane curve singularity with representative $f \in \mathcal{O}_{\Sigma, z} \subset \mathbb{C}\{x, y\}$.

- (a) The following are known to be invariants of the topological type \mathcal{S} of the plane curve singularity (C, z) .
- (1) $\text{mult}(\mathcal{S}) = \text{mult}(C, z) = \text{ord}(f)$ is the *multiplicity* of \mathcal{S} .
 - (2) $\mu(\mathcal{S}) = \mu(C, z) = \dim_{\mathbb{C}} \mathbb{C}\{x, y\} / \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$ is the *Milnor number* of \mathcal{S} .
 - (3) $\tau^{\text{es}}(\mathcal{S}) = \tau^{\text{es}}(C, z)$ is the codimension of the μ -constant stratum in the semiuniversal deformation of (C, z) .
 - (4) $\kappa(\mathcal{S}) = \kappa(C, z)$, the intersection multiplicity of f with a generic polar curve.
 - (5) $\nu^s(\mathcal{S}) = \nu^s(C, z)$ shall be the minimal integer m such that $m_{\Sigma, z}^m \subseteq I^s(C, z)$ and is called the *topological deformation determinacy*. (Cf. Definition 2.6 for the definition of $I^s(C, z)$.)
 - (6) $\delta(\mathcal{S}) = \delta(C, z) = \dim_{\mathbb{C}} (n_* \mathcal{O}_{\tilde{C}, z} / \mathcal{O}_{C, z})$ is the *delta invariant* of \mathcal{S} , where $n : (\tilde{C}, z) \rightarrow (C, z)$ is a normalisation of (C, z) .
 - (7) $r(\mathcal{S}) = r(C, z)$ is the number of branches of (C, z) .
- (b) For the analytical type \mathcal{S} of (C, z) we have three additional invariants:
- (1) The *Tjurina number* of \mathcal{S} is defined as $\tau(\mathcal{S}) = \tau(C, z) = \dim_{\mathbb{C}} \mathbb{C}\{x, y\} / \left(f, \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$.
 - (2) $\nu^a(\mathcal{S}) = \nu^a(C, z)$ shall be the minimal integer m such that $m_{\Sigma, z}^m \subseteq I^a(C, z)$ and is called the *analytical deformation determinacy*. (Cf. Remark 2.7 for the definition of $I^a(C, z)$.)
 - (3) $\text{mod}(\mathcal{S}) = \text{mod}(C, z)$, the *modality* of the singularity \mathcal{S} . (Cf. [AGZV85] p. 184.)

Throughout the thesis we will very often consider topological and analytical singularity types at the same time. We therefore introduce the notation $\tau^*(\mathcal{S})$ and $\nu^*(\mathcal{S})$, where for a topological singularity type $\tau^*(\mathcal{S}) = \tau^{\text{es}}(\mathcal{S})$ and $\nu^*(\mathcal{S}) = \nu^s(\mathcal{S})$, while for an analytical singularity type $\tau^*(\mathcal{S}) = \tau(\mathcal{S})$ and $\nu^*(\mathcal{S}) = \nu^a(\mathcal{S})$.

2.3 Remark

For the convenience of the reader we would like to gather some known relations between the above topological and analytical invariants of the singularity types corresponding to some reduced plane curve singularity (C, z) .

- (a) $\mu(\mathcal{S}) \leq \mu(\mathcal{S}) + r(\mathcal{S}) - 1 = 2\delta(\mathcal{S})$. (Cf. [Mil68] Chapter 10.)

- (b) $\delta(\mathcal{S}) \leq \tau^{\text{es}}(\mathcal{S})$, since the δ -constant stratum of the semiuniversal deformation of (C, z) contains the μ -constant stratum and since its codimension is just $\delta(\mathcal{S})$ (see also [DiH88]).
- (c) $\kappa(\mathcal{S}) = \mu(\mathcal{S}) + \text{mult}(\mathcal{S}) - 1$.
- (d) $\kappa(\mathcal{S}) \leq 2\delta(\mathcal{S})$. (Cf. [Los98] Lemma 5.12.)
- (e) $\tau^{\text{es}}(\mathcal{S}) = \mu(\mathcal{S}) - \text{mod}(\mathcal{S})$. (Cf. [AGZV85] p. 245.)
- (f) $\tau^{\text{es}}(\mathcal{S}) \leq \tau(\mathcal{S})$, since $\tau(\mathcal{S})$ is the dimension of the base space of the semiuniversal deformation of (C, z) .
- (g) $\tau(\mathcal{S}) \leq \mu(\mathcal{S})$, by definition.
- (h) $\nu^s(\mathcal{S}) \leq \tau^{\text{es}}(\mathcal{S})$. (Cf. [GLS00] Lemma 1.5.)
- (i) $\nu^s(\mathcal{S}) \leq \delta(\mathcal{S})$, if all branches of (C, z) have at least multiplicity three. (Cf. [GLS00] Lemma 1.5.)
- (j) $\nu^a(\mathcal{S}) \leq \tau(\mathcal{S})$. (Cf. [GLS00] Remark 1.9.)
- (k) $\nu^s(\mathcal{S}) \leq \nu^a(\mathcal{S})$, since $I^a(C, z) \subseteq I^s(C, z)$ (cf. Definition 2.6).

Combining these results we get:

$$\nu^s(\mathcal{S}) \leq \left\{ \begin{array}{l} \tau^{\text{es}}(\mathcal{S}) \\ \nu^a(\mathcal{S}) \end{array} \right\} \leq \tau(\mathcal{S}) \leq \mu(\mathcal{S}) \leq \kappa(\mathcal{S}) \leq 2\delta(\mathcal{S}) \leq 2\tau^{\text{es}}(\mathcal{S}) \leq 2\tau(\mathcal{S}).$$

2.4 Remark

It is well known that the simple singularities, that is the singularities with modality zero, form two infinite series A_μ , given by $x^{\mu+1} - y^2$ for $\mu \geq 1$, and D_μ , given by $y \cdot (x^2 - y^{\mu-2})$ for $\mu \geq 4$, together with three exceptional singularities E_μ for $\mu = 6, 7, 8$, given by $x^3 - y^4$, $x^3 - xy^3$ and $x^3 - y^5$ respectively. Moreover, they are the only plane curve singularities \mathcal{S} with $\mu(\mathcal{S}) = \tau(\mathcal{S}) \leq 8$.

The first non-simple family of singularities X_9 , that is, the only one-modular family of singularities of Milnor number 9, is given by $x^4 - y^4 + a \cdot x^2y^2$. The modulus corresponds to the crossratio of the four intersecting lines. Topologically all these singularities are indeed equivalent. Note that, apart from the simple singularities, these are the only singularities \mathcal{S} with $\tau^{\text{es}}(\mathcal{S}) \leq 8$, as a view at the classification table in [AGZV85] Chapter 15 shows, taking the restrictions on $\tau^{\text{es}}(\mathcal{S})$ given in Remark 2.3 into account.

Moreover, from the classification we deduce for an analytical singularity type \mathcal{S} some results which will be useful in Lemma V.3.10.

$\text{mult}(\mathcal{S})$	$\text{mod}(\mathcal{S})$	$\mu(\mathcal{S})$
2	0	$1 \leq \mu(\mathcal{S})$

$\text{mult}(\mathcal{S})$	$\text{mod}(\mathcal{S})$	$\mu(\mathcal{S})$
3	0	$4 \leq \mu(\mathcal{S})$
	1	$10 \leq \mu(\mathcal{S}) \leq 14$
	2	$16 \leq \mu(\mathcal{S})$
	3	$22 \leq \mu(\mathcal{S})$
	≥ 4	$28 \leq \mu(\mathcal{S})$
4	1	$9 \leq \mu(\mathcal{S}) \leq 12$
	2	$15 \leq \mu(\mathcal{S})$
	≥ 3	$22 \leq \mu(\mathcal{S})$

And finally:

$\text{mod}(\mathcal{S})$	$\mu(\mathcal{S})$	$\tau(\mathcal{S})$	$\text{mult}(\mathcal{S})$
0	$1 \leq \mu(\mathcal{S}) = \tau(\mathcal{S}) \leq 8$	$1 \leq \mu(\mathcal{S}) = \tau(\mathcal{S}) \leq 8$	2, 3
1	$9 \leq \mu(\mathcal{S}) \leq 14$	$9 \leq \tau(\mathcal{S}) \leq 14$	3, 4
2	$15 \leq \mu(\mathcal{S})$	$13 \leq \mu(\mathcal{S}) - \text{mod}(\mathcal{S}) \leq \tau(\mathcal{S})$	3, 4
≥ 3	$16 \leq \mu(\mathcal{S})$	$\mu(\mathcal{S}) - \text{mod}(\mathcal{S}) \leq \tau(\mathcal{S})$	$3 \leq \text{mult}(\mathcal{S})$

2.5 Definition

Given distinct points $z_1, \dots, z_r \in \Sigma$ and non-negative integers m_1, \dots, m_r we denote by $X(\underline{m}; \underline{z}) = X(m_1, \dots, m_r; z_1, \dots, z_r)$ the zero-dimensional subscheme of Σ defined by the ideal sheaf $\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}$ with stalks

$$\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma, z} = \begin{cases} \mathfrak{m}_{\Sigma, z_i}^{m_i}, & \text{if } z = z_i, i = 1, \dots, r, \\ \mathcal{O}_{\Sigma, z}, & \text{else.} \end{cases}$$

We call a scheme of the type $X(\underline{m}; \underline{z})$ an *ordinary fat point scheme*.

2.6 Definition

Let $C \subset \Sigma$ be a reduced curve.

- (a) The scheme $X^{\text{ea}}(C)$ in Σ is defined via the ideal sheaf $\mathcal{J}_{X^{\text{ea}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X^{\text{ea}}(C)/\Sigma, z} = I^{\text{ea}}(C, z) = I^{\text{ea}}(f) = \left(f, \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) \subseteq \mathcal{O}_{\Sigma, z},$$

where x, y denote local coordinates of Σ at z and $f \in \mathcal{O}_{\Sigma, z}$ a local equation of C . $I^{\text{ea}}(C, z)$ is called the *Tjurina ideal* of the singularity (C, z) , and it is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point of C . We call $X^{\text{ea}}(C)$ the *equianalytical singularity scheme* of C .

- (b) We define the zero-dimensional subscheme $X^{\text{es}}(C)$ of Σ via the ideal sheaf $\mathcal{J}_{X^{\text{es}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X^{\text{es}}(C)/\Sigma, z} = I^{\text{es}}(C, z) = \{g \in \mathcal{O}_{\Sigma, z} \mid f + \varepsilon g \text{ is equisingular over } \mathbb{C}[\varepsilon]/(\varepsilon^2)\},$$

where $f \in \mathcal{O}_{\Sigma, z}$ is a local equation of C at z . $I^{\text{es}}(C, z)$ is called the *equisingularity ideal* of the singularity (C, z) , and it is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point. $I^{\text{es}}(C, z)/I^{\text{ea}}(C, z)$ can be identified with the tangent space of the equisingular stratum in the semiuniversal deformation of (C, z) (cf. [Wah74b], [DiH88], and Definition III.2.1). We call $X^{\text{es}}(C)$ the *equisingularity scheme* of C .

- (c) The scheme $X_{\text{fix}}^{\text{ea}}(C)$ in Σ is defined via the ideal sheaf $\mathcal{J}_{X_{\text{fix}}^{\text{ea}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X_{\text{fix}}^{\text{ea}}(C)/\Sigma, z} = I_{\text{fix}}^{\text{ea}}(C, z) = I_{\text{fix}}^{\text{ea}}(f) = (f) + \mathfrak{m}_{\Sigma, z} \cdot \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) \subseteq \mathcal{O}_{\Sigma, z},$$

where x, y denote local coordinates of Σ at z and $f \in \mathcal{O}_{\Sigma, z}$ a local equation of C . $I_{\text{fix}}^{\text{ea}}(C, z)$ is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point of C .

- (d) We define the zero-dimensional subscheme $X_{\text{fix}}^{\text{es}}(C)$ of Σ via the ideal sheaf $\mathcal{J}_{X_{\text{fix}}^{\text{es}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X_{\text{fix}}^{\text{es}}(C)/\Sigma, z} = I_{\text{fix}}^{\text{es}}(C, z) = \left\{ g \in \mathcal{O}_{\Sigma, z} \mid \begin{array}{l} f + \varepsilon g \text{ is equisingular over } \mathbb{C}[\varepsilon]/(\varepsilon^2) \\ \text{along the trivial section} \end{array} \right\},$$

where $f \in \mathcal{O}_{\Sigma, z}$ is a local equation of C at z . $I_{\text{fix}}^{\text{es}}(C, z)$ is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point.

- (e) The scheme $X^{\text{a}}(C)$ in Σ is defined via the ideal sheaf $\mathcal{J}_{X^{\text{a}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X^{\text{a}}(C)/\Sigma, z} = I^{\text{a}}(C, z) \subseteq \mathcal{O}_{\Sigma, z},$$

where we refer for the somewhat lengthy definition of $I^{\text{a}}(C, z)$ to Remark 2.7. $I^{\text{a}}(C, z)$ is called the *analytical singularity ideal* of the singularity (C, z) , and it is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point. We call $X^{\text{a}}(C)$ the *analytical singularity scheme* of C .

- (f) The scheme $X^{\text{s}}(C)$ in Σ is defined via the ideal sheaf $\mathcal{J}_{X^{\text{s}}(C)/\Sigma}$ with stalks

$$\mathcal{J}_{X^{\text{s}}(C)/\Sigma, z} = I^{\text{s}}(C, z) = \left\{ g \in \mathcal{O}_{\Sigma, z} \mid g \text{ goes through the cluster } \mathcal{Cl}(C, T^*(C, z)) \right\},$$

where $T^*(C, z)$ denotes the essential subtree of the complete embedded resolution tree of (C, z) . $I^{\text{s}}(C, z)$ is called the *singularity ideal* of the singularity (C, z) , and it is of course $\mathcal{O}_{\Sigma, z}$ whenever z is a smooth point. We call $X^{\text{s}}(C)$ the *singularity scheme* of C . (Cf. [Los98] Section 2.2.1.)

Throughout the thesis we will frequently treat topological and analytical singularities at the same time. Whenever we do so, we will write $X^*(C)$ for $X^{\text{es}}(C)$ respectively for $X^{\text{ea}}(C)$, and similarly $X_{\text{fix}}^*(C)$ for $X_{\text{fix}}^{\text{es}}(C)$ respectively for $X_{\text{fix}}^{\text{ea}}(C)$, and $X(C)$ for $X^{\text{s}}(C)$ respectively for $X^{\text{a}}(C)$.

2.7 Remark

Let (C, z) be an isolated plane curve singularity of analytical type \mathcal{S} with representative $f \in \mathcal{O}_{\Sigma, z}$.

(a) A collection of ideals $\mathcal{I}(f) = \{I(g) \mid g \in \mathcal{O}_{\Sigma, z} : g \sim_c f\}$ is said to be *suitable*, if the ideals $I(g) \in \mathcal{I}$ satisfy the following properties:

(1) $g \in I(g)$.

(2) For a generic element in $h \in I(g)$ we have $h \sim_c g$ and $I(h) = I(g)$; more precisely, for any $d \geq 0$ the set of polynomial $h \in I(g) \cap \mathbb{C}[x, y]_{\leq d}$ such that $h \sim_c g$ and $I(h) = I(g)$ is open dense in $I(g) \cap \mathbb{C}[x, y]_{\leq d}$, where $\mathbb{C}[x, y]_{\leq d}$ are the polynomials of degree at most d .

(3) $I(g) = \psi^*I(f)$, if $\psi : \mathcal{O}_{\Sigma, z} \rightarrow \mathcal{O}_{\Sigma, z}$ is an isomorphism and $u \in \mathcal{O}_{\Sigma, z}$ is a unit such that $g = \psi^*(u \cdot f)$.

(4) $\exists m > 0 : I(g) = I(\text{jet}_m(g))$, i. e. $I(g)$ is determined by $\text{jet}_m(g)$.

We note that (3) implies that $I(g)$ only depends on the ideal $\langle g \rangle$, and that the isomorphism class of $I(f)$ is an invariant of \mathcal{S} . We call $\deg \mathcal{I}(f) = \dim_{\mathbb{C}}(\mathcal{O}_{\Sigma, z}/I(f))$ the *degree* of $\mathcal{I}(f)$. (Cf. [Los98] Definition 2.39.)

(b) If we define

$$\tilde{\mathcal{I}}^a(g) = \{h \in \mathcal{O}_{\Sigma, z} \mid I^{ea}(h) \subseteq I^{ea}(g)\}$$

for $g \in \mathcal{O}_{\Sigma, z}$ with $g \sim_c f$, then the collection $\tilde{\mathcal{I}}(f) = \{\tilde{\mathcal{I}}^a(g) \mid g \in \mathcal{O}_{\Sigma, z} : g \sim_c f\}$ is suitable. (Cf. [Los98] Definition 2.42.)

By [Los98] Lemma 2.44 we know $\deg \tilde{\mathcal{I}}(f) \leq 3\tau(\mathcal{S})$.

(c) Among the suitable collections of ideals we choose one with minimal degree, and we call it $\mathcal{I}^a(f)$. The elements of $\mathcal{I}^a(f)$ are parametrised by $G := \{g \in \mathcal{O}_{\Sigma, z} \mid g \sim_c f\}$, and we denote the element corresponding to $g \in G$ by $I^a(g)$. We then define

$$I^a(C, z) = I^a(f).$$

Note that $\mathcal{I}^a(f)$ exists in view of (b), and that its definition depends only on \mathcal{S} . Moreover, $\deg \mathcal{I}^a(f) \leq 3\tau(\mathcal{S})$. (Cf. [Los98] Definition 2.40.)

2.8 Remark

If $C \subset \Sigma$ is a reduced curve, then $\mathcal{J}_{X^a(C)/\Sigma} \subseteq \mathcal{J}_{X_{\text{fix}}^{ea}(C)/\Sigma} \subseteq \mathcal{J}_{X^{ea}(C)/\Sigma}$ and $\mathcal{J}_{X^s(C)/\Sigma} \subseteq \mathcal{J}_{X_{\text{fix}}^{es}(C)/\Sigma} \subseteq \mathcal{J}_{X^{es}(C)/\Sigma}$.

In particular, the vanishing of $h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D))$ implies the vanishing of $h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D))$, and the vanishing of the latter implies that of $h^1(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(D))$.

To see the first assertion, consider the exact sequence

$$0 \rightarrow \mathcal{J}_{X(C)/\Sigma}(D) \rightarrow \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D) \rightarrow (\mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}/\mathcal{J}_{X(C)/\Sigma})(D) \rightarrow 0.$$

The long exact cohomology sequence then leads to

$$H^1(\Sigma, \mathcal{J}_{X/\Sigma}(D)) \rightarrow H^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*/\Sigma}(D)) \rightarrow H^1(\Sigma, (\mathcal{J}_{X_{\text{fix}}^*/\Sigma}/\mathcal{J}_{X/\Sigma})(D)) = 0,$$

where the latter vanishes since its support is zero-dimensional. This proves the claim. The second assertion follows analogously.

2.9 Remark

In [Los98] Proposition 2.19 and 2.20 and in Remark 2.40 (see also [GLS00] and 2.41) it is shown that, fixing a point $z \in \Sigma$ and a topological respectively analytical type \mathcal{S} , the singularity schemes respectively analytical singularity schemes having the same topological respectively analytical type are parametrised by an irreducible Hilbert scheme, which we are going to denote by $\text{Hilb}_z(\mathcal{S})$. This then leads to an irreducible family

$$\text{Hilb}(\mathcal{S}) = \coprod_{z \in \Sigma} \text{Hilb}_z(\mathcal{S}).$$

For a more careful study of these objects we refer to [Los98] Remark 2.21 and Remark 2.41.

In particular, equisingular respectively equianalytical singularities have singularity schemes respectively analytical singularity schemes of the same degree (see also [GLS98c] or [Los98] Lemma 2.8). The same is of course true, regarding the equisingularity scheme respectively the equianalytical singularity scheme. This leads to the following definition.

2.10 Definition

If $C \subset \Sigma$ is a reduced curve such that z is a singular point of topological respectively analytical type \mathcal{S} , then we define $\deg(X^s(\mathcal{S})) = \deg(X^s(C), z)$, $\deg(X_{\text{fix}}^{\text{es}}(\mathcal{S})) = \deg(X_{\text{fix}}^{\text{es}}(C), z)$ and $\deg(X^{\text{es}}(\mathcal{S})) = \deg(X^{\text{es}}(C), z)$ respectively $\deg(X^a(\mathcal{S})) = \deg(X^a(C), z)$, $\deg(X_{\text{fix}}^{\text{ea}}(\mathcal{S})) = \deg(X_{\text{fix}}^{\text{ea}}(C), z)$ and $\deg(X^{\text{ea}}(\mathcal{S})) = \deg(X^{\text{ea}}(C), z)$.

To express the degree of the schemes $X^*(\mathcal{S})$ and $X_{\text{fix}}^*(\mathcal{S})$ in terms of the invariants from Remark 2.2 for an analytical respectively topological singularity type \mathcal{S} is much simpler, since by definition

$$\deg(X^{\text{es}}(\mathcal{S})) = \tau^{\text{es}}(\mathcal{S}) \quad \text{and} \quad \deg(X^{\text{ea}}(\mathcal{S})) = \tau(\mathcal{S}),$$

while $\deg(X_{\text{fix}}^*(\mathcal{S})) = \deg(X^*(\mathcal{S})) + 2$, and hence

$$\deg(X_{\text{fix}}^{\text{es}}(\mathcal{S})) = \tau^{\text{es}}(\mathcal{S}) + 2 \quad \text{and} \quad \deg(X_{\text{fix}}^{\text{ea}}(\mathcal{S})) = \tau(\mathcal{S}) + 2.$$

2.11 Remark

We note that for a topological respectively analytical singularity type \mathcal{S}

$$\dim \text{Hilb}_z(\mathcal{S}) = \deg(X(\mathcal{S})) - \deg(X^*(\mathcal{S})) - 2$$

for any $z \in \Sigma$, and thus

$$\dim \text{Hilb}(\mathcal{S}) = \deg(X(\mathcal{S})) - \deg(X^*(\mathcal{S})).$$

For the convenience of the reader we reproduce the proof from [GLS02] Lemma 2.1.49 at the end of this chapter.

2.12 Remark

- (a) A simple calculation shows (see also [Los98] p. 28)

\mathcal{S}	$\deg(X^a(\mathcal{S})) = \deg(X^s(\mathcal{S}))$	
A_μ	$\frac{3}{2} \cdot \mu + 2,$	μ even,
A_μ	$\frac{3}{2} \cdot \mu + \frac{3}{2},$	μ odd,
D_μ	$\frac{3}{2} \cdot \mu,$	μ even,
D_μ	$\frac{3}{2} \cdot \mu + \frac{1}{2},$	$\mu \neq 5$ odd,
D_5	$\frac{3}{2} \cdot \mu - \frac{1}{2} = 7,$	$\mu = 5,$
E_6	$\frac{3}{2} \cdot \mu - 1 = 8,$	$\mu = 6,$
E_7	$\frac{3}{2} \cdot \mu - \frac{1}{2} = 10,$	$\mu = 7,$
E_6	$\frac{3}{2} \cdot \mu - 1 = 11,$	$\mu = 8.$

In particular, if \mathcal{S} is a simple singularity, then $\deg(X^s(\mathcal{S})) \leq \frac{3}{2}\mu(\mathcal{S}) + 2$, where $\mu(\mathcal{S})$ denotes the Milnor number of \mathcal{S} .

- (b) If
- \mathcal{S}
- is an analytical singularity type, then (cf. [Los98] Lemma 2.44)

$$\deg(X^a(\mathcal{S})) \leq 3\tau(\mathcal{S}), \quad (2.1)$$

where $\tau(\mathcal{S})$ denotes the Tjurina number of \mathcal{S} .

- (c) Let
- \mathcal{S}
- be any topological singularity type, then by [Los98] Lemma 2.8

$$\deg(X^s(\mathcal{S})) = \delta(\mathcal{S}) + \sum_{q \in T^*(C,z)} m_q,$$

where (C, z) is a plane curve singularity of type \mathcal{S} , $T^*(C, z)$ is the essential subtree of the complete embedded resolution tree of (C, z) , m_q is the multiplicity of (C, z) at the infinitely near point $q \in T^*(C, z)$, and $\delta(\mathcal{S})$ the delta invariant of \mathcal{S} .

In [Los98] p. 103 it is shown that if \mathcal{S} is not a simple singularity type, then

$$\deg(X^s(\mathcal{S})) \leq \frac{3}{2} \cdot (\mu(\mathcal{S}) - 1).$$

Combining this with (a) we get for \mathcal{S} arbitrary

$$\deg(X^s(\mathcal{S})) \leq \frac{3}{2}\mu(\mathcal{S}) + 2. \quad (2.2)$$

2.13 Definition

- (a) Given topological or analytical singularity types
- $\mathcal{S}_1, \dots, \mathcal{S}_r$
- and a divisor
- $D \in \text{Div}(\Sigma)$
- , we denote by
- $V = V_{|D|}(\mathcal{S}_1, \dots, \mathcal{S}_r)$
- the locally closed subspace of
- $|D|_{\mathbb{1}}$
- of reduced curves in the linear system
- $|D|_{\mathbb{1}}$
- having precisely
- r
- singular points of types
- $\mathcal{S}_1, \dots, \mathcal{S}_r$
- .

By $V^{\text{reg}} = V_{|\mathcal{D}|}^{\text{reg}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ we denote the open³ subset

$$V^{\text{reg}} = \{C \in V \mid h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) = 0\} \subseteq V,$$

where $X(C) = X^s(C)$ respectively $X^a(C)$, and by $V^{\text{irr}} = V_{|\mathcal{D}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ we denote the open subset

$$V^{\text{irr}} = \{C \in V \mid h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D)) = 0\} \subseteq V,$$

where $X_{\text{fix}}^*(C) = X_{\text{fix}}^{\text{es}}(C)$ respectively $X_{\text{fix}}^{\text{ea}}(C)$

Similarly, we use the notation $V^{\text{irr}} = V_{|\mathcal{D}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ to denote the open subset of irreducible curves in the space V , and we set $V^{\text{irr,reg}} = V_{|\mathcal{D}|}^{\text{irr,reg}}(\mathcal{S}_1, \dots, \mathcal{S}_r) = V^{\text{irr}} \cap V^{\text{reg}}$ and $V^{\text{irr,fix}} = V_{|\mathcal{D}|}^{\text{irr,fix}}(\mathcal{S}_1, \dots, \mathcal{S}_r) = V^{\text{irr}} \cap V^{\text{fix}}$, which are open in V^{reg} respectively V^{fix} , and hence in V .

If a type \mathcal{S} occurs $k > 1$ times, we rather write $k\mathcal{S}$ than $\mathcal{S}, \dots, \mathcal{S}$.

- (b) Analogously, $V_{|\mathcal{D}|}(m_1, \dots, m_r) = V_{|\mathcal{D}|}(\underline{m})$ denotes the locally closed subspace of $|\mathcal{D}|_1$ of reduced curves having precisely r ordinary singular points of multiplicities m_1, \dots, m_r . (Cf. [GrS99] or [Los98] 1.3.2)
- (c) Let $V = V_{|\mathcal{D}|}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ respectively $V = V_{|\mathcal{D}|}(\underline{m})$. We say V is *T-smooth* at $C \in V$ if the germ (V, C) is smooth of the (expected) dimension $\dim |\mathcal{D}|_1 - \deg(X^*(C))$, where $X^*(C) = X^{\text{es}}(C)$, $X^*(C) = X^{\text{es}}(C)$ or $X^*(C) = X(\underline{m}; \underline{z})$ with $\text{Sing}(C) = \{z_1, \dots, z_r\}$ respectively.

We call these families of curves *equisingular families of curves*.

2.14 Remark

With the notation of Definition 2.13 and by [Los98] Proposition 2.1 (see also [GrK89], [GrL96], [GLS00]) T-smoothness of V at C is implied by the vanishing of $H^1(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(C))$. This is due to the fact that the tangent space of V at C may be identified with $H^0(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(C))/H^0(\Sigma, \mathcal{O}_\Sigma)$.

In particular, the subvarieties V^{reg} , V^{fix} , $V^{\text{irr,reg}}$ and $V^{\text{irr,fix}}$ are T-smooth, that is smooth of expected dimension $h^0(\Sigma, \mathcal{O}_\Sigma(C)) - \deg(X^*(C)) - 1$.

2.15 Definition

Let $D \in \text{Div}(\Sigma)$ be a divisor, $\mathcal{S}_1, \dots, \mathcal{S}_r$ distinct topological or analytical singularity types, and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

- (a) We denote by \tilde{B} the irreducible parameter space

$$\tilde{B} = \tilde{B}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) = \prod_{i=1}^r \text{Sym}^{k_i}(\text{Hilb}(\mathcal{S}_i)),$$

and by $B = B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ the non-empty open, irreducible and dense, subspace

$$B = \left\{ ([X_{1,1}, \dots, X_{1,k_1}], \dots, [X_{r,1}, \dots, X_{r,k_r}]) \in \tilde{B} \mid \begin{array}{l} \text{supp}(X_{i,j}) \cap \text{supp}(X_{s,t}) = \emptyset \\ \forall 1 \leq i, s \leq r, 1 \leq j \leq k_i, 1 \leq t \leq k_s \end{array} \right\}.$$

³See Theorem V.1.1.

Note that $\dim(B)$ does not depend on Σ , more precisely, with the notation of Remark 2.11 we have

$$\dim(B) = \sum_{i=1}^r k_i \cdot \left(\deg(X(\mathcal{S}_i)) - \deg(X^*(\mathcal{S}_i)) \right).$$

- (b) Let us set $n = \sum_{i=1}^r k_i \deg(X(\mathcal{S}_i))$. We then define an injective morphism

$$\begin{aligned} \psi = \psi(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) : B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) &\longrightarrow \text{Hilb}_\Sigma^n \\ ([X_{1,1}, \dots, X_{1,k_1}], \dots, [X_{r,1}, \dots, X_{r,k_r}]) &\longmapsto \bigcup_{i=1}^r \bigcup_{j=1}^{k_i} X_{i,j}, \end{aligned}$$

where Hilb_Σ^n denotes the smooth connected Hilbert scheme of zero-dimensional schemes of degree n on Σ (cf. [Los98] Section 1.3.1).

- (c) We denote by $\Psi = \Psi_D(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ the fibration of $V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ induced by $B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$, that is, the morphism Ψ is given by

$$\begin{aligned} \Psi : V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) &\longrightarrow B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) \\ C &\longmapsto ([X_{1,1}, \dots, X_{1,k_1}], \dots, [X_{r,1}, \dots, X_{r,k_r}]) \end{aligned}$$

where $\text{Sing}(C) = \{z_{i,j} \mid i = 1, \dots, r, j = 1, \dots, k_i\}$, $X_{i,j} = X(C, z_{i,j})$ and $(C, z_{i,j}) \cong \mathcal{S}_i$ for all $i = 1, \dots, r, j = 1, \dots, k_i$.

- (d) Denoting by $m = k_1 + \dots + k_r$ the number of imposed singularities we define the fibration $\Phi = \Phi_D(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ by

$$\begin{aligned} \Phi : V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) &\longrightarrow \text{Sym}^m(\Sigma) \\ C &\longmapsto \text{Sing}(C), \end{aligned}$$

sending a curve C to the unordered tuple of its singular points.

2.16 Remark

- (a) With the notation of Definition 2.13 and Definition 2.15 note that for $C \in V = V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ the fibre $\Psi^{-1}(\Psi(C))$ is the open dense subset of the linear system $|\mathcal{J}_{X(C)/\Sigma}(D)|_1$ consisting of the curves C' with $X(C') = X(C)$. In particular, the fibres of Ψ restricted to V^{reg} are irreducible, and since for $C \in V^{\text{reg}}$ the cohomology group $H^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D))$ vanishes, they are equidimensional of dimension

$$\begin{aligned} h^0(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) - 1 &= h^0(\Sigma, \mathcal{O}_\Sigma(D)) - \deg(X(C)) - 1 \\ &= h^0(\Sigma, \mathcal{O}_\Sigma(D)) - \sum_{i=1}^r k_i \cdot \deg(X(\mathcal{S}_i)) - 1. \end{aligned}$$

- (b) Note that for $C \in V$ the fibre $\Phi^{-1}(\Phi(C))$ has at C the tangent space $H^0(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)}(D))/H^0(\mathcal{O}_\Sigma)$, so that

$$\dim(\Phi^{-1}(\Phi(C))) \leq h^0(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)}(D)) - 1. \quad (2.3)$$

Moreover, suppose that $h^1(\Sigma, \mathcal{J}_{\mathcal{X}_{\text{fix}}^*(C)/\Sigma}(D)) = 0$, then the germ of the fibration at C

$$(\Phi, C) : (V, C) \rightarrow (\text{Sym}^m(\Sigma), \text{Sing}(C))$$

is smooth of fibre dimension $h^0(\Sigma, \mathcal{J}_{\mathcal{X}_{\text{fix}}^*(C)}(D)) - 1$, i. e. locally at C the morphism Φ is a projection of the product of the smooth base space with the smooth fibre. This implies in particular, that close to C there is a curve having its singularities in very general position. (Cf. [Los98] Proposition 2.1 (e).)

Proof of Remark 2.11: Since $\dim \text{Hilb}(\mathcal{S})$ is independent of Σ , we may as well suppose that $\Sigma = \mathbb{P}_c^2$, and we let H be a line in \mathbb{P}_c^2 .

For a reduced curve $C \subset \mathbb{P}_c^2$ we set $X(C) = X^s(C)$ and $X^*(C) = X^{\text{es}}(C)$ respectively $X(C) = X^a(C)$ and $X^*(C) = X^{\text{ea}}(C)$.

Since $\deg(X) = \deg(X(\mathcal{S}))$ is independent of $X \in \text{Hilb}(\mathcal{S})$, there is an integer $m > 0$ such that for $d > m$

$$h^1(\mathbb{P}_c^2, \mathcal{J}_{X/\mathbb{P}_c^2}(d)) = 0$$

for any $X \in \text{Hilb}(\mathcal{S})$.

Let $k > 0$ be the determinacy bound of \mathcal{S} , that is, any representative $f \in \mathcal{O}_{\mathbb{P}_c^2, z} = \mathbb{C}\{x, y\}$ of \mathcal{S} depends only on the k -jet of f . Hence, for $d > k$ the morphism

$$\Psi = \Psi_{\text{dH}}(\mathcal{S}) : V_{|\text{dH}|}(\mathcal{S}) \rightarrow B(\mathcal{S}) = \text{Hilb}(\mathcal{S})$$

is surjective.

Let us now fix some $d > \max\{k, m\}$. For each $C \in V_{|\text{dH}|}(\mathcal{S})$ the fibre $\Psi^{-1}(\Psi(C))$ is the open dense subset of $|\mathcal{J}_{X(C)/\mathbb{P}_c^2}(d)|$, consisting of curves $C' \in V_{|\text{dH}|}(\mathcal{S})$ with $X(C') = X(C)$. From the long exact cohomology sequence of

$$0 \rightarrow \mathcal{J}_{X(C)/\Sigma} \rightarrow \mathcal{O}_{\mathbb{P}_c^2}(d) \rightarrow \mathcal{O}_{X(C)} \rightarrow 0$$

it follows

$$h^0(\mathbb{P}_c^2, \mathcal{J}_{X(C)/\mathbb{P}_c^2}(d)) = h^0(\mathbb{P}_c^2, \mathcal{O}_{\mathbb{P}_c^2}(d)) - \deg(X(\mathcal{S})).$$

In particular the fibres all have the same dimension

$$\dim \Psi^{-1}(\Psi(C)) = h^0(\mathbb{P}_c^2, \mathcal{O}_{\mathbb{P}_c^2}(d)) - \deg(X(\mathcal{S})) - 1.$$

We therefore get

$$\begin{aligned} \dim \text{Hilb}(\mathcal{S}) &= \dim \Psi(V_{|\text{dH}|}(\mathcal{S})) = \dim(V_{|\text{dH}|}(\mathcal{S})) - \dim \Psi^{-1}(\Psi(C)) \\ &= \dim(V_{|\text{dH}|}(\mathcal{S})) - h^0(\mathbb{P}_c^2, \mathcal{O}_{\mathbb{P}_c^2}(d)) + \deg(X(\mathcal{S})) + 1. \end{aligned}$$

Moreover, by Remark 2.8 we know that also $h^1(\mathbb{P}_c^2, \mathcal{J}_{X^*(C)/\mathbb{P}_c^2}(d)) = 0$ for any $C \in V_{|\text{dH}|}(\mathcal{S})$, and thus in view of Remark 2.14 $V_{|\text{dH}|}(\mathcal{S})$ is \mathbf{T} -smooth, that is

$$\dim(V_{|\text{dH}|}(\mathcal{S})) = h^0(\mathbb{P}_c^2, \mathcal{O}_{\mathbb{P}_c^2}(d)) - \deg(X^*(\mathcal{S})) - 1,$$

which finishes the claim. \square

CHAPTER II

A Vanishing Theorem

In Chapter III we deduce the existence of an irreducible curve with prescribed topological or analytical singularities from the existence of an irreducible curve C with ordinary multiple points z_1, \dots, z_r of certain multiplicities m_1, \dots, m_r which has the T-smoothness property, that is such that $h^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0$. If we want to derive numerical conditions for the existence in terms of invariants of the singularities and the divisor D from this theorem, we need to have such numerical conditions for the vanishing of $H^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D))$. Section 1 is devoted to the proof of a suitable vanishing theorem. We generalise an approach used by Geng Xu in [Xu95] for the plane case. The basic idea is to give conditions such that $(\pi^*D - \sum_{i=1}^r m_i E_i) - K_{\tilde{\Sigma}}$ is big and nef, where $\pi : \text{Bl}_{\underline{z}}(\Sigma) = \tilde{\Sigma} \rightarrow \Sigma$ denotes the blow up of Σ in z_1, \dots, z_r , and then to apply the Kawamata–Viehweg Vanishing Theorem.

1. The Vanishing Theorem

1.1 Theorem

Let $m_1 \geq \dots \geq m_r \geq 0$ be non-negative integers, $\alpha \in \mathbb{R}$ with $\alpha > 1$, $k_\alpha = \max \{n \in \mathbb{N} \mid n < \frac{\alpha}{\alpha-1}\}$ and let $D \in \text{Div}(\Sigma)$ be a divisor satisfying the following three conditions¹

$$(1.1) \quad (D - K_\Sigma)^2 \geq \max \left\{ \alpha \cdot \sum_{i=1}^r (m_i + 1)^2, (k_\alpha \cdot m_1 + k_\alpha)^2 \right\},$$

$$(1.2) \quad (D - K_\Sigma) \cdot B \geq k_\alpha \cdot (m_1 + 1) \text{ for any irreducible curve } B \text{ with } B^2 = 0 \text{ and } \dim |B|_a > 0, \text{ and}$$

$$(1.3) \quad D - K_\Sigma \text{ is nef.}$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), \pi^*D - \sum_{i=1}^r m_i E_i \right) = 0.$$

In particular,

$$H^\nu(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0.$$

Proof: By the Kawamata–Viehweg Vanishing Theorem (cf. [Kaw82] and [Vie82]) it suffices to show that $A = (\pi^*D - \sum_{i=1}^r m_i E_i) - K_{\tilde{\Sigma}}$ is big and nef, i. e. we have to show:

¹The proof uses Kawamata–Viehweg vanishing which needs characteristic zero for the ground field.

- (a) $A^2 > 0$, and
 (b) $A \cdot B' \geq 0$ for any irreducible curve B' in $\tilde{\Sigma} = \text{Bl}_{\underline{z}}(\Sigma)$.

Note that $A = \pi^*(D - K_{\Sigma}) - \sum_{i=1}^r (m_i + 1)E_i$, and thus by Hypothesis (1.4) we have

$$A^2 = (D - K_{\Sigma})^2 - \sum_{i=1}^r (m_i + 1)^2 > 0,$$

which gives condition (a).

For condition (b) we observe that an irreducible curve B' on $\tilde{\Sigma}$ is either the strict transform of an irreducible curve B in Σ or is one of the exceptional curves E_i . In the latter case we have

$$A \cdot B' = A \cdot E_i = m_i + 1 > 0.$$

We may, therefore, assume that $B' = \tilde{B}$ is the strict transform of an irreducible curve B on Σ having multiplicity $\text{mult}_{z_i}(B) = n_i$ at z_i , $i = 1, \dots, r$. Then

$$A \cdot B' = (D - K_{\Sigma}) \cdot B - \sum_{i=1}^r (m_i + 1)n_i,$$

and thus condition (b) is equivalent to

$$(b') \quad (D - K_{\Sigma}) \cdot B \geq \sum_{i=1}^r (m_i + 1)n_i.$$

Since \underline{z} is in very general position Lemma 2.1 applies in view of Corollary A.3. Using the Hodge Index Theorem E.4, Hypothesis (1.4), Lemma 2.1, and the Cauchy-Schwarz Inequality we get the following sequence of inequalities:

$$\begin{aligned} ((D - K_{\Sigma}) \cdot B)^2 &\geq (D - K_{\Sigma})^2 \cdot B^2 \geq \alpha \cdot \left(\sum_{i=1}^r (m_i + 1)^2 \right) \cdot \left(\sum_{i=1}^r n_i^2 - n_{i_0} \right) \\ &= \sum_{i=1}^r (m_i + 1)^2 \cdot \sum_{i=1}^r n_i^2 + (\alpha - 1) \cdot \left(\sum_{i=1}^r (m_i + 1)^2 \cdot \left(\sum_{i=1}^r n_i^2 - \frac{\alpha}{\alpha-1} \cdot n_{i_0} \right) \right) \\ &\geq \left(\sum_{i=1}^r (m_i + 1) \cdot n_i \right)^2 + (\alpha - 1) \cdot \left(\sum_{i=1}^r (m_i + 1)^2 \cdot \left(\sum_{i=1}^r n_i^2 - \frac{\alpha}{\alpha-1} \cdot n_{i_0} \right) \right), \end{aligned}$$

where $i_0 \in \{1, \dots, r\}$ is such that $n_{i_0} = \min\{n_i \mid n_i \neq 0\}$. Since $D - K_{\Sigma}$ is nef, condition (b') is satisfied as soon as we have

$$\sum_{i=1}^r n_i^2 \geq \frac{\alpha}{\alpha-1} \cdot n_{i_0}.$$

If this is not fulfilled, then $n_i < \frac{\alpha}{\alpha-1}$ for all $i = 1, \dots, r$, and thus

$$\sum_{i=1}^r (m_i + 1) \cdot n_i \leq k_{\alpha} \cdot (m_1 + 1).$$

Hence, for the remaining considerations (b') may be replaced by the worst case

$$(D - K_{\Sigma}) \cdot B \geq k_{\alpha} \cdot (m_1 + 1).$$

Note that since the z_i are in very general position and $z_{i_0} \in B$ we have that $B^2 \geq 0$ and $\dim |B|_a > 0$ (cf. Corollary A.6). If $B^2 > 0$ then we are done by the Hodge Index Theorem E.4 and Hypothesis (1.4), since $D - K_\Sigma$ is nef:

$$(D - K_\Sigma).B \geq \sqrt{(D - K_\Sigma)^2} \geq \sqrt{(k_\alpha \cdot m_1 + k_\alpha)^2} \geq k_\alpha \cdot (m_1 + 1).$$

It remains to consider the case $B^2 = 0$ which is covered by Hypothesis (1.5).

For the “in particular” part we just note that $\mathcal{J}_{X(m_1; z_1)/\Sigma} \otimes \cdots \otimes \mathcal{J}_{X(m_r; z_r)/\Sigma} \otimes \mathcal{O}_\Sigma(D) \cong \mathcal{J}_{X(\underline{m}; z)/\Sigma} \otimes \mathcal{O}_\Sigma(D)$ (see Lemma C.4) and that, using the Leray spectral sequence (compare [Laz97] Lemma 5.1)

$$H^\nu \left(\Sigma, \bigotimes_{i=1}^r \mathcal{J}_{X(m_i; z_i)/\Sigma} \otimes \mathcal{O}_\Sigma(D) \right) = H^\nu \left(\tilde{\Sigma}, \pi^* D - \sum_{i=1}^r m_i E_i \right).$$

□

Choosing the constant $\alpha = 2$ in Theorem 1.1, then $\frac{\alpha}{\alpha-1} = 2$ and thus $k_\alpha = 1$. We therefore get the following corollary, which has the advantage that the conditions look simpler, and that the hypotheses on the “exceptional” curves are not too hard.

1.2 Corollary

Let $m_1, \dots, m_r \in \mathbb{N}_0$, and $D \in \text{Div}(\Sigma)$ be a divisor satisfying the following three conditions

$$(1.4) \quad (D - K_\Sigma)^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.5) \quad (D - K_\Sigma).B > \max\{m_i \mid i = 1, \dots, r\} \text{ for any irreducible curve } B \text{ with } B^2 = 0 \text{ and } \dim |B|_a > 0, \text{ and}$$

$$(1.6) \quad D - K_\Sigma \text{ is nef.}$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), \pi^* D - \sum_{i=1}^r m_i E_i \right) = 0.$$

In particular,

$$H^\nu(\Sigma, \mathcal{J}_{X(\underline{m}; z)/\Sigma}(D)) = 0.$$

1.3 Remark

Condition (1.3) respectively Condition (1.6) are in several respects “expectable”. First, Theorem 1.1 is a corollary of the Kawamata–Viehweg Vanishing Theorem, and if we take all m_i to be zero, our assumptions should basically be the same, i. e. $D - K_\Sigma$ nef and big. The latter is more or less just (1.1) respectively (1.4). Secondly, we want to apply the theorem to an existence problem. A divisor being nef means it is somehow close to being effective, or better its linear system is close to being non-empty. If we want that some linear system $|D|_l$ contains a curve with certain properties, then it seems not to be so unreasonable to restrict to systems where already $|D - K_\Sigma|_l$,

or even $|D - L - K_\Sigma|_1$ with L some fixed divisor, is of positive dimension, thus nef.

In many interesting examples, such as $\mathbb{P}_\mathbb{C}^2$, Condition (1.2) respectively (1.5) turn out to be obsolete or easy to handle. So finally the most restrictive obstruction seems to be (1.1) respectively (1.4).

If we consider the situation where the largest multiplicity m_1 occurs in a large number, more precisely, if $m_1 = \dots = m_{l_\alpha}$ with $l_\alpha = \min \{n \in \mathbb{N} \mid \alpha \cdot n \geq k_\alpha^2\}$, then Condition (1.1) comes down to

$$(1.1') \quad (D - K_\Sigma)^2 \geq \alpha \cdot \sum_{i=1}^r (m_i + 1)^2.$$

1.4 Remark

Even though we said that condition (1.4) was the really restrictive condition we would like to understand better what condition (1.5) means. We therefore show in Appendix B that an algebraic system $|B|_a$ of dimension greater than zero with B irreducible and $B^2 = 0$ gives rise to a fibration $f : \Sigma \rightarrow H$ of Σ over a smooth projective curve H whose fibres are just the elements of $|B|_a$.

2. Generalisation of a Lemma of Geng Xu

Throughout the proof of Theorem 1.1 we need the following generalisation of a lemma of Geng Xu.

2.1 Lemma

Let $\underline{z} = (z_1, \dots, z_r) \in \Sigma^r$ be in very general position, $\underline{n} \in \mathbb{N}_0^r$, and let $B \subset \Sigma$ be an irreducible curve with $\text{mult}_{z_i}(B) \geq n_i$, then

$$B^2 \geq \sum_{i=1}^r n_i^2 - \min\{n_i \mid n_i \neq 0\}.$$

2.2 Remark

- (a) A proof for the above lemma in the case $\Sigma = \mathbb{P}_\mathbb{C}^2$ can be found in [Xu94] and in the case $r = 1$ in [EiL93]. Here we just extend the arguments given there to the slightly more general situation.
- (b) For better estimates of the self intersection number of curves in the situation where one has some knowledge on equisingular deformations inside the algebraic system see [GuS84].
- (c) With the notation of Lemma A.2 respectively Corollary A.3 the assumption in Lemma 2.1 could be formulated more precisely as “let $B \subset \Sigma \subseteq \mathbb{P}_\mathbb{C}^N$ be an irreducible curve such that $V_{B, \underline{n}} = \Sigma^r$ ”, or “let $\underline{z} \in \Sigma^r \setminus V$ ”.²

²Since B is irreducible, the general element in $|B|_a$ will be irreducible. Since $V_{B, \underline{n}} = \Sigma^r$ there will be some family of curves in Hilb_Σ^h satisfying the requirements of Lemma 2.3.

- (d) Note, that one cannot expect to get rid of the “ $-\min\{n_i \mid n_i \neq 0\}$ ”. E. g. $\Sigma = \text{Bl}_z(\mathbb{P}_C^2)$, the projective plane blown up in a point z , and $B \subset \Sigma$ the strict transform of a line through z . Let now $r = 1$, $n_1 = 1$ and $z_1 \in \Sigma$ be any point. Then there is of course a (unique) curve $B_1 \in |\mathcal{B}|_a$ through z_1 , but $B^2 = 0 < 1 = n_1^2$.

Idea of the proof: Set $e_1 := n_1 - 1$ and $e_i := n_i$ for $i \neq 1$, where w. l. o. g. $n_1 = \min\{n_i \mid n_i \neq 0\}$. By assumption there is a family $\{C_t\}_{t \in \mathbb{C}}$ in $|\mathcal{B}|_a$ satisfying the requirements of Lemma 2.3. Setting $C := C_0$ the proof is done in three steps:

Step 1: We show that $H^0(C, \mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C)) \neq 0$. (Lemma 2.3)

Step 2: We deduce that $H^0(C, \pi_* \mathcal{O}_{\tilde{C}}(-\sum_{i=1}^r e_i E_i) \otimes \mathcal{O}_C(C)) \neq 0$. (Lemma 2.4)

Step 3: It follows that $\deg(\pi_* \mathcal{O}_{\tilde{C}}(-\sum_{i=1}^r e_i E_i) \otimes \mathcal{O}_C(C)) \geq 0$, but this degree is just $C^2 - \sum_{i=1}^r e_i n_i$.

□

2.3 Lemma

Given $e_1, \dots, e_r \in \mathbb{N}_0$, $r \geq 1$. Let $\{C_t\}_{t \in \mathbb{U}}$, $\mathbb{U} \subseteq \mathbb{C}$ an open neighbourhood of 0, be a non-trivial family of curves in Σ together with a section $\mathbb{U} \rightarrow \Sigma : t \mapsto z_{1,t} \in C_t$ such that

$$\text{mult}_{z_{1,t}}(C_t) \geq e_1 + 1 \quad \text{for all } t \in \mathbb{U}$$

and fixed points $z_2, \dots, z_r \in \Sigma$ such that

$$\text{mult}_{z_i}(C_t) \geq e_i \quad \text{for all } i = 2, \dots, r \text{ and } t \in \mathbb{U}.$$

Then with $z_1 = z_{1,0}$

$$H^0(C, \mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C)) \neq 0,$$

i. e. there is a non-trivial section of the normal bundle of C , vanishing at z_i to the order of at least e_i for $i = 1, \dots, r$.³

Proof: We stick to the convention $n_1 = e_1 + 1$ and $n_i = e_i$ for $i = 2, \dots, r$, and we set $z_{i,t} := z_i$ for $i = 2, \dots, r$ and $t \in \mathbb{U}$. Let $\Delta \subset \mathbb{U}$ be a small disc around 0 with coordinate t , and choose coordinates (x_i, y_i) on Σ around z_i such that

- $z_{i,t} = (a_i(t), b_i(t))$ for $t \in \Delta$ with $a_i, b_i \in \mathbb{C}\{t\}$,
- $z_i = (a_i(0), b_i(0)) = (0, 0)$, and
- $F_i(x_i, y_i, t) = f_{i,t}(x_i, y_i) \in \mathbb{C}\{x_i, y_i, t\}$, where $C_t = \{f_{i,t} = 0\}$ locally at $z_{i,t}$ (for $t \in \Delta$).

We view $\{C_t\}_{t \in \Delta}$ as a non-trivial deformation of C , which implies that the image of $\frac{\partial}{\partial t}|_{t=0} \in T_0(\Delta)$ under the Kodaira-Spencer map is a non-zero section s of $H^0(C, \mathcal{O}_C(C))$. s is locally at z_i given by $\frac{\partial F_i}{\partial t}|_{t=0}$.

Idea: Show that $\frac{\partial F_i}{\partial t}|_{t=0} \in (x_i, y_i)^{e_i}$, which are the stalks of $\mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C)$ at the z_i , and hence s is actually a global section of the subsheaf $\mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C)$.

³Note, that $\mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C) = m_{\Sigma, z_1}^{e_1} \cdots m_{\Sigma, z_r}^{e_r} \cdot \mathcal{O}_C(C)$.

Set $\Phi_{i,t}(x_i, y_i) := F_{i,t}(x_i + a_i(t), y_i + b_i(t), t) = \sum_{k=0}^{\infty} \varphi_{i,k}(x_i, y_i) \cdot t^k \in \mathbb{C}\{x_i, y_i, t\}$. By assumption for any $t \in \Delta$ the multiplicity of $\Phi_{i,t}$ at $(0, 0)$ is at least n_i , i. e. $\Phi_{i,t}(x_i, y_i) \in (x_i, y_i)^{n_i}$ for every fixed complex number $t \in \Delta$. Hence, $\varphi_{i,k}(x_i, y_i) \in (x_i, y_i)^{n_i}$ for every k .⁴

On the other hand we have

$$\begin{aligned} \varphi_{i,1}(x_i, y_i) &= \frac{\partial \Phi_{i,t}(x_i, y_i)}{\partial t} \Big|_{t=0} \\ &= \left\langle \left(\frac{\partial F_i}{\partial x_i}(x_i, y_i, 0), \frac{\partial F_i}{\partial y_i}(x_i, y_i, 0), \frac{\partial F_i}{\partial t}(x_i, y_i, 0) \right), (\dot{a}_i(0), \dot{b}_i(0), 1) \right\rangle \\ &= \frac{\partial f_{i,0}}{\partial x_i}(x_i, y_i) \cdot \dot{a}_i(0) + \frac{\partial f_{i,0}}{\partial y_i}(x_i, y_i) \cdot \dot{b}_i(0) + \frac{\partial F_i}{\partial t}(x_i, y_i, 0). \end{aligned}$$

Since $f_{i,0} \in (x_i, y_i)^{n_i}$, we have $\frac{\partial f_{i,0}}{\partial x_i}(x_i, y_i), \frac{\partial f_{i,0}}{\partial y_i}(x_i, y_i) \in (x_i, y_i)^{n_i-1}$, and hence $\frac{\partial F_i}{\partial t}(x_i, y_i, 0) \in (x_i, y_i)^{e_i}$. For this note that $\dot{a}_i(0) = \dot{b}_i(0) = 0$, if $i \neq 1$. \square

The main idea of the next lemma is the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker}(\beta) & \longrightarrow & H^0(\Sigma, \pi_* \mathcal{O}_{\tilde{\Sigma}}(-\sum_{i=1}^r e_i E_i) \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C)) & \xrightarrow{\beta} & H^0(C, \mathcal{J}_{X(\underline{e}; \underline{z})/\Sigma} \cdot \mathcal{O}_C(C)) \longrightarrow 0 \\ & & & & \downarrow \alpha & \dashrightarrow \bar{\beta} & \nearrow \gamma \\ & & & & H^0(\Sigma, \pi_* \mathcal{O}_{\tilde{C}}(-\sum_{i=1}^r e_i E_i) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)) & & \end{array}$$

i. e. the fact that $\text{Ker}(\alpha) \subseteq \text{Ker}(\beta)$, or in other words, that β factorises over α .

2.4 Lemma

Given $e_1, \dots, e_r \in \mathbb{N}_0$ and $z_1, \dots, z_r \in \Sigma$, $r \geq 1$.

The canonical morphism⁵ $\mathcal{J}_{X(e_1; z_1)/\Sigma} \otimes \dots \otimes \mathcal{J}_{X(e_r; z_r)/\Sigma} \otimes \mathcal{O}_C(C) \longrightarrow \mathcal{J}_{X(\underline{e}; \underline{z})/\Sigma} \cdot \mathcal{O}_C(C)$ induces a surjective morphism β on the level of global sections.⁶

If $s \in H^0(C, \mathcal{J}_{X(e_1; z_1)/\Sigma} \otimes_{\mathcal{O}_{\Sigma}} \dots \otimes_{\mathcal{O}_{\Sigma}} \mathcal{J}_{X(e_r; z_r)/\Sigma} \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C))$, but not in $\text{Ker}(\beta)$, then s induces a non-zero section \tilde{s} in $H^0(C, \pi_* \mathcal{O}_{\tilde{C}}(-\sum_{i=1}^r e_i E_i) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C))$.

Proof: Set $E := -\sum_{i=1}^r e_i E_i$.

We start with the structure sequence for \tilde{C} :

$$0 \longrightarrow \mathcal{O}_{\tilde{\Sigma}}(-\tilde{C}) \longrightarrow \mathcal{O}_{\tilde{\Sigma}} \longrightarrow \mathcal{O}_{\tilde{C}} \longrightarrow 0.$$

Tensoring with the locally free sheaf $\mathcal{O}_{\tilde{\Sigma}}(E)$ and then applying π_* we get a morphism:

$$\pi_* \mathcal{O}_{\tilde{\Sigma}}(E) \longrightarrow \pi_* \mathcal{O}_{\tilde{C}}(E).$$

Now tensoring by $\mathcal{O}_C(C)$ over \mathcal{O}_{Σ} we have an exact sequence:

$$0 \longrightarrow \text{Ker}(\gamma) \longrightarrow \pi_* \mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C) \xrightarrow{\gamma} \pi_* \mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C).$$

And finally taking global sections, we end up with:

$$0 \longrightarrow H^0(\Sigma, \text{Ker}(\gamma)) \longrightarrow H^0(\Sigma, \pi_* \mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)) \xrightarrow{\alpha} H^0(\Sigma, \pi_* \mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)).$$

⁴See Lemma C.1.

⁵I. e. $m_{\Sigma, z_1}^{e_1} \otimes \dots \otimes m_{\Sigma, z_r}^{e_r} \otimes \mathcal{O}_C(C) \longrightarrow m_{\Sigma, z_1}^{e_1} \cdot \dots \cdot m_{\Sigma, z_r}^{e_r} \cdot \mathcal{O}_C(C)$.

⁶ β is surjective, since $\text{supp}(\text{Ker}(\beta)) \subseteq \{z_1, \dots, z_r\}$ by Lemma C.4, and hence $H^1(\Sigma, \text{Ker}(\beta)) = 0$.

Since the sheaves we look at are actually \mathcal{O}_C -sheaves and since C is a closed subscheme of Σ , the global sections of the sheaves as sheaves on Σ and as sheaves on C coincide (cf. [Har77] III.2.10 - for more details, see Corollary C.3). Furthermore, $\pi_* \mathcal{O}_{\tilde{\Sigma}}(E) = \bigotimes_{i=1}^r \mathcal{J}_{X(e_i; z_i)/\Sigma}$.

Thus it suffices to show that $\text{Ker}(\alpha) \subseteq \text{Ker}(\beta)$.

Since $\pi| : \tilde{\Sigma} \setminus (\bigcup_{i=1}^r E_i) \rightarrow \Sigma \setminus \{z_1, \dots, z_r\}$ is an isomorphism, we have that $\text{supp}(\text{Ker}(\gamma)) \subseteq \{z_1, \dots, z_r\}$ is finite by Lemma C.5. Hence, by Lemma C.6 $\text{Ker}(\gamma)$ is a torsion sheaf, and thus

$$\text{Ker}(\alpha) = H^0(\Sigma, \text{Ker}(\gamma)) \subseteq H^0\left(\Sigma, \text{Tor}\left(\bigotimes_{i=1}^r \mathcal{J}_{X(e_i; z_i)/\Sigma} \otimes \mathcal{O}_C(C)\right)\right).$$

Let now $t \in \text{Ker}(\alpha)$ be given. We have to show that $\beta(t) = 0$, i. e. $\beta_z(t_z) = 0$ for every $z \in \Sigma$. If $z \notin \{z_1, \dots, z_r\}$, then $t_z = 0$. Thus we may assume $z = z_k$. As we have shown,

$$t_{z_k} \in \text{Tor}\left(\mathfrak{m}_{\Sigma, z_k}^{e_k} \otimes_{\mathcal{O}_{\Sigma, z_k}} \mathcal{O}_{C, z_k}\right) = \text{Tor}\left(\mathfrak{m}_{\Sigma, z_k}^{e_k} / f_{z_k} \mathfrak{m}_{\Sigma, z_k}^{e_k}\right) = (f_{z_k}) / f_{z_k} \mathfrak{m}_{\Sigma, z_k}^{e_k},$$

where f_{z_k} is a local equation of C at z_k . Therefore, there exists a $0 \neq g_{z_k} \in \mathcal{O}_{\Sigma, z_k}$ such that $t_{z_k} = f_{z_k} g_{z_k} \pmod{f_{z_k} \mathfrak{m}_{\Sigma, z_k}^{e_k}} \equiv f_{z_k} \otimes g_{z_k}$ (note that $f_{z_k} \in \mathfrak{m}_{\Sigma, z_k}^{n_k} \subseteq \mathfrak{m}_{\Sigma, z_k}^{e_k}$!). But then $\beta_{z_k}(t_{z_k})$ is just the residue class of $f_{z_k} g_{z_k}$ in $\mathfrak{m}_{\Sigma, z_k}^{e_k} \mathcal{O}_{C, z_k} = \mathfrak{m}_{\Sigma, z_k}^{e_k} / (f_{z_k})$, and is thus zero. \square

Proof of Lemma 2.1: Using the notation of the idea of the proof given on page 17, we have, by Lemma 2.3, a non-zero section $s \in H^0(C, \mathcal{J}_{X(\underline{e}; z)/\Sigma} \cdot \mathcal{O}_C(C))$. This lifts under the surjection β to a section $s' \in H^0(C, \bigotimes_{i=1}^r \mathcal{J}_{X(e_i; z_i)/\Sigma} \otimes \mathcal{O}_C(C))$ which is not in the kernel of β . Again setting $E := -\sum_{i=1}^r e_i E_i$, by Lemma 2.4, we have a non-zero section $\tilde{s} \in H^0(C, \pi_* \mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C))$, where by the projection formula the latter is just $H^0(C, \pi_*(\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\tilde{C}}} \pi^* \mathcal{O}_C(C))) =_{\text{def}} H^0(\tilde{C}, \mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\tilde{C}}} \pi^* \mathcal{O}_C(C))$.

Since $\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\tilde{C}}} \pi^* \mathcal{O}_C(C)$ has a global section and since \tilde{C} is irreducible and reduced, we get by Lemma D.2:

$$\begin{aligned} 0 \leq \deg(\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\tilde{C}}} \pi^* \mathcal{O}_C(C)) &= \deg(\mathcal{O}_{\tilde{C}}(E)) + \deg(\pi^* \mathcal{O}_C(C)) \\ &= E \cdot \tilde{C} + \deg(\mathcal{O}_C(C)) = \sum_{i=1}^r -e_i n_i + C^2. \end{aligned}$$

\square

3. Examples

In this section we are going to examine the conditions in the vanishing theorem (Corollary 1.2). Unless otherwise stated, $r \geq 1$ is a positive integer, and $m_1, \dots, m_r \in \mathbb{N}_0$ are non-negative.

3.a. The Classical Case - $\Sigma = \mathbb{P}_\mathbb{C}^2$

Since in $\mathbb{P}_\mathbb{C}^2$ there are no irreducible curves of self-intersection number zero, condition (1.5) is redundant. Moreover, condition (1.6) takes in view of (1.4) the form $d + 3 \geq \sqrt{2}$. Corollary 1.2 thus takes the following form, where $L \in |\mathcal{O}_{\mathbb{P}_\mathbb{C}^2}(1)|_l$ is a generic line.

1.2a Corollary

Let d be any integer such that

$$(1.4a) \quad (d + 3)^2 \geq 2 \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.6a) \quad d \geq -1.$$

Then for $z_1, \dots, z_r \in \mathbb{P}_\mathbb{C}^2$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\mathbb{P}_\mathbb{C}^2), d \pi^* L - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.b. Geometrically Ruled Surfaces

Throughout this section we use the notation and the results from Section G.a, in particular Lemma G.2 for the irreducible curves with selfintersection zero on Σ .

1.2b Corollary

Given two integers $a, b \in \mathbb{Z}$ satisfying

$$(1.4b) \quad a(b - (\frac{a}{2} - 1)e) \geq \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.5b.i) \quad a > \max\{m_i \mid i = 1, \dots, r\},$$

$$(1.5b.ii) \quad b > \max\{m_i \mid i = 1, \dots, r\}, \text{ if } e = 0,$$

$$(1.5b.iii) \quad 2(b - (\frac{a}{2} - 1)e) > \max\{m_i \mid i = 1, \dots, r\}, \text{ if } e < 0, \text{ and}$$

$$(1.6b) \quad b \geq (a - 1)e, \text{ if } e > 0.$$

For $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), (a - 2) \cdot \pi^* C_0 + (b - 2 + 2g) \cdot \pi^* F - \sum_{i=1}^r m_i E_i \right) = 0.$$

Proof: Note that if the invariant e is non-positive, then $(b - (\frac{a}{2} - 1)e) > 0$ implies

$$b \geq (a - 1)e, \tag{3.1}$$

so that this inequality is fulfilled for any choice of e .

Setting $D = (a - 2)C_0 + (b - 2 + 2g)F$ we have

$$(D - K_\Sigma)^2 = (aC_0 + (b + e)F)^2 = 2a \left(b - \left(\frac{a}{2} - 1 \right) e \right) \geq 2 \sum_{i=1}^r (m_i + 1)^2,$$

which is just (1.4b.i/ii/iii). Similarly, by (1.5b) and Lemma G.2 condition (1.5) is satisfied.⁷ Finally, in view of Lemma G.1, condition (1.6b) implies that $D - K_\Sigma$ is nef. \square

3.c. Products of Curves

As we have seen in Proposition G.12, for a generic choice of smooth projective curves of genera $g_1 \geq 1$ and $g_2 \geq 1$ respectively the surface $\Sigma = C_1 \times C_2$ has Picard number two and according to Remark G.11 the only irreducible curves $B \subset \Sigma$ with selfintersection $B^2 = 0$ are the fibres C_1 and C_2 , and for any irreducible curve $B \sim_a aC_1 + bC_2$ the coefficients a and b must be non-negative. Taking into account that $K_\Sigma \sim_a (2g_2 - 2)C_1 + (2g_1 - 2)C_2$ Corollary 1.2 comes down to the following.

1.2c Corollary

Let C_1 and C_2 be two generic curves with $g(C_i) = g_i \geq 1$, $i = 1, 2$, and let $a, b \in \mathbb{Z}$ be integers satisfying

$$(1.4c) \quad (a - 2g_2 + 2) \cdot (b - 2g_1 + 2) \geq \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

$$(1.5c) \quad (a - 2g_2 + 2), (b - 2g_1 + 2) > \max\{m_i \mid i = 1, \dots, r\},$$

then for $z_1, \dots, z_r \in \Sigma = C_1 \times C_2$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), a\pi^*C_1 + b\pi^*C_2 - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.d. Products of Elliptic Curves

That C_1 and C_2 be “generic” in the above sense means for elliptic curves just that they are non-isogenous.

In view of (1.5d) and Lemma G.19 (iv) the condition (1.6) becomes obsolete, and Corollary 1.2 has the following form, taking Lemma G.19 (iii) and $K_\Sigma = 0$ into account.

1.2d Corollary

Let C_1 and C_2 be two non-isogenous elliptic curves, $a, b \in \mathbb{Z}$ be such that

$$(1.4d) \quad ab \geq \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

⁷To see this, let $B \sim_a a'C_0 + b'F$ be an irreducible curve with $B^2 = 0$. Then by Lemma G.2 either $a' = 0$ and $b' = 1$, or $e = 0$, $a' \geq 1$ and $b' = 0$, or $e < 0$, $a' \geq 2$, and $b' = \frac{a'}{2}e < 0$. In the first case, $(D - K_\Sigma).B = a > \max\{m_i \mid i = 1, \dots, r\}$ by (1.5b.i). In the second case, $(D - K_\Sigma).B = ba' \geq b > \max\{m_i \mid i = 1, \dots, r\}$ by (1.5b.ii). And finally, in the third case, we have $(D - K_\Sigma).B = a' \cdot (b - (\frac{a'}{2} - 1)e) > \max\{m_i \mid i = 1, \dots, r\}$ by (1.5b.iii).

$$(1.5d) \quad a, b > \max\{m_i \mid i = 1, \dots, r\},$$

then for $z_1, \dots, z_r \in \Sigma = C_1 \times C_2$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), a \pi^* C_1 + b \pi^* C_2 - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.e. Surfaces in \mathbb{P}_c^3

Since we consider the case of rational surfaces separately the following considerations thus give a full answer for the “general case” of a surface in \mathbb{P}_c^3 .

1.2e Corollary

Let $\Sigma \subset \mathbb{P}_c^3$ be a surface in \mathbb{P}_c^3 of degree n , $H \in \text{NS}(\Sigma)$ be the algebraic class of a hyperplane section, and d an integer satisfying

$$(1.4e) \quad n \cdot (d - n + 4)^2 \geq 2 \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

$$(1.5e) \quad (d - n + 4) \cdot H \cdot B > \max\{m_i \mid i = 1, \dots, r\} \text{ for any irreducible curve } B \text{ with } B^2 = 0 \text{ and } \dim |B|_a \geq 1, \text{ and}$$

$$(1.6e) \quad d \geq n - 4,$$

then for $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), d \pi^* H - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.1 Remark

- (a) If $\text{NS}(\Sigma) = H\mathbb{Z}$, then (1.5e) is redundant, since there are no irreducible curves B with $B^2 = 0$. Otherwise we would have $B \sim_a kH$ for some $k \in \mathbb{Z}$ and $k^2 n = B^2 = 0$ would imply $k = 0$, but then $H \cdot B = 0$ in contradiction to H being ample (see Lemma E.1).
- (b) By a Theorem of Noether a generic surface in \mathbb{P}_c^3 has Picard number one. However, a quadric in \mathbb{P}_c^3 or the K3-surface given by $w^4 + x^4 + y^4 + z^4 = 0$ contain irreducible curves of self-intersection zero.
- (c) If $\sum_{i=1}^r (m_i + 1)^2 > \frac{n}{2} m_i^2$ for all $i = 1, \dots, r$ then again (1.5e) becomes obsolete in view of (1.4e), since $H \cdot B > 0$ anyway. The above inequality is, for instance, fulfilled if the highest multiplicity occurs at least $\frac{n}{2}$ times.
- (d) In the existence theorems the condition depending on curves of self-intersection will vanish in any case, see Section III.3.e.

3.f. K3-Surfaces

3.f.i. *Generic K3-Surfaces.* Since a generic K3-surface does not possess an elliptic fibration the following version of Corollary 1.2 applies for generic K3-surfaces. (cf. [FrM94] I.1.3.7)

1.2f.i Corollary

Let Σ be a K3-surface which is not elliptic, and let D a divisor on Σ satisfying

$$(1.4f) \quad D^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

$$(1.6f) \quad D \text{ nef,}$$

then for $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), \pi^*D - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.f.ii. *K3-Surfaces with an Elliptic Structure.* The hypersurface in $\mathbb{P}_\mathbb{C}^3$ given by the equation $x^4 + y^4 + z^4 + u^4 = 0$ is an example of a K3-surface which is endowed with an elliptic fibration. Among the elliptic K3-surfaces the general one will possess a unique elliptic fibration while there are examples with infinitely many different such fibrations. (cf. [FrM94] I.1.3.7)

1.2f.ii Corollary

Let Σ be a K3-surface which possesses an elliptic fibration, and let D be a divisor on Σ satisfying

$$(1.4f) \quad D^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.5f) \quad D \cdot B > \max\{m_i \mid i = 1, \dots, r\} \text{ for any irreducible curve } B \text{ with } B^2 = 0, \text{ and}$$

$$(1.6f) \quad D \text{ nef,}$$

then for $z_1, \dots, z_r \in \Sigma$ in very general position and $\nu > 0$

$$H^\nu \left(\text{Bl}_{\underline{z}}(\Sigma), \pi^*D - \sum_{i=1}^r m_i E_i \right) = 0.$$

3.2 Remark

If Σ is generic among the elliptic K3-surfaces, i. e. admits exactly one elliptic fibration, then condition (1.5f) means that a curve in $|D|_1$ meets a general fibre in at least $k = \max\{m_i \mid i = 1, \dots, r\}$ distinct points.

4. Yet Another Vanishing Theorem

In [GLS98c] a different approach is used for the existence of curves with prescribed topological singularity types in the plane case. It relies once more on vanishing theorems for singularity schemes and may be applied to other surfaces as well, once we have the corresponding vanishing theorems. It is the aim of this section to generalise the first of these vanishing theorems (cf. [GLS98c] Lemma 3.1 or [Los98] Lemma 3.10), and we claim that the second one (cf. [GLS98c] Lemma 4.1 or [Los98] Lemma 3.11) can be treated analogously.

The proof uses an induction based on the reduction of a singularity scheme X by a smooth curve L , that is, replacing X by $X : L$. Unfortunately, the class of singularity schemes is not closed under reduction, it therefore becomes necessary to consider a wider class of zero-dimensional schemes. We only give the definitions and state the results, which are necessary for Theorem 4.5. For a more thorough investigation of these schemes, including many examples, we refer to [GLS98c] Section 2 or [Los98] Section 2.2.

4.1 Definition

Let $(C, z) \subset (\Sigma, z)$ be a reduced plane curve singularity with complete embedded resolution tree $T(C, z)$, and let $T^*(C, z)$ denote the essential subtree of $T(C, z)$.

- (a) If T^* is a finite, connected subtree of $T(C, z)$ containing the essential subtree $T^*(C, z)$, we call the zero-dimensional scheme $X(C, T^*)$ defined by the ideal sheaf with stalks

$$\mathcal{I}_{X(C, T^*)/\Sigma, z} = I(C, T^*) = \left\{ g \in \mathcal{O}_{\Sigma, z} \mid g \text{ goes through the cluster } \mathcal{C}l(C, T^*) \right\}$$

and $\mathcal{I}_{X(C, T^*)/\Sigma, z'} = \mathcal{O}_{\Sigma, z'}$, whenever $z' \neq z$, a *generalised singularity scheme* (with centre z).

We denote by \mathcal{GS} the class of zero-dimensional schemes in Σ which may be obtained that way.

- (b) The subclass of \mathcal{GS} of zero-dimensional schemes of the form $X(C, T^*)$ with centre z , where the plane curve singularity (C, z) has only *smooth branches*, is denoted by \mathcal{GS}_1 .
- (c) The subclass of \mathcal{GS} of zero-dimensional schemes of the form $X(C, T^*)$ with centre z where $T^* = T^*(C, z)$ is denoted by \mathcal{S} , and its members are called *singularity schemes*.

We note that in this sense the singularity schemes $X^s(C)$ introduced in Definition I.2.6 are finite unions of singularity schemes. (See Remark 4.2.)

- (d) By \mathcal{OS} we denote the subclass of \mathcal{S} of ordinary fat point schemes $X(m; z)$.
- (e) Since the support of a scheme $X \in \mathcal{GS}$ consists of a unique point z , we may define $\text{mult}(X) = \text{mult}(X, z)$.
- (f) Let $X = X(C, T^*) \in \mathcal{GS}$ and $L \subset \Sigma$ be a curve which is smooth at z . We then define $T^* \cap L = \{q \in T^* \mid \text{the strict transform of } L \text{ goes through } q\}$, that is, $T^* \cap L$ is the maximal subtree T of T^* such that L goes through the cluster $\mathcal{C}l(C, T)$.
- (g) Two generalised singularity schemes $X_0, X_1 \in \mathcal{GS}$ with centre z are called *isomorphic*, if they are isomorphic as subschemes of Σ . We then write $X_0 \cong X_1$.

- (h) An *equimultiple family* of plane curve singularities over a (reduced) algebraic scheme T over \mathbb{C} is a flat family

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & S \times T \\ & \swarrow \sigma & \searrow \\ & T & \end{array}$$

of reduced plane curve singularities $(\mathcal{C}_t, \sigma(t)) \subset \Sigma \times \{t\} \cong \Sigma$ with section σ , which admits a simultaneous embedded resolution, together with sections σ_q through infinitely near points, defining a family \mathcal{T}^* of finite connected trees T_t^* with $T^*(\mathcal{C}_t, \sigma(t)) \subseteq T_t^* \subset T(\mathcal{C}_t, \sigma(t))$, such that the total transform of \mathcal{C} is equimultiple along σ_q , $\sigma_q(t) \in T_t^*$. (Cf. [GLS98c] Definition 2.9, and [Los98] Definition 2.14.)

- (i) Two generalised singularity schemes $X_0, X_1 \in \mathcal{GS}$ with centre z are called *equivalent*, if there exist germs (C_0, z) and (C_1, z) defining X_0 respectively X_1 , and a T^* -equimultiple family of plane curve singularities over some (reduced) open connected subset T of $\mathbb{A}_{\mathbb{C}}^1$ having (C_0, T_0^*) and (C_1, T_1^*) as fibres. We then write $X_0 \sim X_1$. (Cf. [GLS98c] Definition 2.9, and [Los98] Definition 2.28.)

4.2 Remark

The classes \mathcal{GS} , \mathcal{GS}_1 and \mathcal{OS} are closed with respect to the equivalence relation \sim and with respect to reduction by a smooth curve L . (Cf. [GLS98c] Proposition 2.11 and Lemma 2.13, or [Los98] Proposition 2.23 and Lemma 2.31.)

4.3 Remark

The concepts introduced so far immediately generalise to multigerms (C, \underline{z}) , and also the remaining part of this section does not change in this situation (cf. [GLS98c] p. 545). Moreover, the singularity schemes introduced in Definition I.2.6 then become precisely the elements of \mathcal{S} . Just in order to save some notation we will avoid the multigerms.

4.4 Definition

Let $L \subset \Sigma$ be a smooth curve such that the corresponding divisor is very ample.

- (a) Let $f_1, \dots, f_r : \text{Num}(\Sigma) \times \bigcup_{n \geq 1} \text{Hilb}_{\Sigma}^n \rightarrow \mathbb{Q}$ be some functions possibly depending on the numerical class of L , and let $\mathcal{K} \subseteq \bigcup_{n \geq 1} \text{Hilb}_{\Sigma}^n$ be some class of zero-dimensional schemes (not necessarily closed under reduction by L).

We then call (f_1, \dots, f_r) a *tuple of L-reduction resistant conditions on \mathcal{K}* if, whenever $f_i(D, X) \geq 0$, $i = 1, \dots, r$, for some zero-dimensional scheme $X \in \mathcal{K}$ and some divisor class $D \in \text{Num}(\Sigma)$ with $D.L > L^2$, then also $f_i(D - L, X : L) \geq 0$, $i = 1, \dots, r$.

We say, an L-reduction resistant condition f is *satisfied* by some zero-dimensional scheme X and some divisor class D , if $f(D, X) \geq 0$.

(b) We call a tuple (f_1, \dots, f_r) of L -reduction resistant conditions on \mathcal{OS} *suitable* if for any ordinary fat point scheme $X = X(\underline{m}, \underline{z}) \subset \Sigma$ and any divisor D , with $D \cdot L > L^2$, satisfying f_i , for $i = 1, \dots, r$, there exists a scheme $X' \in \mathcal{OS}$ with

- (1) $X' \sim X$,
- (2) $\deg(X' \cap L) = \deg(X \cap L)$, and
- (3) $h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(D)) = 0$.

Given a generalised singularity scheme X and a divisor D we would like to know under which circumstances $H^1(\Sigma, \mathcal{J}_{X/\Sigma}(D))$ vanishes. However, it turns out that this is hard to answer, and indeed, for the applications in the existence theorems (cf. [GLS98c]) we can do with less. There, we may replace X by an equivalent scheme X' for which the cohomology group vanishes, and it is the aim of Theorem 4.5 to provide numerical conditions ensuring the existence of X' . Well, actually the theorem reduces the problem to finding conditions which guarantee the existence of X' if X is an ordinary fat point scheme (cf. (4.1)), which is much simpler to handle and where the geometry of Σ will come into play.

4.5 Theorem

Let $L \subset \Sigma$ be a smooth curve of genus $g = g(L)$ such that the corresponding divisor is very ample, and let $D \in \text{Div}(\Sigma)$ with $D \cdot L \geq L^2$ and $X \in \mathcal{GS}_1$ such that:

(4.1) *There is a suitable tuple of L -reduction resistant conditions on \mathcal{OS} which are satisfied by X and D .*

(4.2) $\deg(X) \leq \beta \cdot \left(\left(\frac{D \cdot L}{L^2} - \text{mult}(X) \right)^2 - \frac{2\alpha g}{(\alpha-1)} \cdot \frac{D \cdot L}{L^2} \right)$, and

(4.3) $\deg(X \cap L) \leq \frac{D \cdot L}{L^2} - \frac{\alpha \cdot L^2}{D \cdot L} \cdot \deg(X) - 2g$,

where $\alpha = 1 + \sqrt{2}$ and $\beta = 3 - 2 \cdot \sqrt{2}$.

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(D)) = 0.$$

Proof: We do the proof by induction on $d = D \cdot L$, starting with $d = L^2$. We note that we may suppose that X is not an ordinary fat point scheme, since the theorem holds for these by Assumption (4.1). This gives in particular the induction basis, since for $d = L^2$ (4.2) can only be satisfied if $X = \emptyset$.

Setting \mathcal{K} the subclass of \mathcal{GS}_1 which are not ordinary fat point schemes, Condition (4.2) and Condition (4.3) are L -reduction resistant on \mathcal{K} by Lemma 4.7. We then may replace X and D by $X : L$ and $D - L$, satisfying now (4.1)–(4.3). Thus, by induction there is a scheme $Y' \sim X : L$ in \mathcal{GS}_1 such that $\deg(Y' \cap L) = \deg((X : L) \cap L)$ and $h^1(\Sigma, \mathcal{J}_{Y'/\Sigma}(D - L)) = 0$. Due to the

Semicontinuity Theorem for h^1 (cf. [Har77] III.12.8) and Lemma 4.8 (a) we then find a scheme $X' \in \mathcal{GS}$ such that $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$. Moreover, in view of Remark 4.2 we have $X' \in \mathcal{GS}_1$, and by Lemma III.1.1 and Condition (4.3) we find

$$h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(D)) = 0,$$

which finishes the proof. \square

4.6 Remark

(a) We note that for an ordinary fat point scheme $X(m; z)$ the cohomology group $H^1(\Sigma, \mathcal{J}_{X(m; z)/\Sigma}(D))$ vanishes if and only if

(1) $h^1(\Sigma, \mathcal{O}_\Sigma(D)) = 0$, and

(2) $H^0(\Sigma, \mathcal{O}_\Sigma(D)) \rightarrow H^0(\Sigma, \mathcal{O}_{X(m; z)}(D)) = \mathcal{O}_{\Sigma, z}/\mathfrak{m}_{\Sigma, z}^m$ is surjective.

(b) We call a surface Σ *regular with respect to* L if for any $d \geq 1$ we have $h^1(\Sigma, \mathcal{O}_\Sigma(dL)) = 0$.

We then see that the examples which we are considering throughout this thesis are all regular with respect to the obvious choices of very ample divisors.

- (1) Complete intersections are regular with respect to hyperplane sections by Proposition G.20. In particular \mathbb{P}_c^2 and surfaces in \mathbb{P}_c^3 are so.
- (2) On a Hirzebruch surfaces F_e , $e \geq 0$, a divisor $L = \alpha C_0 + \beta F$ is very ample if and only if $\alpha > 0$ and $\beta > \alpha e$ (cf. [Har77] V.2.18). Then, however, F_e is regular with respect to L (cf. Remark G.4 and Lemma G.6).
- (3) For a geometrically ruled surface Σ with invariant e over a curve C of genus $g = g(C) \geq 1$, we may choose a suitable integer $l \geq \{2g-1, e+2g-1\}$ such that there is a very ample divisor $L \sim_a C_0 + lF$. By Lemma G.5 Σ is then regular with respect to L .
- (4) Let $\Sigma = C_1 \times C_2$ be a product of two smooth projective curves of genera $g_i = g(C_i)$ and denote by $\text{pr}_i : \Sigma \rightarrow C_i$ the canonical projection, $i = 1, 2$. If $L = \text{pr}_2^* a \otimes \text{pr}_1^* b$ with $a \in \text{Div}(C_2)$ of degree $a \geq 2g_2 + 1$ and $b \in \text{Div}(C_1)$ of degree $b \geq 2g_1 + 1$, then L is very ample by Lemma G.7, and since a and b are non-special by Riemann-Roch (cf. [Har77] IV.1.3.4) the Künneth formula in Lemma G.8 implies that $h^1(\Sigma, L) = 0$. But then Σ is regular with respect to L .
- (5) If Σ is a product of two elliptic curves, then $K_\Sigma = 0$, and thus by the Kodaira Vanishing Theorem Σ is regular with respect to any very ample divisor.
- (6) If Σ is a K3-surface, again the Kodaira Vanishing Theorem implies that Σ is regular with respect to any very ample divisor L .

- (c) If $L \in \text{Div}(\Sigma)$ is very ample such that Σ is regular with respect to L and if $D = d \cdot L$ is a multiple of the very ample divisor L , then (4.1) just becomes obsolete.

In this situation Condition (4.2) and Condition (4.3) take the form:

$$(4.2^*) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot \left((d - \text{mult}(X))^2 - (2 + \sqrt{2}) \cdot g \cdot d \right),$$

$$(4.3^*) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X) - 2g.$$

- (d) In the case of the Hirzebruch surface $\Sigma = \mathbb{F}_0 = \mathbb{P}_c^1 \times \mathbb{P}_c^1$ we would like to use the chance to replace (4.1) by some condition, which applies to arbitrary divisors $D = aC_0 + bF$ on Σ rather than only to multiples of $L = C_0 + F$. We claim that (4.1) may be replaced by

$$D \cdot F = a \geq \text{mult}(X) - 1 \quad \text{and} \quad D \cdot C_0 = b \geq \text{mult}(X) - 1. \quad (4.4)$$

Proof: (a) and (b) are obvious.

- (c) L is a hyperplane section of some embedding $\Sigma \hookrightarrow \mathbb{P}_c^N$, and we thus may find two curves $L_x, L_y \in |L|_1$ such that the germs $L_{x,z}$ and $L_{y,z}$ at z are local coordinates of (Σ, z) , that is $\mathfrak{m}_{\Sigma,z} = (L_{x,z}, L_{y,z})$.

If $X = X(m; z)$ is an ordinary fat point scheme such that (4.2) and (4.3) are satisfied by D and X . We then deduce from (4.11) below, which is an immediate consequence of (4.2), that

$$m = \text{mult } X \leq \frac{D \cdot L}{2 \cdot L^2} < d.$$

But then the map in (a) is surjective.

- (d) The global sections of $\mathcal{O}_\Sigma(D)$ may be identified with the polynomials in $\mathbb{C}[x_0, x_1, y_0, y_1]$ which are bihomogenous of bidegree (a, b) . Assuming that $z = ((0 : 1), (0 : 1))$ the map in (a) comes down to

$$H^0(\Sigma, aC_0 + bF) \rightarrow \mathcal{O}_{\Sigma,z}/\mathfrak{m}_{\Sigma,z}^m \cong \mathbb{C}[x_0, y_0]/(x_0, y_0)^m : f \mapsto f(x_0, 1, y_0, 1),$$

which is surjective by Assumption (4.4).

It remains to show that the conditions are L -reduction resistant on \mathcal{O}_S . However, if $X(m; z)$ is an ordinary fat point scheme and L is smooth at z , then $\text{mult}(X(m; z) : L) = \text{mult}(X(m; z)) - 1$, while $D - L = (a - 1) \cdot C_0 + (b - 1) \cdot F$, which finishes the claim. □

4.7 Lemma

Let \mathcal{K} denote the subclass of \mathcal{GS}_1 of schemes which are not ordinary fat point schemes, let $L \subset \Sigma$ be a smooth curve of genus $g = g(L)$ such that the corresponding divisor is very ample, and let $\alpha = 1 + \sqrt{2}$, $\beta = 3 - 2 \cdot \sqrt{2}$ and $\gamma = \frac{2\alpha}{\alpha-1} \cdot g = (2 + \sqrt{2}) \cdot g$.

We define two conditions $f_A, f_B : \text{Num}(\Sigma) \times \bigcup_{n \geq 1} \text{Hilb}_\Sigma^n \rightarrow \mathbb{Q}$ by

$$f_A(D, X) = A(D, X) - \deg(X) \quad \text{and} \quad f_B(D, X) = B(D, X) - \deg(X \cap L),$$

where

$$A(D, X) = \beta \cdot \left(\left(\frac{D \cdot L}{L^2} - \text{mult}(X) \right)^2 - \gamma \cdot \frac{D \cdot L}{L^2} \right)$$

and

$$B(D, X) = \frac{D \cdot L}{L^2} - \frac{\alpha \cdot L^2}{D \cdot L} \cdot \text{deg}(X) - 2g.$$

The tuple (f_A, f_B) is L -reduction resistant on \mathcal{K} .

Proof: For $X \in \mathcal{K}$ and $D \in \text{Num}(\Sigma)$ with $D \cdot L > L^2$, $\text{deg}(X) \leq A(D, X)$ and $\text{deg}(X \cap L) \leq B(D, X)$ we have to show

$$\text{deg}(X : L) \leq A(D - L, X : L) \quad (4.5)$$

and

$$\text{deg}((X : L) \cap L) \leq B(D - L, X : L). \quad (4.6)$$

For this we consider four different cases, where the first case shall illustrate that the constants α and β are chosen optimal.

Step 1: Some useful considerations on α , β and γ .

We claim that $\alpha > 0$ is minimal and $\beta > 0$ maximal such that

$$\beta \cdot (\alpha^2 + \alpha) = \alpha - 1. \quad (4.7)$$

To see this we consider for a fixed $b > 0$ the equation

$$b \cdot a^2 + (b - 1) \cdot a + 1 = 0. \quad (4.8)$$

The discriminant $(b - 1)^2 - 4b$ vanishes if and only if $b = \beta$, and in this case $a = -\frac{\beta-1}{2\beta} = \alpha$, which proves the claim.

Furthermore, by the definition of α , β and γ we have

$$(\alpha + 1) \cdot \beta \cdot \gamma = 2g, \quad (4.9)$$

$$1 - \alpha\beta - 2\beta = 3\sqrt{2} - 4 > 0, \quad (4.10)$$

and $\text{mult}(X)^2 \leq 2 \cdot \text{deg}(X) \leq 2 \cdot A(D, X) \leq 2\beta \cdot \left(\frac{D \cdot L}{L^2} - \text{mult}(X) \right)^2$, which implies

$$\text{mult}(X) < \frac{1+\sqrt{2\beta}}{\sqrt{2\beta}} \cdot \text{mult}(X) \leq \frac{D \cdot L}{L^2}. \quad (4.11)$$

Step 2: Suppose that $\text{deg}(X \cap L) = B(D, X)$.

Since $\beta \cdot (\alpha^2 + \alpha) \leq \alpha - 1$ by (4.7), we have

$$\begin{aligned} (L^2)^2 \cdot (\alpha^2 + \alpha) \cdot \text{deg}(X) &\leq (L^2)^2 \cdot \frac{(\alpha-1)}{\beta} \cdot A(D, X) \\ &\leq (\alpha - 1) \cdot (D \cdot L)^2 - L^2 \cdot (\alpha - 1) \cdot \gamma \cdot D \cdot L \\ &< (\alpha - 1) \cdot (D \cdot L)^2 + (L^2 - 2\alpha\gamma \cdot L^2) \cdot D \cdot L. \end{aligned}$$

Taking the assumption of Step 2 into account, this implies

$$\frac{\alpha \cdot L^2}{D \cdot L} \cdot \text{deg}(X) \geq 1 + \frac{\alpha \cdot L^2}{D \cdot L - L^2} \cdot (\text{deg}(X) - B(D, X)) = \frac{L^2}{L^2} + \frac{\alpha \cdot L^2}{D \cdot L - L^2} \cdot \text{deg}(X : L).$$

But then (4.6) follows:

$$\deg((X : L) \cap L) \leq \deg(X \cap L) \leq B(D, X) \leq B(D - L, X : L).$$

Due to (4.9)–(4.11) we have

$$\begin{aligned} C(D, X) &:= (D \cdot L)^2 \cdot (1 - \alpha\beta - 2\beta) + \alpha\beta \cdot \text{mult}(X) \cdot L^2 \cdot (2 \cdot D \cdot L - L^2 \cdot \text{mult}(X)) + \\ &D \cdot L \cdot L^2 \cdot (2\beta \cdot \text{mult}(X) + \beta + (\alpha + 1) \cdot \beta\gamma - 2g) \geq 0. \end{aligned}$$

In view of the assumption of Step 2 and (4.9) a tedious calculation shows

$$\begin{aligned} \deg(X : L) &= \deg(X) - \deg(X \cap L) = \deg(X) - B(D, X) \\ &= \left(1 + \frac{\alpha \cdot L^2}{D \cdot L}\right) \cdot \deg(X) - \frac{D \cdot L}{L^2} + 2g \leq \left(1 + \frac{\alpha \cdot L^2}{D \cdot L}\right) \cdot A(D, X) - \frac{D \cdot L}{L^2} + 2g \\ &= \beta \cdot \left(\left(\frac{D \cdot L - L^2}{L^2} - \text{mult}(X) \right)^2 - \gamma \cdot \frac{D \cdot L - L^2}{L^2} \right) - \frac{C(D, X)}{D \cdot L \cdot L^2} \\ &\leq \beta \cdot \left(\left(\frac{D \cdot L - L^2}{L^2} - \text{mult}(X) \right)^2 - \gamma \cdot \frac{D \cdot L - L^2}{L^2} \right) \leq A(D - L, X : L), \end{aligned}$$

which gives (4.5).

Step 3: We now suppose that $X = X(C, T^*)$ satisfies the following conditions:

- (a) $\deg(X \cap L) < B(D, X)$.
- (b) There exists an irreducible branch Q of (C, z) such that $T^* \cap L = T^* \cap Q$.
- (c) There exists no irreducible branch Q' of (C, z) such that $T^* \cap L \subsetneq T^* \cap Q'$.

Since X is not an ordinary fat point scheme, the tree T^* has at least three vertices, and due to (c) the tree $T^* \cap L$, therefore, has at least two. But then $\deg(X : L) < \deg(X)$ and $\deg((X : L) \cap L) \leq \deg(X \cap L) - 2$. Moreover, since $\alpha\beta < 1$ and by (4.11) we have

$$\alpha \cdot \deg(X) < \left(\frac{D \cdot L}{L^2} - \text{mult}(X) \right)^2 \leq \frac{D \cdot L}{L^2} \cdot (D \cdot L - L^2),$$

and thus, taking (a) into account, we get (4.6):

$$\begin{aligned} \deg((X : L) \cap L) &< B(D, X) - 2 \\ &= B(D - L, X : L) + \frac{\alpha \cdot L^2}{D \cdot L \cdot (D \cdot L - L^2)} \cdot \deg(X) - 1 \leq B(D - L, X : L). \end{aligned}$$

By (b) and Lemma 4.8 (b) we know that $\text{mult}(X : L) = \text{mult}(X) - 1$. Therefore,

$$\deg(X : L) < \deg(X) \leq A(D, X) = A(D - L, X : L) - \gamma,$$

which implies (4.5).

Step 4: We now suppose that $X = X(C, T^*)$ satisfies the following conditions:

- (a) $\deg(X \cap L) < B(D, X)$.
- (b) There exists an irreducible branch Q of (C, z) such that
 - (1) $T^* \cap Q = \{z = q_0 < q_1 < \dots < q_l\}$,
 - (2) $T^* \cap L = \{z = q_0 < q_1 < \dots < q_s\}$, $s < l$, and
 - (3) $\text{mult}(C_{(q_{s+1}), q_{s+1}}) = m$ with $\deg(X \cap L) + m \geq B(D, X)$.

Due to (4.10) and (4.11) we have

$$C'(D, X) := (1 - \alpha\beta - 2\beta) \cdot \left(\frac{D \cdot L}{L^2} - \text{mult}(X)\right) + \beta + \frac{\alpha\beta \cdot \text{mult}(X) \cdot L^2}{D \cdot L} \cdot \left(\frac{D \cdot L}{L^2} - \text{mult}(X)\right) \geq 0.$$

But then, taking (3) and (4.9) into account, a simple calculation shows

$$\begin{aligned} \deg(X : L) &= \deg(X) - \deg(X \cap L) \leq \deg(X) - B(D, X) + \text{mult}(X) \\ &= \left(1 + \frac{\alpha \cdot L^2}{D \cdot L}\right) \cdot \deg(X) - \left(\frac{D \cdot L}{L^2} - \text{mult}(X)\right) + 2g \\ &\leq \left(1 + \frac{\alpha \cdot L^2}{D \cdot L}\right) \cdot A(D, X) - \left(\frac{D \cdot L}{L^2} - \text{mult}(X)\right) + 2g \\ &= \beta \cdot \left(\left(\frac{D \cdot L - L^2}{L^2} - \text{mult}(X)\right)^2 - \gamma \cdot \frac{D \cdot L - L^2}{L^2}\right) - C'(D, X) \\ &\leq \beta \cdot \left(\left(\frac{D \cdot L - L^2}{L^2} - \text{mult}(X)\right)^2 - \gamma \cdot \frac{D \cdot L - L^2}{L^2}\right) \leq A(D - L, X : L), \end{aligned}$$

which gives (4.5).

Consider next the concave function $\varphi : \mathbb{R} \rightarrow \mathbb{R} : x \mapsto \beta \cdot (1 + \alpha) \cdot \left(2x - \frac{L^2}{D \cdot L} \cdot x^2\right)$, which takes its maximum at $x = \frac{D \cdot L}{L^2}$ and is thus concavely increasing on the interval $\left[0, \frac{\sqrt{2\beta}}{1 + \sqrt{2\beta}} \cdot \frac{D \cdot L}{L^2}\right]$. In particular, since the line $\{y = x\}$ intersects the graph of φ in $x = 0$ and $x = \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} \cdot \frac{D \cdot L}{L^2}$, $\varphi(x) \geq x$ for all $0 \leq x \leq \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} \cdot \frac{D \cdot L}{L^2}$.

Now consider a second function $\psi : \left[0, \frac{\sqrt{2\beta}}{1 + \sqrt{2\beta}} \cdot \frac{D \cdot L}{L^2}\right] \rightarrow \mathbb{R}$ given by

$$\psi(x) := \begin{cases} x, & \text{for } 0 \leq x \leq \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} \cdot \frac{D \cdot L}{L^2}, \\ \sqrt{2\beta \cdot \left(\frac{D \cdot L}{L^2} - x\right)^2 - x^2}, & \text{for } \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} \cdot \frac{D \cdot L}{L^2} \leq x \leq \frac{\sqrt{2\beta}}{1 + \sqrt{2\beta}} \cdot \frac{D \cdot L}{L^2}. \end{cases}$$

This function is concave as well and takes its maximum at $x = \frac{\sqrt{\beta}}{1 + \sqrt{\beta}} \cdot \frac{D \cdot L}{L^2}$, where ψ and φ coincide. In particular, ψ is bounded from above by φ .

Note that by Lemma 4.8 (c) we have $\frac{\text{mult}(X)^2 + m^2}{2} < \deg(X) \leq \beta \cdot \left(\frac{D \cdot L}{L^2} - \text{mult}(X)\right)^2$; and $m \leq \text{mult}(X)$ anyway. We thus find

$$m \leq \psi(\text{mult}(X)) \leq \varphi(\text{mult}(X)),$$

since by (4.11) $\text{mult}(X)$ takes its values in the domain of definition of ψ .

Taking (4.8) and (4.9) into account, we deduce from this

$$\begin{aligned} C''(D, X) &:= \frac{D \cdot L}{L^2} \cdot (\alpha - 1) + (1 - \alpha m - 2\alpha g) - \frac{L^2}{D \cdot L} \cdot (\alpha + \alpha^2) \cdot \deg(X) \\ &\geq \frac{D \cdot L}{L^2} \cdot (\alpha - 1) + (1 - \alpha m - 2\alpha g) - \frac{L^2}{D \cdot L} \cdot (\alpha + \alpha^2) \cdot A(D, X) \\ &\geq \alpha \cdot \left(\varphi(\text{mult}(X)) - m\right) \geq 0. \end{aligned}$$

Moreover, we note that due to (3) we have

$$\deg(X : L) = \deg(X) - \deg(X \cap L) \leq \deg(X) - B(D, X) + m.$$

But then

$$\begin{aligned} \deg((X : L) \cap L) &\leq \deg(X \cap L) \leq B(D, X) \\ &= \frac{D \cdot L - L^2}{L^2} - \alpha \cdot \frac{L^2}{D \cdot L - L^2} \cdot (\deg(X) - B(D, X) + m) - 2g - \frac{L^2}{D \cdot L - L^2} \cdot C''(D, X) \\ &\leq \frac{D \cdot L - L^2}{L^2} - \alpha \cdot \frac{L^2}{D \cdot L - L^2} \cdot \deg(X : L) - 2g = B(D - L, X : L), \end{aligned}$$

which gives (4.6).

Step 5: We finally suppose that $X = X(C, T^*)$ satisfies the following conditions:

- (a) $\deg(X \cap L) < B(D, X)$.
- (b) There exists an irreducible branch Q of (C, z) such that
 - (1) $T^* \cap Q = \{z = q_0 < q_1 < \dots < q_l\}$,
 - (2) $T^* \cap L = \{z = q_0 < q_1 < \dots < q_s\}$, $s < l$, and
 - (3) $\text{mult}(C_{(q_{s+1})}, q_{s+1}) = m$ with $\deg(X \cap L) + m < B(D, X)$.

Then by Lemma 4.8 (d) we may specialise the point q_{s+1} onto the curve L and consider the new scheme $X_1 \sim X$ with $\deg(X_1 \cap L) = \deg(X \cap L) + m$. (Note that $X_1 \sim X$ implies that $X_1 \in \mathcal{K}$.) By the Semicontinuity Theorem for h^1 (cf. [Har77] III.12.8) the vanishing of $H^1(\Sigma, \mathcal{J}_{X''/\Sigma}(D))$ for some $X'' \sim X_1$ with $\deg(X'' \cap L) = \deg(X_1 \cap L)$ implies the vanishing of $H^1(\Sigma, \mathcal{J}_{X/\Sigma}(D))$ itself. Thus, specialising points to L we come down to the cases which we have already studied. \square

4.8 Lemma

Let $X = X(C, T^*) \in \mathcal{GS}$, let Q be a smooth branch of (C, z) and let $L \subset \Sigma$ be a curve which is smooth at z . Then:

- (a) For almost all $Y' \sim X : L$ in \mathcal{GS} with $\deg(Y' \cap L) = \deg((X : L) \cap L)$ there exists a generalised singularity scheme $X' \sim X$ in \mathcal{GS} such that $Y' = X' : L$ and $\deg(X' \cap L) = \deg(X \cap L)$.
- (b) If $T^* \cap Q \subseteq T^* \cap L$, then $\text{mult}(X : L) = \text{mult}(X) - 1$.
- (c) $\deg(X) = \sum_{q \in T^*} \frac{m_q \cdot (m_q + 1)}{2}$.
- (d) If $T^* \cap L \subseteq T \subseteq T^* \cap Q$ is a connected subtree, then there exists an analytical isomorphism $\varphi : (\Sigma, z) \rightarrow (\Sigma, z)$ mapping (C, z) to a plane curve singularity (C_1, z) such that the subtree T_1^* corresponding to T^* satisfies:
 - (1) $X_1 = X(C_1, T_1^*) \cong X$,
 - (2) $T_1^* \cap L$ is the subtree of T_1^* corresponding to T , and
 - (3) $T_1^* \cap L = T_1^* \cap Q_1$, where $Q_1 = \varphi(Q)$ is a smooth branch of (C_1, z) .

In particular, $X_1 \sim X$ and $\deg(X_1 \cap L) = \deg(X \cap L) + \sum_{q \in T \setminus (T^* \cap L)} m_q$.

Proof: (a) Cf. [GLS98c] Proposition 2.18 or [Los98] Proposition 2.32.

(b) Cf. [GLS98c] Lemma 2.15 or [Los98] Lemma 2.26.

(c) Cf. [GLS98c] Lemma 2.6 or [Los98] Lemma 2.8.

(d) Cf. [GLS98c] Lemma 2.14 or [Los98] Lemma 2.25.

\square

5. Examples

We are now stating the results which we derive from Theorem 4.5 in view of Remark 4.6. We leave it to the reader to exploit the possibilities, which (4.1) offers, in a better way.

5.a. The Classical Case - $\Sigma = \mathbb{P}_\mathbb{C}^2$

Choosing L as a line in $\mathbb{P}_\mathbb{C}^2$, which has genus $g = g(L) = 0$, we get precisely the same result as in [GLS98c].

4.5a Theorem

Let $d > 1$ be an integer and let $X \in \mathcal{GS}_1$ such that:

$$(4.2a) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot (d - \text{mult}(X))^2,$$

$$(4.3a) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X).$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\mathbb{P}_\mathbb{C}^2, \mathcal{J}_{X'/\Sigma}(dL)) = 0.$$

5.b. Hirzebruch Surfaces

Working now with the very ample divisor $L = C_0 + (e + 1) \cdot L$, which has genus $g = g(L) = 0$, the theorem takes for divisors dL precisely the same form as in the case of $\mathbb{P}_\mathbb{C}^2$. We might as well work with any other very ample divisor L , of possibly larger genus $g = g(L)$. The Theorem then takes the same form as for K3-surfaces.

For the Hirzebruch surface $\mathbb{F}_0 = \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$ we also get another, somewhat more general version.

4.5b Theorem

Let $a, b > 1$ be two integers and let $X \in \mathcal{GS}_1$ such that:

$$(4.1b) \quad a, b \geq \text{mult}(X) - 1,$$

$$(4.2b) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot (d - \text{mult}(X))^2,$$

$$(4.3b) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X).$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1, \mathcal{J}_{X'/\Sigma}(aC_0 + bF)) = 0.$$

5.c. Geometrically Ruled Surfaces

Using the notation of Section G.a $\pi : \Sigma \rightarrow C$ is a geometrically ruled surface with invariant e . As in Remark 4.6 we may choose some integer $l \geq \max\{2g - 1, e + 2g - 1\}$ such that $L = C_0 + lF$ is very ample. Its genus is then just $g(L) = g(C)$.

4.5c Theorem

Let $d > 1$ be an integer and let $X \in \mathcal{GS}_1$ such that

$$(4.2c) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot \left((d - \text{mult}(X))^2 - (2 + \sqrt{2}) \cdot g(C) \cdot d \right),$$

$$(4.3c) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X) - 2g(C).$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(dL)) = 0.$$

5.d. Products of Curves

Let $\Sigma = C_1 \times C_2$ be a product of two smooth projective curves of genera g_1 and g_2 respectively. Then we may choose some very ample divisor $L \sim_{\alpha} (2g_2 + 1) \cdot C_1 + (2g_1 + 1) \cdot C_2$ of genus $g = 8g_1g_2 + g_1 + g_2$.

4.5d Theorem

Let $d > 1$ be an integer and let $X \in \mathcal{GS}_1$ such that

$$(4.2d) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot \left((d - \text{mult}(X))^2 - (2 + \sqrt{2}) \cdot (8g_1g_2 + g_1 + g_2) \cdot d \right),$$

$$(4.3d) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X) - 2 \cdot (8g_1g_2 + g_1 + g_2).$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1\left(C_1 \times C_2, \mathcal{J}_{X'/\Sigma}(d \cdot (2g_2 + 1) \cdot C_1 + d \cdot (2g_1 + 1) \cdot C_2)\right) = 0.$$

5.e. Surfaces in $\mathbb{P}_{\mathbb{C}}^3$

For a surface in $\mathbb{P}_{\mathbb{C}}^3$ of degree n with L a hyperplane section, the formula for the genus $g = g(L)$ comes down to $g = \frac{(n-1) \cdot (n-2)}{2}$.

4.5e Theorem

Let $d > 1$ be an integer and let $X \in \mathcal{GS}_1$ such that

$$(4.2e) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot \left((d - \text{mult}(X))^2 - \frac{2+\sqrt{2}}{2} \cdot (n-1) \cdot (n-2) \cdot d \right),$$

$$(4.3e) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X) - (n-1) \cdot (n-2).$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(dL)) = 0.$$

If Σ was a complete intersection of type (d_1, \dots, d_{N-2}) and L is a hyperplane section, then we just would have to replace the genus $g = g(L) = \frac{d_1 \cdots d_{N-2} \cdot \left(\sum_{i=1}^{N-2} d_i - N \right) + 2}{2}$.

5.f. K3-Surfaces

If Σ is a K3-surface and L any very ample divisor of genus $g(L) = g$, Theorem 4.5 comes down to.

4.5f Theorem

Let $d > 1$ be an integer and let $X \in \mathcal{GS}_1$ such that

$$(4.2f) \quad \deg(X) \leq (3 - 2 \cdot \sqrt{2}) \cdot \left((d - \text{mult}(X))^2 - (2 + \sqrt{2}) \cdot g \cdot d \right),$$

$$(4.3f) \quad \deg(X \cap L) \leq d - \frac{1+\sqrt{2}}{d} \cdot \deg(X) - 2g.$$

Then there is some $X' \in \mathcal{GS}_1$ with $X' \sim X$ and $\deg(X' \cap L) = \deg(X \cap L)$ such that

$$h^1(\Sigma, \mathcal{J}_{X'/\Sigma}(dL)) = 0.$$

CHAPTER III

Existence

The question whether there exist plane curves of given degree having singular points of given types was already studied by the Italian geometers during the last century. A very nice and inspiring result in this direction is that of Severi (cf. [Sev21]) which says that there is an irreducible plane curve of degree d with r nodes as only singularities if and only if $0 \leq r \leq \frac{(d-1) \cdot (d-2)}{2}$. In our terminology this means

$$V_{|d \cdot H|}^{\text{irr}}(rA_1) \neq \emptyset \iff 0 \leq r \leq \frac{(d-1) \cdot (d-2)}{2},$$

where H is a line in \mathbb{P}_c^2 . Severi deduced this by showing that nodes on plane curves can be “smoothed” independently.

Looking for generalisations of this result one might concentrate on different aspects such as

- looking at ordinary multiple points of higher multiplicities on plane curves, or
- looking at arbitrary singularities on plane curves, or
- looking at nodal curves on arbitrary surfaces, or, finally,
- looking at arbitrary singularities on arbitrary surfaces.

All these generalisations have one problem in common – we may hardly expect criteria which are necessary *and* sufficient for the existence at the same time. Already when replacing “nodes” by “cusps” on plane curves there is no such complete answer known – except for small degrees d , $d \leq 10$ –, and the existence of irreducible plane curves with more cusps than the number of conditions imposed by them should allow shows that one may hardly expect such a result (cf. [Hir92]). Similar problems arise in the other situations, and the reason is mainly that in general the conditions which the singular points impose are not independent.

The direction of looking at plane curves with ordinary multiple points of arbitrary multiplicities was very much inspired by a conjecture of Nagata who proposed in [Nag59] that $\sqrt{r} \cdot \sum_{i=1}^r m_i$ should be a lower bound for the degree of a plane curve passing through r general points with multiplicities m_1, \dots, m_r , whenever $r \geq 9$. He himself gave a complete answer for the case $r \leq 8$ and when r is a square. Many efforts have been taken to fully prove the conjecture and much progress has been made – cf. [Nag59, Seg62, ArC81, Hir85, Har86, Gim87, Hir89, Ran89, CiM98,

Sei99, Bru99, Eva99, Mig00, AlH00, Roé01, Mig01]. However, a full proof is still missing, though several special cases have been cracked and the generally known bounds have been improved largely. We refer to **[Cil01]** and **[Har01]** for an overview on the state of the art for the Nagata Conjecture and for the related conjectures of Segre **[Seg62]**, Gimigliano **[Gim87]** and of Harbourne–Hirschowitz **[Har85a, Hir89]**.

Also for the existence of plane curves with one or with several singularities of given topological singularity types there is a large number of results, giving necessary or sufficient conditions for the existence or just giving strange examples – cf. **[GTU66, ArC81, Bru81, ArC83, GuN83, Var83, Ura83, Tan84, Ura84, GrM85, Ivi85, Koe86, Hir86, Ura86, Deg87, Shu87a, GrM88, Hir92, Bar93b, Sak93, Shu93, Deg95, Wal95, Wal96, GLS98c, Los98, Shu98, Shu99, Mig01, Shu01]**. For an overview on the results known we refer to **[Los98]**, **[GrS99]** or **[GLS02]**. Just recently the asymptotically best known conditions for the existence of curves in the plane with arbitrary topological singularities have been considerably improved using Castelnuovo function arguments, and at the same time the first general conditions for analytical singularities have been found (cf. **[Shu01]**). The existence of an irreducible plane curve of degree d with topological singularity types $\mathcal{S}_1, \dots, \mathcal{S}_r$, among them k nodes, m cusps and t singularities of type A_{2l} with $l \geq 2$, as only singular points is ensured by

$$6k + 10m + \frac{49}{6} \cdot t + \frac{625}{48} \cdot \sum_{\mathcal{S}_i \neq A_1, A_2} \delta(\mathcal{S}_i) \leq d^2 - 2d + 3,$$

where $\delta(\mathcal{S}_i)$ is the delta-invariant of \mathcal{S}_i , and similarly for analytical singularity types

$$6k + 10m + 9 \cdot \sum_{\mathcal{S}_i \neq A_1, A_2} \mu(\mathcal{S}_i) \leq d^2 - 2d + 3,$$

suffices, where $\mu(\mathcal{S}_i)$ is the Milnor number of \mathcal{S}_i . (Cf. **[Shu01]**.) We will compare this particular result to the corresponding special case of our general results in Section 3.a and see that our results are weaker – which was to be expected, since the techniques applied in **[Shu01]** are optimized for the plane case.

In the past the question of the existence of curves with prescribed analytical singularity types has attained much less attention, due to the lack of suitable methods to tackle the problem. However, the Viro method which Shustin uses in **[Shu99]** Section 6 on hypersurfaces in $\mathbb{P}_{\mathbb{C}}^n$ and which we also use here allows to glue topological singularities as well as analytical ones, so that our main results are valid for both kinds of singularity types. (See also **[Shu01]**.)

The basic idea may be described as follows: suppose we have a “suitable” irreducible curve $C \in |D|_l$ with ordinary multiple points z_1, \dots, z_r of multiplicities m_1, \dots, m_r as singularities, and suppose we have “good” affine representatives g_1, \dots, g_r for the singularity types $\mathcal{S}_1, \dots, \mathcal{S}_r$, then we may glue locally

at the z_i the equations g_i into the curve C . In Definition 2.1 we define what it precisely means that a representative is “good”; that the curve C then is “suitable”, just means that the m_i -jet of C at z_i coincides with the m_i -jet of g_i , where $m_i = \deg(g_i)$, and that $V_{|D|}(m_1, \dots, m_r)$ is smooth at C of the expected dimension. Finding general conditions for the existence thus comes down to finding conditions for the existence of curves with ordinary multiple points satisfying the T-smoothness property and to finding upper bounds for the degree of a good representative. Section 1 is devoted to the first problem, and we get satisfactory results in Theorem 1.2 and Corollary 1.3, while for the second problem we are bound to known results (cf. [Shu01]).

The most restrictive among the *sufficient* conditions which we find in Corollary 2.5 is (2.7) respectively (2.11) and could be characterised as a condition of the type

$$\sum_{i=1}^r \delta(\mathcal{S}_i) \leq \alpha D^2 + \beta D \cdot K + \gamma,$$

for topological singularity types, respectively for analytical ones as

$$\sum_{i=1}^r \mu(\mathcal{S}_i) \leq \alpha D^2 + \beta D \cdot K + \gamma,$$

where K is some fixed divisor class and α, β and γ are some absolute constants. (See also Remark II.1.3.) However, it seems that this condition is of the right type and with the right “exponents” for the invariants of the \mathcal{S}_i as well as for the divisor D , since there are also *necessary* conditions of this type, e. g.

$$\sum_{i=1}^r \mu(\mathcal{S}_i) \leq 2 \cdot \sum_{i=1}^r \delta(\mathcal{S}_i) \leq D^2 + D \cdot K_\Sigma + 2,$$

which follows from the genus formula: If D is an irreducible curve with precisely r singular points of topological or analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ and $\nu : \tilde{D} \rightarrow D$ its normalisation, then $p_a(D) = g(\tilde{D}) + \delta(D) \geq \delta(D)$, where $\delta(D) = \dim_{\mathbb{C}}(\nu_* \mathcal{O}_{\tilde{D}} / \mathcal{O}_D)$ is the delta invariant of D (cf. [BPV84] II.11). Moreover, by definition $\delta(D) = \sum_{z \in \text{Sing}(D)} \delta(D, z)$, and it is well known that $2\delta(D, z) = \mu(D, z) + r(D, z) - 1 \geq \mu(D, z)$ (see Remark I.2.2). Using now the genus formula we get: $D^2 + D \cdot K_\Sigma + 2 = 2p_a(D) \geq 2 \cdot \sum_{i=1}^r \delta(\mathcal{S}_i) \geq \sum_{i=1}^r \mu(\mathcal{S}_i)$.

1. Existence Theorem for Ordinary Fat Point Schemes

In order to be able to apply the existence theorem for ordinary fat point schemes to the general case it is important that the existing curve has the T-smoothness property, that is that some H^1 vanishes. This vanishing is ensured by reducing the problem to a “lower degree” and applying the following lemma as a kind of induction step.

1.1 Lemma

Let $\mathbb{L} \subset \Sigma$ be a smooth curve and $X \subset \Sigma$ a zero-dimensional scheme. If $D \in \text{Div}(\Sigma)$ such that

$$(1.1) \quad h^1(\Sigma, \mathcal{J}_{X:L/\Sigma}(D - L)) = 0, \text{ and}$$

$$(1.2) \quad \deg(X \cap L) \leq D \cdot L + 1 - 2g(L),$$

then

$$h^1(\Sigma, \mathcal{J}_{X/\Sigma}(D)) = 0.$$

Proof: Condition (1.2) implies

$$2g(L) - 2 < D \cdot L - \deg(X \cap L) = \deg(\mathcal{O}_L(D)) + \deg(\mathcal{J}_{X \cap L/L}) = \deg(\mathcal{J}_{X \cap L/L}(D)),$$

and thus by Riemann-Roch (cf. [Har77] IV.1.3.4)

$$h^1(\mathcal{J}_{X \cap L/L}(D)) = 0.$$

Consider now the exact sequence

$$0 \longrightarrow \mathcal{J}_{X:L/\Sigma}(D - L) \xrightarrow{\cdot L} \mathcal{J}_{X/\Sigma}(D) \longrightarrow \mathcal{J}_{X \cap L/L}(D) \longrightarrow 0.$$

The result then follows from the corresponding long exact cohomology sequence

$$0 = H^1(\mathcal{J}_{X:L/\Sigma}(D - L)) \longrightarrow H^1(\mathcal{J}_{X/\Sigma}(D)) \longrightarrow H^1(\mathcal{J}_{X \cap L/L}(D)) = 0.$$

□

1.2 Theorem

Given $m_1, \dots, m_r \in \mathbb{N}$, not all zero, and $z_1, \dots, z_r \in \Sigma$, in very general position.

Let $L \in \text{Div}(\Sigma)$ be very ample over \mathbb{C} , and let $D \in \text{Div}(\Sigma)$ be such that

$$(1.3) \quad h^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D - L)) = 0, \text{ and}$$

$$(1.4) \quad D \cdot L - 2g(L) \geq m_i + 1 \text{ for all } i = 1, \dots, r.$$

Then there exists an irreducible curve $C \in |D|_L$ with ordinary singular points of multiplicity m_i at z_i for $i = 1, \dots, r$ and no other singular points. Furthermore,

$$h^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0,$$

and in particular, $V_{|D|}(\underline{m})$ is T -smooth at C .

Addendum: Given local coordinates x_j, y_j in z_j and open dense subsets $U_j \subseteq \mathbb{C}[x_j, y_j]_{m_j}$,¹ $j = 1, \dots, r$, the curve C may be chosen such that the m_j -jet at z_j of a section defining C belongs to U_j .

Idea of the proof: For each z_j find a curve $C_j \in |\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)|_L$ with an ordinary singular point of multiplicity m_j and show that this linear system has no other base points than z_1, \dots, z_r . Then the generic element is smooth outside z_1, \dots, z_r and has an ordinary singularity of multiplicity m_j in z_j for $j = 1, \dots, r$.

¹For the definition of $\mathbb{C}[x, y]_d$ see Page 44.

Proof: Let $z_0 \in \Sigma$ be a generic point in the sense that for any $z^* \in \Sigma$ there is a smooth connected curve, say L_{z^*} , in $|L|_1$ through z_0 and z^* containing at most one of the z_i , $i = 1, \dots, r$. (Cf. Lemma E.3.)

If $Z \subseteq \Sigma$ is any zero-dimensional subscheme and $z^* \in \Sigma$ is any point, we define the zero-dimensional scheme $Z \cup \{z^*\}$ by the ideal sheaf

$$\mathcal{I}_{Z \cup \{z^*\}/\Sigma, z} = \begin{cases} \mathcal{I}_{Z/\Sigma, z}, & \text{if } z \neq z^*, \\ \mathfrak{m}_{\Sigma, z^*} \cdot \mathcal{I}_{Z/\Sigma, z^*}, & \text{if } z = z^*. \end{cases}$$

Writing X for $X(\underline{m}; \underline{z})$ we introduce in particular the zero-dimensional schemes $X_j = X \cup \{z_j\}$ for $j = 0, \dots, r$.

Step 1: $h^1(\mathcal{I}_{X_0 \cup \{z^*\}/\Sigma}(D)) = h^1(\mathcal{I}_{X_j/\Sigma}(D)) = 0$ for $z^* \in \Sigma \setminus \{z_0\}$ and $j = 0, \dots, r$.

Since L_{z^*} passes through z_0, z^* and at most one z_j , $j \in \{1, \dots, r\}$, where z^* might be this z_j , Condition (1.4) implies

$$\deg(X_0 \cap L_{z^*}) \leq m_j + 2 \leq D.L + 1 - 2g(L). \quad (1.5)$$

Moreover, if $z^* = z_j$ for some $j \in \{1, \dots, r\}$ or if L_{z^*} does not pass through any z_j , $j \in \{1, \dots, r\}$, then $X_0 \cup \{z^*\} : L_{z^*} = X$, otherwise we get the exact sequence

$$0 \longrightarrow \mathcal{I}_{X/\Sigma} \longrightarrow \mathcal{I}_{X_0 \cup \{z^*\}; L_{z^*}/\Sigma} \longrightarrow \mathfrak{m}_{\Sigma, z_j}^{m_j-1} / \mathfrak{m}_{\Sigma, z_j}^{m_j} \longrightarrow 0,$$

but in any case we have by Condition (1.3) that

$$h^1(\mathcal{I}_{X_0 \cup \{z^*\}; L_{z^*}/\Sigma}(D - L)) = 0. \quad (1.6)$$

(1.5) and (1.6) allow us to apply Lemma 1.1 in order to obtain

$$h^1(\mathcal{I}_{X_0 \cup \{z^*\}/\Sigma}(D)) = 0.$$

The inclusion $\mathcal{I}_{X_0 \cup \{z_j\}/\Sigma} \hookrightarrow \mathcal{I}_{X_j/\Sigma}$, for $j = 1, \dots, r$, respectively the inclusion $\mathcal{I}_{X_0 \cup \{z^*\}/\Sigma} \hookrightarrow \mathcal{I}_{X_0/\Sigma}$, for some $z^* \neq z_0$, then imply that for any $j = 0, \dots, r$ also

$$h^1(\mathcal{I}_{X_j/\Sigma}(D)) = 0.$$

Step 2: For each $j = 1, \dots, r$ there exists a curve $C_j \in |D|_1$ with an ordinary singular point of multiplicity m_j at z_j and with $\text{mult}_{z_i}(C_j) \geq m_i$ for $1 \leq i \neq j$.

Consider the exact sequence

$$0 \longrightarrow \mathcal{I}_{X_j/\Sigma} \longrightarrow \mathcal{I}_{X/\Sigma} \longrightarrow \mathfrak{m}_{\Sigma, z_j}^{m_j} / \mathfrak{m}_{\Sigma, z_j}^{m_j+1} \longrightarrow 0$$

twisted by D and the corresponding long exact cohomology sequence

$$H^0(\mathcal{I}_{X/\Sigma}(D)) \rightarrow \mathfrak{m}_{\Sigma, z_j}^{m_j} / \mathfrak{m}_{\Sigma, z_j}^{m_j+1} \rightarrow H^1(\mathcal{I}_{X_j/\Sigma}(D)) \rightarrow H^1(\mathcal{I}_{X/\Sigma}(D)) \rightarrow 0. \quad (1.7)$$

\parallel Step 1
 0

Thus we may choose the C_j to be given by a section in $H^0(\mathcal{J}_{X/\Sigma}(D))$ where the m_j tangent directions at z_j are all different – *indeed, we may choose the m_j -jet at z_j of the section defining C_j arbitrary!*

Step 3: The base locus of $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ is $\{z_0, \dots, z_r\}$.

Suppose $z^* \in \Sigma$ was an additional base point of $|\mathcal{J}_{X_0/\Sigma}(D)|_1$. By Step 1

$$h^1(\mathcal{J}_{X_0 \cup \{z^*\}/\Sigma}(D)) = 0,$$

and thus as in Step 2

$$h^0(\mathcal{J}_{X_0/\Sigma}(D)) = h^0(\mathcal{J}_{X_0 \cup \{z^*\}/\Sigma}(D)) + 1.$$

But by assumption z^* is a base point, and thus

$$h^0(\mathcal{J}_{X_0/\Sigma}(D)) = h^0(\mathcal{J}_{X_0 \cup \{z^*\}/\Sigma}(D)),$$

which gives us the desired contradiction.

Step 4: The base locus of $|\mathcal{J}_{X/\Sigma}(D)|_1$ is $\{z_1, \dots, z_r\}$.

By Step 3 and since $|\mathcal{J}_{X_0/\Sigma}(D)|_1 \subset |\mathcal{J}_{X/\Sigma}(D)|_1$ we know that the base locus is contained in $\{z_0, \dots, z_r\}$.

Let now $z_0^* \in \Sigma$ be a second generic point like z_0 , then we see with the same arguments that the base locus $|\mathcal{J}_{X/\Sigma}(D)|_1$ is contained in $\{z_0^*, z_1, \dots, z_r\}$, and thus it is $\{z_1, \dots, z_r\}$.

Step 5: There is an irreducible curve $C \in |\mathcal{J}_{X/\Sigma}(D)|_1$, which has an ordinary singular point of multiplicity m_i at z_i for $i = 1, \dots, r$ and no other singular points – indeed, there is an open and dense subset U of curves in $|\mathcal{J}_{X/\Sigma}(D)|_1$ satisfying these requirements.

Because of Step 2 the generic element in $|\mathcal{J}_{X/\Sigma}(D)|_1$ has an ordinary singular point of multiplicity m_i at z_i and is by Bertini's Theorem (cf. [Har77] III.10.9.2) smooth outside its base locus. It remains to show that $|\mathcal{J}_{X/\Sigma}(D)|_1$ contains an irreducible curve, then the generic curve is irreducible as well.

Suppose therefore that all curves in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ are reducible. Then by a Theorem of Bertini (cf. Theorem E.5) there is a one-dimensional family \mathcal{F} of curves such that the irreducible components of any curve in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ belong to \mathcal{F} . Since any curve in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ passes through z_0 , any curve in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ has an irreducible component passing through z_0 . By Step 3 the linear system has no fixed component, and thus each curve of \mathcal{F} must pass through z_0 . By assumption each curve in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ has several components and now each of these components passes through z_0 . Thus, each curve in $|\mathcal{J}_{X_0/\Sigma}(D)|_1$ is singular in z_0 . However, since z_0 is generic there will be a generic curve in $|\mathcal{J}_{X/\Sigma}(D)|_1$ through z_0 , which then has to be smooth in z_0 , as we have seen above. This gives the desired contradiction.

Step 6: $h^1(\mathcal{J}_{X/\Sigma}(D)) = 0$, which follows immediately from equation (1.7).

Step 7: $V_{|\mathcal{D}|}(\underline{m})$ is T -smooth at C .

By [GLS98c], Lemma 2.7, we have

$$\mathcal{J}_{X/\Sigma} \subseteq \mathcal{J}_{X^{es}(C)/\Sigma},$$

and thus by Step 6

$$h^1(\mathcal{J}_{X^{es}(C)/\Sigma}(D)) = 0,$$

which proves the claim.

Step 8: Prove the addendum.

Fixing local coordinates x_j, y_j at z_j we may consider the *linear* maps

$$\varphi_j : H^0(\Sigma, \mathcal{J}_{X/\Sigma}(D)) \rightarrow \mathbb{C}[x_j, y_j]_{m_j} : s \mapsto \text{jet}_{m_j}(s_{z_j}),$$

where $\text{jet}_{m_j}(s_{z_j})$ denotes the m_j -jet of the section s in the local coordinates at z_j . According to the considerations in Step 2 we know that the φ_j are indeed surjective. Thus the sets $\varphi_j^{-1}(U_j)$ are open dense subsets of $H^0(\Sigma, \mathcal{J}_{X/\Sigma}(D))$, and if $V \subset H^0(\Sigma, \mathcal{J}_{X/\Sigma}(D))$ denotes the open dense subset of sections corresponding to the curves in the set U from Step 5, then $W = V \cap \bigcap_{j=1}^r \varphi_j^{-1}(U_j)$ is non-empty and a curve defined by a section in W satisfies the requirements of the addendum. \square

1.3 Corollary

Let $m_1, \dots, m_r \in \mathbb{N}_0$, not all zero, and let $L \in \text{Div}(\Sigma)$ be very ample over \mathbb{C} . Suppose $D \in \text{Div}(\Sigma)$ such that

$$(1.8) \quad (D - L - K_\Sigma)^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.9) \quad (D - L - K_\Sigma) \cdot B > \max\{m_1, \dots, m_r\} \quad \text{for any irreducible curve } B \subset \Sigma \\ \text{with } B^2 = 0 \text{ and } \dim |B|_a \geq 1,$$

$$(1.10) \quad D \cdot L - 2g(L) > \max\{m_1, \dots, m_r\}, \text{ and}$$

$$(1.11) \quad D - L - K_\Sigma \text{ is nef.}$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position there exists an irreducible curve $C \in |\mathcal{D}|_1$ with ordinary singular points of multiplicity m_i at z_i for $i = 1, \dots, r$ and no other singular points. Furthermore,

$$h^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0,$$

and in particular, $V_{|\mathcal{D}|}(\underline{m})$ is T -smooth in C .

Addendum: Given local coordinates x_j, y_j in z_j and open dense subsets $U_j \subseteq \mathbb{C}[x_j, y_j]_{m_j}$, $j = 1, \dots, r$, the curve C may be chosen such that the m_j -jet at z_j of a section defining C belongs to U_j .

Proof: Follows from Theorem 1.2 and Corollary II.1.2. \square

2. Existence Theorem for General Singularity Schemes

Notation

Throughout this section we will be concerned with equisingular respectively equianalytical families of curves. We are treating these two cases simultaneously, using the suffixes “es” in the equisingular case, that is whenever topological singularity types are considered, “ea” in the equianalytical case, that is when analytical singularity types are considered, and finally, using “*” when both cases are considered at once.

In the following we will denote by $\mathbb{C}[x, y]_d$, respectively by $\mathbb{C}[x, y]_{\leq d}$ the \mathbb{C} -vector spaces of polynomials of degree d , respectively of degree at most d . If $f \in \mathbb{C}[x, y]_{\leq d}$ we denote by $f_k \in \mathbb{C}[x, y]_k$ for $k = 0, \dots, d$ the homogeneous part of degree k of f , and thus $f = \sum_{k=0}^d f_k$. By $\underline{a} = (a_{k,l} \mid 0 \leq k+l \leq d)$ we will denote the coordinates of $\mathbb{C}[x, y]_{\leq d}$ with respect to the basis $\{x^k y^l \mid 0 \leq k+l \leq d\}$.

For any $f \in \mathbb{C}[x, y]_{\leq d}$ the tautological family

$$\mathbb{C}[x, y]_{\leq d} \times \mathbb{C}^2 \supset \bigcup_{g \in \mathbb{C}[x, y]_{\leq d}} \{g\} \times g^{-1}(0) \longrightarrow \mathbb{C}[x, y]_{\leq d}$$

induces a deformation of the plane curve singularity $(f^{-1}(0), 0)$ whose base space is the germ $(\mathbb{C}[x, y]_{\leq d}, f)$ of $\mathbb{C}[x, y]_{\leq d}$ at f . Given any deformation $(X, x) \hookrightarrow (\mathcal{X}, x) \rightarrow (S, s)$ of a plane curve singularity (X, x) , we will denote by $S^* = (S^*, s)$ the germ of the equisingular respectively equianalytical stratum of (S, s) .² Thus, fixed an $f \in \mathbb{C}[x, y]_{\leq d}$, $\mathbb{C}[x, y]_{\leq d}^* = (\mathbb{C}[x, y]_{\leq d}^*, f)$ is the (local) equisingular respectively equianalytical stratum of $\mathbb{C}[x, y]_{\leq d}$ at f .

2.1 Definition

- (a) We say the family $\mathbb{C}[x, y]_{\leq d}$ is topologically respectively analytically *T-smooth* at $f \in \mathbb{C}[x, y]_{\leq d}$ if for any $e \geq d$ there exists a $\Lambda \subset \{(k, l) \in \mathbb{N}_0^2 \mid 0 \leq k+l \leq d\}$ with $\#\Lambda = \tau^* = \tau^*(f^{-1}(0), 0)$ such that $\mathbb{C}[x, y]_{\leq e}^*$ is given by equations

$$a_{k,l} = \phi_{k,l}(\underline{a}_{(1)}, \underline{a}_{(2)}), \quad (k, l) \in \Lambda,$$

with $\phi_{k,l} \in \mathbb{C}\{\underline{a}_{(1)}, \underline{a}_{(2)}\}$ where $\underline{a}_{(0)} = (a_{k,l} \mid (k, l) \in \Lambda)$, $\underline{a}_{(1)} = (a_{k,l} \mid 0 \leq k+l \leq d, (k, l) \notin \Lambda)$, and $\underline{a}_{(2)} = (a_{k,l} \mid d+1 \leq k+l \leq e)$.

- (b) A polynomial $f \in \mathbb{C}[x, y]_{\leq d}$ is said to be a *good representative* of the topological respectively analytical singularity type \mathcal{S} in $\mathbb{C}[x, y]_{\leq d}$ if it meets the following conditions:
- $\text{Sing}(f^{-1}(0)) = \{p \in \mathbb{C}^2 \mid f(p) = 0, \frac{\partial f}{\partial x}(p) = 0, \frac{\partial f}{\partial y}(p) = 0\} = \{0\}$,
 - $(f^{-1}(0), 0) \sim_t \mathcal{S}$ respectively $(f^{-1}(0), 0) \sim_c \mathcal{S}$,
 - $\mathbb{C}[x, y]_{\leq d}$ is T-smooth at f , and

²That is, S^* is the analytical space germ parametrising the subfamily of $(\mathcal{X}, x) \rightarrow (S, s)$ of singularities which are topologically respectively analytically equivalent to (X, x) .

- (d) f_d is a generic reduced d -form, that is, there is an open subset $U \subseteq \mathbb{C}[x, y]_d$ of reduced d -forms such that for each $g_d \in U$ there is an $f \in \mathbb{C}[x, y]_{\leq d}$ satisfying (a)-(c) and $f_d = g_d$.
- (c) Given a topological respectively analytical singularity type \mathcal{S} we define $e^s(\mathcal{S})$ respectively $e^a(\mathcal{S})$ to be the minimal number d such that \mathcal{S} has a good representative of degree d .

Whenever we consider topological and analytical singularity types at the same time, we will use $e^*(\mathcal{S})$ to denote $e^s(\mathcal{S})$ and $e^a(\mathcal{S})$ respectively.

2.2 Remark

- (a) The condition for T-smoothness just means that for any $e \geq d$ the equisingular respectively equianalytical stratum $\mathbb{C}[x, y]_{\leq e}^*$ is smooth at the point f of the expected codimension in $(\mathbb{C}[x, y]_{\leq e}, f)$.
- (b) Note that for a polynomial of degree d the highest homogeneous part f_d defines the normal cone, i. e. the intersection of the curve $\{\hat{f} = 0\}$ with the line at infinity in $\mathbb{P}_\mathbb{C}^2$, where \hat{f} is the homogenisation of f . Thus the condition “ f_d reduced” just means that the line at infinity intersects the curve transversally in d different points.
- (c) If $f \in \mathbb{C}[x, y, z]_d$ is an irreducible polynomial such that $(0 : 0 : 1)$ is the only singular point of the plane curve $\{f = 0\} \subset \mathbb{P}_\mathbb{C}^2$, then a linear change of coordinates of the type $(x, y, z) \mapsto (x, y, z + ax + by)$ will ensure that the dehomogenisation \check{f} of f satisfies “ \check{f}_d reduced”. Note for this that the coordinate change corresponds to choosing a line in $\mathbb{P}_\mathbb{C}^2$, not passing through $(0 : 0 : 1)$ and meeting the curve in d distinct points. Therefore, the bounds for $e^s(\mathcal{S})$ and $e^a(\mathcal{S})$ given in [Shu01] do apply here.
- (d) In order to obtain good numerical conditions for the existence it is vital to find good upper bounds for $e^s(\mathcal{S})$ and $e^a(\mathcal{S})$. We gather here the best known results in this direction obtained by Eugenii Shustin in [Shu01] (see also [GLS02] Chapter V.4).

If \mathcal{S} is a simple singularity, then of course $e^s(\mathcal{S}) = e^a(\mathcal{S})$.

\mathcal{S}	$e^s(\mathcal{S}) = e^a(\mathcal{S})$	\mathcal{S}	$e^s(\mathcal{S}) = e^a(\mathcal{S})$
A_1	2	D_4	3
A_2	3	D_5	4
$A_\mu, \mu = 3, \dots, 7$	4	$D_\mu, \mu \leq 6, \dots, 10$	5
$A_\mu, \mu = 8, \dots, 10$	5	$D_\mu, \mu \leq 11, \dots, 13$	6
$A_\mu, \mu \geq 1$	$\leq 2 \cdot \lfloor \sqrt{\mu + 5} \rfloor$	$D_\mu, \mu \geq 1$	$\leq 2 \cdot \lfloor \sqrt{\mu + 7} \rfloor$

\mathcal{S}	$e^s(\mathcal{S}) = e^a(\mathcal{S})$	\mathcal{S}	$e^s(\mathcal{S}) = e^a(\mathcal{S})$	\mathcal{S}	$e^s(\mathcal{S}) = e^a(\mathcal{S})$
E_6	4	E_7	4	E_8	5

For non-simple singularities the invariants for $e^s(\mathcal{S})$ and $e^a(\mathcal{S})$ differ in general. If $\mathcal{S} \notin \{A_1, A_2\}$ is a topological singularity type, then

$$e^s(\mathcal{S}) \leq \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S})} - 1,$$

and if $\mathcal{S} \notin \{A_1, A_2\}$ is an analytical singularity type, then

$$e^a(\mathcal{S}) \leq 3 \cdot \sqrt{\mu(\mathcal{S})} - 1.$$

Finally, in Lemma V.3.10 we are going to show that for an arbitrary topological or analytical singularity type we have the inequality

$$e^*(\mathcal{S}) \leq \tau^*(\mathcal{S}) + 1.$$

- (e) For refined results using the techniques of the following proof we refer to [Shu99].

2.3 Theorem (Existence)

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, and suppose there exists an irreducible curve $C \in |D|_1$ with $\text{Sing}(C) = \{z_1, \dots, z_r\}$ such that in local coordinates at z_i the m_i -jet of a section defining C coincides with the m_i -jet of a good representative of \mathcal{S}_i ,³ where $m_i = e^s(\mathcal{S}_i)$ respectively $m_i = e^a(\mathcal{S}_i)$, $i = 1, \dots, r$, and

$$h^1(\Sigma, \mathcal{J}_{X(\underline{m}; z)/\Sigma}(D)) = 0.$$

Then there exists an irreducible curve $\tilde{C} \in |D|_1$ with precisely r singular points of types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

Idea of the proof: The basic idea is to glue locally at the z_i equations of good representatives for the \mathcal{S}_i into the curve C . Let us now explain more detailed what we mean by this.

If $g_i = \sum_{k+l=0}^{m_i} a_{k,l}^{i,\text{fix}} x_i^k y_i^l$, $i = 1, \dots, r$, are good representatives of the \mathcal{S}_i , then we are looking for a family F_t , $t \in (\mathbb{C}, 0)$, in $H^0(\Sigma, \mathcal{O}_\Sigma(D))$ which in local coordinates x_i, y_i at z_i looks like

$$F_t^i = \sum_{k+l=0}^{m_i-1} t^{m_i-k-l} \tilde{a}_{k,l}^i(t) \cdot x_i^k y_i^l + \sum_{k+l=m_i} (a_{k,l}^{i,\text{fix}} + t \cdot \text{h.o.t.}) \cdot x_i^k y_i^l + \text{h.o.t.},$$

where the $\tilde{a}_{k,l}^i(t)$ should be convergent power series in t with $\tilde{a}_{k,l}^i(0) = a_{k,l}^{i,\text{fix}}$. The curve defined by F_0 will just be the curve C , while F_t^i for $t \neq 0$ can be transformed, by $(x_i, y_i) \mapsto (tx_i, ty_i)$, into a member of some family

$$\tilde{F}_t^i = \sum_{k+l=0}^{m_i-1} \tilde{a}_{k,l}^i(t) \cdot x_i^k y_i^l + \sum_{k+l=m_i} (a_{k,l}^{i,\text{fix}} + t \cdot \text{h.o.t.}) \cdot x_i^k y_i^l + \text{h.o.t.}, \quad t \in \mathbb{C},$$

³This implies in particular that C has an ordinary m_i -fold point.

with

$$\tilde{F}_0^i = g_i.$$

Using now the T-smoothness property of g_i , $i = 1, \dots, r$, we can choose the $\tilde{a}_{k,l}^i(t)$ such that this family is equisingular respectively equianalytical. Hence, for small $t \neq 0$, the curve given by F_t will have the right singularities at the z_i . Finally, the knowledge on the singularities of C and the Conservation of Milnor numbers will ensure that the curve given by F_t has no further singularities, for $t \neq 0$ sufficiently small.

The proof will be done in several steps. First of all we are going to fix some notation by choosing good representatives for the \mathcal{S}_i (Step 1.1) and by choosing a basis of $H^0(\Sigma, \mathcal{O}_\Sigma(D))$ which reflects the “independence” of the coordinates at the different z_i ensured by $h^1(\Sigma, \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0$ (Step 1.1). In a second step we are making an “Ansatz” for the family F_t , and, for the local investigation of the singularity type, we are switching to some other families \tilde{F}_t^i , $i = 1, \dots, r$ (Step 2.1). We, then, reduce the problem of F_t , for $t \neq 0$ small, having the right singularities to a question about the equisingular respectively equianalytical strata of some families of polynomials (Step 2.2), which in Step 2.3 will be solved. The final step serves to show that the curves F_t have only the singularities which we controlled in the previous steps.

Proof:

Step 1.1: By the definition of $m_i = e^s(\mathcal{S}_i)$ respectively $m_i = e^a(\mathcal{S}_i)$, we may choose good representatives

$$g_i = \sum_{k+l=0}^{m_i} a_{k,l}^{i, \text{fix}} x_i^k y_i^l \in \mathbb{C}[x_i, y_i]_{\leq m_i}$$

for the \mathcal{S}_i , $i = 1, \dots, r$. Let $\underline{a}^{i, \text{fix}} = (a_{k,l}^{i, \text{fix}} \mid 0 \leq k+l \leq m_i)$ and $\underline{a}^{\text{fix}} = (\underline{a}^{1, \text{fix}}, \dots, \underline{a}^{r, \text{fix}})$.

Step 1.2: Parametrise $|D|_i = \mathbb{P}(H^0(\mathcal{O}_\Sigma(D)))$.

Consider the following exact sequence:

$$0 \longrightarrow \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D) \longrightarrow \mathcal{O}_\Sigma(D) \longrightarrow \bigoplus_{i=1}^r \mathcal{O}_{\Sigma, z_i} / \mathfrak{m}_{\Sigma, z_i}^{m_i} \longrightarrow 0.$$

Since $h^1(\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)) = 0$, the long exact cohomology sequence gives

$$H^0(\mathcal{O}_\Sigma(D)) = \bigoplus_{i=1}^r \mathbb{C}\{x_i, y_i\} / (x_i, y_i)^{m_i} \oplus H^0(\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D)),$$

where x_i, y_i are local coordinates of (Σ, z_i) .

We, therefore, can find a basis $\{s_{k,l}^i, s_j \mid j = 1, \dots, e, i = 1, \dots, r, 0 \leq k+l \leq m_i - 1\}$ of $H^0(\mathcal{O}_\Sigma(D))$, with $e = h^0(\mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma}(D))$, such that⁴

⁴Throughout this proof we will use the multi index notation $\alpha = (\alpha_1, \alpha_2) \in \mathbb{N}^2$ and $|\alpha| = \alpha_1 + \alpha_2$.

(1) C is the curve defined by s_1 ,

(2) $(s_1)_{z_i} = \sum_{|\alpha| \geq m_i} B_\alpha^i x_i^{\alpha_1} y_i^{\alpha_2}$ for $i = 1, \dots, r$, with $B_\alpha^i = a_{\alpha_1, \alpha_2}^{i, \text{fix}}$ when $|\alpha| = m_i$,

$$(3) \quad (s_{k,l}^j)_{z_i} = \begin{cases} x_i^k y_i^l + \sum_{|\alpha| \geq m_i} A_{\alpha, k, l}^{j, i} x_i^{\alpha_1} y_i^{\alpha_2}, & \text{if } i = j, \\ \sum_{|\alpha| \geq m_i} A_{\alpha, k, l}^{j, i} x_i^{\alpha_1} y_i^{\alpha_2}, & \text{if } i \neq j. \end{cases}$$

Let us now denote the coordinates of $H^0(\mathcal{O}_\Sigma(D))$ w. r. t. this basis by $(\underline{a}, \underline{b}) = (\underline{a}^1, \dots, \underline{a}^r, \underline{b})$ with $\underline{a}^i = (a_{k,l}^i \mid 0 \leq k+l \leq m_i - 1)$ and $\underline{b} = (b_j \mid j = 1, \dots, e)$.

Thus the family

$$F_{(\underline{a}, \underline{b})} = \sum_{i=1}^r \sum_{k+l=0}^{m_i-1} a_{k,l}^i s_{k,l}^i + \sum_{j=1}^e b_j s_j, \quad (\underline{a}, \underline{b}) \in \mathbb{C}^N \text{ with } N = e + \sum_{i=1}^r \binom{m_i+1}{2},$$

parametrises $H^0(\mathcal{O}_\Sigma(D))$.

Step 2: We are going to glue the good representatives for the \mathcal{S}_i into the curve C . More precisely, we are constructing a subfamily F_t , $t \in (\mathbb{C}, 0)$, in $H^0(\mathcal{O}_\Sigma(D))$ such that $F_0 = s_1$ and locally in z_i , $i = 1, \dots, r$, the F_t , for small $t \neq 0$, can be transformed into members of a fixed \mathcal{S}_i -equisingular respectively \mathcal{S}_i -equianalytical family.

Step 2.1: ‘‘Ansatz’’ and first reduction for a local investigation.

Let us make the following ‘‘Ansatz’’:

$$b_1 = 1, \quad b_2 = \dots = b_e = 0,$$

$$a_{k,l}^i = t^{m_i-k-l} \cdot \tilde{a}_{k,l}^i, \quad \text{for } i = 1, \dots, r, \quad 0 \leq k+l \leq m_i - 1.$$

This gives rise to a family

$$F_{(t, \underline{\tilde{a}})} = s_1 + \sum_{i=1}^r \sum_{k+l=0}^{m_i-1} t^{m_i-k-l} \tilde{a}_{k,l}^i s_{k,l}^i \in H^0(\mathcal{O}_\Sigma(D))$$

with $t \in \mathbb{C}$ and $\underline{\tilde{a}} = (\tilde{a}^1, \dots, \tilde{a}^r)$ where $\tilde{a}^i = (\tilde{a}_{k,l}^i \mid 0 \leq k+l \leq m_i - 1) \in \mathbb{C}^{N_i}$ with $N_i = \binom{m_i+1}{2}$. In particular, $F_{(0, \underline{\tilde{a}})} = s_1$.

Fixing $i \in \{1, \dots, r\}$, in local coordinates at z_i the family looks like

$$F_{(t, \underline{\tilde{a}})}^i := (F_{(t, \underline{\tilde{a}})})_{z_i} = \sum_{k+l=0}^{m_i-1} t^{m_i-k-l} \tilde{a}_{k,l}^i x_i^k y_i^l + \sum_{|\alpha| \geq m_i} \varphi_\alpha^i(t, \underline{\tilde{a}}) x_i^{\alpha_1} y_i^{\alpha_2},$$

with

$$\varphi_\alpha^i(t, \underline{\tilde{a}}) = B_\alpha^i + \sum_{j=1}^r \sum_{k+l=0}^{m_j-1} t^{m_j-k-l} \tilde{a}_{k,l}^j A_{\alpha, k, l}^{j, i},$$

and thus

$$\varphi_\alpha^i(0, \underline{\tilde{a}}) = B_\alpha^i = a_{\alpha_1, \alpha_2}^{i, \text{fix}}, \quad \text{for } |\alpha| = m_i.$$

For $t \neq 0$ the transformation $\psi_t^i : (x_i, y_i) \mapsto (tx_i, ty_i)$ is indeed a coordinate transformation, and thus $F_{(t, \underline{a})}^i$ is contact equivalent to

$$\tilde{F}_{(t, \underline{a})}^i := t^{-m_i} \cdot F_{(t, \underline{a})}^i(tx_i, ty_i) = \sum_{k+l=0}^{m_i-1} \tilde{a}_{k,l}^i x_i^k y_i^l + \sum_{|\alpha| \geq m_i} t^{|\alpha|-m_i} \varphi_\alpha^i(t, \underline{a}) x_i^{\alpha_1} y_i^{\alpha_2}.$$

Note that for this new family in $\mathbb{C}\{x_i, y_i\}$ we have

$$\tilde{F}_{(0, \underline{a}^{\text{fix}})}^i = \sum_{k+l=0}^{m_i} a_{k,l}^{i, \text{fix}} x_i^k y_i^l = g_i,$$

and hence it gives rise to a deformation of $(g_i^{-1}(0), 0)$.

Step 2.2: Reduction to the investigation of the equisingular respectively equianalytical strata of certain families of polynomials.

It is basically our aim to verify the \underline{a} as convergent power series in t such that the corresponding family is equisingular respectively equianalytical. However, since the $\tilde{F}_{(t, \underline{a})}^i$ are power series in x_i and y_i , we cannot right away apply the T-smoothness property of g_i , but we rather have to reduce to polynomials. For this let e_i be the determinacy bound⁵ of S_i and define

$$\hat{F}_{(t, \underline{a})}^i := \sum_{k+l=0}^{m_i-1} \tilde{a}_{k,l}^i x_i^k y_i^l + \sum_{|\alpha|=m_i}^{e_i} t^{|\alpha|-m_i} \varphi_\alpha^i(t, \underline{a}) x_i^{\alpha_1} y_i^{\alpha_2} \equiv \tilde{F}_{(t, \underline{a})}^i \pmod{(x_i, y_i)^{e_i+1}}.$$

Thus $\hat{F}_{(t, \underline{a})}^i$ is a family in $\mathbb{C}[x_i, y_i]_{\leq e_i}$, and still

$$\hat{F}_{(0, \underline{a}^{\text{fix}})}^i = \tilde{F}_{(0, \underline{a}^{\text{fix}})}^i = g_i.$$

We claim that it suffices to find $\underline{a}(t) \in \mathbb{C}\{t\}$ with $\underline{a}(0) = (a_{k,l}^{i, \text{fix}} \mid i = 1, \dots, r, 0 \leq k+l \leq m_i-1)$, such that the families $\hat{F}_t^i := \hat{F}_{(t, \underline{a}(t))}^i$, $t \in (\mathbb{C}, 0)$, are in the equisingular respectively equianalytical strata $\mathbb{C}[x_i, y_i]_{\leq e_i}^*$, for $i = 1, \dots, r$.

Since then we have, for small⁶ $t \neq 0$,

$$g_i = \hat{F}_0^i \sim_c \hat{F}_t^i \sim \tilde{F}_{(t, \underline{a}(t))}^i \sim_c F_{(t, \underline{a}(t))}^i = (F_{(t, \underline{a}(t))})_{z_i(t)},$$

by the e_i -determinacy and since ψ_t^i is a coordinate change for $t \neq 0$, which proves condition (2). Note that the singular points z_i will move with t .

Step 2.3: Find $\underline{a}(t) \in \mathbb{C}\{t\}^n$ with $\underline{a}(0) = (a_{k,l}^{i, \text{fix}}, i = 1, \dots, r, 0 \leq k+l \leq m_i-1)$, $n = \sum_{i=1}^r \binom{m_i+1}{2}$, such that the families $\hat{F}_t^i = \hat{F}_{(t, \underline{a}(t))}^i$, $t \in (\mathbb{C}, 0)$, are in the equisingular respectively equianalytical strata $\mathbb{C}[x_i, y_i]_{\leq e_i}^*$, for $i = 1, \dots, r$.

⁵A power series $f \in \mathcal{O}_2 = \mathbb{C}\{x, y\}$ (respectively the singularity $(V(f), 0)$ defined by f) is said to be *finitely determined* with respect to some equivalence relation \sim if there exists some positive integer e such that $f \sim g$ whenever f and g have the same e -jet. If f is finitely determined, the smallest possible e is called the *determinacy bound*. Isolated singularities are finitely determined with respect to analytical equivalence and hence as well with respect to topological equivalence. C. f. [DeP00] Theorem 9.1.3 and Footnote 6.

⁶Here $f \sim g$ means $f \sim_t g$ respectively $f \sim_c g$ in the sense of Definition I.2.1. Note that if f and g are contact equivalent, then there exists even an analytic coordinate change Φ , that is, $f \sim_c g$ implies $f \sim_t g$.

In the sequel we adopt the notation of Definition 2.1 adding indices i in the obvious way.

Since $\mathbb{C}[x_i, y_i]_{\leq m_i}$ is \mathbb{T} -smooth at g_i , for $i = 1, \dots, r$, there exist $\Lambda_i \subseteq \{(k, l) \mid 0 \leq k + l \leq m_i\}$ and power series $\phi_{k,l}^i \in \mathbb{C}\{\tilde{\underline{a}}_{(1)}^i, \tilde{\underline{a}}_{(2)}^i\}$, for $(k, l) \in \Lambda_i$, such that the equisingular respectively equianalytical stratum $\mathbb{C}[x_i, y_i]_{\leq e_i}^*$ is given by the $\tau_i^* = \tau^*(\mathcal{S}_i) = \#\Lambda_i$ equations

$$\tilde{a}_{k,l}^i = \phi_{k,l}^i(\tilde{\underline{a}}_{(1)}^i, \tilde{\underline{a}}_{(2)}^i), \quad \text{for } (k, l) \in \Lambda_i.$$

Setting $\Lambda = \bigcup_{j=1}^r \{j\} \times \Lambda_j$ we use the notation $\tilde{\underline{a}}_{(0)} = (\tilde{\underline{a}}_{(0)}^1, \dots, \tilde{\underline{a}}_{(0)}^r) = (\tilde{a}_{k,l}^i \mid (i, k, l) \in \Lambda)$ and, similarly $\tilde{\underline{a}}_{(1)}$, $\tilde{\underline{a}}_{(2)}$, $\underline{a}_{(1)}^{i, \text{fix}}$, $\underline{a}_{(0)}^{i, \text{fix}}$, and $\underline{a}_{(1)}^{i, \text{fix}}$. Moreover, setting $\tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)}) = (t^{|\alpha| - m_i} \varphi_\alpha^i(t, \tilde{\underline{a}}_{(0)}, \underline{a}_{(1)}^{i, \text{fix}}) \mid m_i \leq |\alpha| \leq e_i)$, we define an analytic map germ

$$\Phi : \left(\mathbb{C} \times \mathbb{C}^{\tau_1^*} \times \dots \times \mathbb{C}^{\tau_r^*}, (0, \underline{a}_{(0)}^{i, \text{fix}}) \right) \rightarrow \left(\mathbb{C}^{\tau_1^*} \times \dots \times \mathbb{C}^{\tau_r^*}, 0 \right)$$

by

$$\Phi_{k,l}^i(t, \tilde{\underline{a}}_{(0)}) = \tilde{a}_{k,l}^i - \phi_{k,l}^i(\underline{a}_{(1)}^{i, \text{fix}}, \tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)})), \quad \text{for } (i, k, l) \in \Lambda,$$

and we consider the system of equations

$$\Phi_{k,l}^i(t, \tilde{\underline{a}}_{(0)}) = 0, \quad \text{for } (i, k, l) \in \Lambda.$$

We show in Step 2.4 that

$$\left(\frac{\partial \Phi_{k,l}^i}{\partial \tilde{a}_{\kappa,\lambda}^j} (0, \underline{a}_{(0)}^{i, \text{fix}}) \right)_{(i,k,l),(j,\kappa,\lambda) \in \Lambda} = \text{id}_{\mathbb{C}^n}.$$

Thus by the Inverse Function Theorem there exist $\tilde{a}_{k,l}^i(t) \in \mathbb{C}\{t\}$ with $\tilde{a}_{k,l}^i(0) = \underline{a}_{k,l}^{i, \text{fix}}$ such that

$$\tilde{a}_{k,l}^i(t) = \phi_{k,l}^i(\underline{a}_{(1)}^{i, \text{fix}}, \tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)}(t))), \quad (i, k, l) \in \Lambda.$$

Now, setting $\tilde{\underline{a}}_{(1)}(t) \equiv \underline{a}_{(1)}^{i, \text{fix}}$, the families $\hat{F}_t^i = \hat{F}_{(t, \tilde{\underline{a}}_{(0)}^i)}^i$ are in the equisingular respectively equianalytical strata $\mathbb{C}[x_i, y_i]_{\leq e_i}^*$, for $i = 1, \dots, r$.

Step 2.4: Show that $\left(\frac{\partial \Phi_{k,l}^i}{\partial \tilde{a}_{\kappa,\lambda}^j} (0, \underline{a}_{(0)}^{i, \text{fix}}) \right)_{(i,k,l),(j,\kappa,\lambda) \in \Lambda} = \text{id}_n$.

For this it suffices to show that

$$\left. \frac{\partial \phi_{k,l}^i(\underline{a}_{(1)}^{i, \text{fix}}, \tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)}))}{\partial \tilde{a}_{\kappa,\lambda}^j} \right|_{t=0, \tilde{\underline{a}}_{(0)} = \underline{a}_{(0)}^{i, \text{fix}}} = 0,$$

for any $(i, k, l), (j, \kappa, \lambda) \in \Lambda$, which is fulfilled since the map

$$\theta : (t, \tilde{\underline{a}}_{(0)}) \mapsto \left(\underline{a}_{(1)}^{i, \text{fix}}, \tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)}) \right)$$

satisfies

$$\left. \frac{\partial \theta}{\partial \tilde{a}_{\kappa,\lambda}^j} \right|_{t=0, \tilde{\underline{a}}_{(0)} = \underline{a}_{(0)}^{i, \text{fix}}} = \left(\underline{0}, \left. \frac{\partial \tilde{\varphi}^i(t, \tilde{\underline{a}}_{(0)})}{\partial \tilde{a}_{\kappa,\lambda}^j} \right|_{t=0, \tilde{\underline{a}}_{(0)} = \underline{a}_{(0)}^{i, \text{fix}}} \right) = 0.$$

For the latter notice that

$$\frac{\partial \tilde{\varphi}^i(t, \tilde{\mathbf{a}}_{(0)})}{\partial \tilde{\mathbf{a}}_{\kappa, \lambda}^j} \Big|_{t=0, \tilde{\mathbf{a}}_{(0)} = \mathbf{a}_{(0)}^{\text{fix}}} = \left(t^{|\alpha| - m_i} \cdot \frac{\partial \varphi_{\alpha}^i(t, \tilde{\mathbf{a}}_{(0)}, \mathbf{a}_{(1)}^{\text{fix}})}{\partial \tilde{\mathbf{a}}_{\kappa, \lambda}^j} \Big|_{t=0, \tilde{\mathbf{a}}_{(0)} = \mathbf{a}_{(0)}^{\text{fix}}} \right) \Big|_{m_i \leq |\alpha| \leq e_i}$$

and for $|\alpha| = m_i$

$$\varphi_{\alpha}^i(t, \tilde{\mathbf{a}}) = \mathbf{a}_{\alpha_1, \alpha_2}^{i, \text{fix}} + t \cdot \sum_{j=1}^r \sum_{k+l=0}^{m_j-1} t^{m_j-1-k-l} \tilde{\mathbf{a}}_{k,l}^j \mathbf{A}_{\alpha, k, l}^{j, i}.$$

Step 3: It finally remains to show that F_t , for small $t \neq 0$, has no other singular points than $z_1(t), \dots, z_r(t)$.

$C_t \in |D|_i$ shall denote the curve defined by F_t , $t \in (\mathbb{C}, 0)$, in particular $C_0 = C$.

Since for any $i = 1, \dots, r$ the family F_t , $t \in (\mathbb{C}, 0)$, induces a deformation of the singularity (C_0, z_i) there are, by the Conservation of Milnor Numbers (cf. [DeP00], Chapter 6), (Euclidean) open neighbourhoods $U(z_i) \subset \Sigma$ and $V(0) \subset \mathbb{C}$ such that for any $t \in V(0)$

$$(2.1) \quad \text{Sing}(C_t) \subset \bigcup_{i=1}^r U(z_i), \text{ i. e. singular points of } C_t \text{ come from singular points of } C_0,$$

$$(2.2) \quad \mu(C_0, z_i) = \sum_{z \in \text{Sing}(F_t^i) \cap U(z_i)} \mu(F_t^i, z), \quad i = 1, \dots, r.$$

Let $i \in \{1, \dots, r\}$. For $t \neq 0$ fixed we consider the transformation ψ_t^i defined by the coordinate change ψ_t^i ,

$$\begin{array}{ccc} \mathbb{C}^2 \supset U(z_i) & \longrightarrow & U_t(z_i) \subset \mathbb{C}^2 \\ \psi & & \psi \\ (x_i, y_i) & \mapsto & \left(\frac{1}{t}x_i, \frac{1}{t}y_i\right), \end{array}$$

and the transformed equations

$$\tilde{F}_t^i(x_i, y_i) = t^{-m_i} F_t^i(tx_i, ty_i) = 0.$$

Then obviously,

$$F_t^i(z) = 0 \iff (\tilde{F}_t^i \circ \psi_t^i)(z) = 0,$$

and

$$\nabla \psi_t^i \equiv \frac{1}{t} \text{id}_{\mathbb{C}^2}.$$

Thus we have

$$(m_i - 1)^2 = \mu(C_0, z_i) = \sum_{z \in \text{Sing}(F_t^i) \cap U(z_i)} \mu(F_t^i, z) = \sum_{z \in \text{Sing}(\tilde{F}_t^i) \cap U_t(z_i)} \mu(\tilde{F}_t^i, z).$$

For $t \neq 0$ very small $U_t(z_i)$ becomes very large, so that, by shrinking $V(0)$ we may suppose that for any $0 \neq t \in V(0)$

$$\text{Sing}(g_i) \subset U_t(z_i),$$

and that for any $z \in \text{Sing}(g_i)$ there is an open neighbourhood $U(z) \subset U_t(z_i)$ such that

$$\mu(g_i, z) = \sum_{z' \in \text{Sing}(\tilde{F}_t^i) \cap U(z)} \mu(\tilde{F}_t^i, z').$$

If we now take into account that g_i has precisely one critical point, z_i , on its zero level, and that the critical points on the zero level of \tilde{F}_t^i all contribute to the Milnor number $\mu(g_i, z_i)$, then we get the following sequence of inequalities:⁷

$$\begin{aligned} (m_i - 1)^2 - \mu(\mathcal{S}_i) &= \sum_{z \in \text{Sing}(g_i)} \mu(g_i, z) - \sum_{z \in \text{Sing}(g_i^{-1}(0))} \mu(g_i, z) \\ &\leq \sum_{z \in \text{Sing}(\tilde{F}_t^i) \cap U_t(z_i)} \mu(\tilde{F}_t^i, z) - \sum_{z \in \text{Sing}((\tilde{F}_t^i)^{-1}(0)) \cap U_t(z_i)} \mu(\tilde{F}_t^i, z) \\ &= \sum_{z \in \text{Sing}(F_t^i) \cap U(z_i)} \mu(F_t^i, z) - \sum_{z \in \text{Sing}((F_t^i)^{-1}(0)) \cap U(z_i)} \mu(F_t^i, z) \\ &\leq \mu(C_0, z_i) - \mu(F_t^i, z_i) = (m_i - 1)^2 - \mu(\mathcal{S}_i). \end{aligned}$$

Hence all inequalities must have been equalities, and, in particular,

$$\text{Sing}(C_t) \cap U(z_i) = \text{Sing}((F_t^i)^{-1}(0)) \cap U(z_i) = \{z_i\},$$

which in view of condition (2.1) finishes the proof.

Note that C_t , being a small deformation of the irreducible reduced curve $C = C_0$, will again be irreducible and reduced. \square

Now applying the existence theorem of Section 1 for ordinary fat point schemes we deduce the following corollary, giving explicit numerical criteria for the existence of curves with prescribed topological respectively analytical singularities on an arbitrary smooth projective surface.

2.4 Corollary

Let $L \in \text{Div}(\Sigma)$ be very ample over \mathbb{C} . Suppose that $D \in \text{Div}(\Sigma)$ and $\mathcal{S}_1, \dots, \mathcal{S}_r$ are topological respectively analytical singularity types with $e^*(\mathcal{S}_1) \geq \dots \geq e^*(\mathcal{S}_r)$ satisfying

$$(2.3) \quad (D - L - K_\Sigma)^2 \geq 2 \sum_{i=1}^r (e^*(\mathcal{S}_i) + 1)^2,$$

$$(2.4) \quad (D - L - K_\Sigma) \cdot B > e^*(\mathcal{S}_1)$$

for any irreducible curve B with $B^2 = 0$ and $\dim |B|_a > 0$,

⁷Note, an ordinary plane curve singularity (X, x) of multiplicity m has Milnor number $\mu(X, x) = \dim_{\mathbb{C}} (\mathbb{C}\{x, y\} / (x^{m-1}, y^{m-1})) = (m-1)^2$. And, for an affine plane curve given by an equation g such that the equation has no critical point at infinity we have by Bézout's Theorem:

$$(m-1)^2 = \sum_{p \in \mathbb{A}^2} i \left(\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}; p \right),$$

where by definition $i(\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}; p)$ and $\mu(g, p)$ are defined as the dimension of the same vector spaces.

$$(2.5) \quad D.L - 2g(L) > e^*(\mathcal{S}_1), \text{ and}$$

$$(2.6) \quad D - L - K_\Sigma \text{ is nef,}$$

then there is an irreducible reduced curve C in $|D|_1$ with r singular points of topological respectively analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

Proof: Due to the definition of a good representative there is an open dense subset U_i of m_i -forms for which there exists a good representative of \mathcal{S}_i , where $m_i = e^*(\mathcal{S}_i)$. The above conditions then imply with the aid of Corollary 1.3 the existence of an irreducible curve $C \in |D|_1$ with $\text{Sing}(C) = \{z_1, \dots, z_r\}$ whose m_i -jet in local coordinates at z_i coincides with the m_i -jet of a good representative. Moreover, $h^1(\Sigma, \mathcal{J}_{X(\underline{m}, \underline{z})/\Sigma}(D)) = 0$ and thus Theorem 2.3 applies. \square

In view of the bounds for m_i in Remark 2.2 we get the following corollary.

2.5 Corollary

Let $L \in \text{Div}(\Sigma)$ be very ample over \mathbb{C} and $D \in \text{Div}(\Sigma)$.

- (a) Suppose that $\mathcal{S}_1, \dots, \mathcal{S}_r$ are topological singularity types with $\delta(\mathcal{S}_1) \geq \dots \geq \delta(\mathcal{S}_r)$, among them k nodes and m cusps, satisfying

$$(2.7) \quad (D - L - K_\Sigma)^2 \geq 18k + 32m + \frac{625}{24} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i),$$

$$(2.8) \quad (D - L - K_\Sigma).B > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1$$

for any irreducible curve B with $B^2 = 0$ and $\dim |B|_a > 0$,

$$(2.9) \quad D.L - 2g(L) > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1, \text{ and}$$

$$(2.10) \quad D - L - K_\Sigma \text{ is nef,}$$

then there is an irreducible reduced curve C in $|D|_1$ with r singular points of topological types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

- (b) Suppose that $\mathcal{S}_1, \dots, \mathcal{S}_r$ are analytical singularity types with $\mu(\mathcal{S}_1) \geq \dots \geq \mu(\mathcal{S}_r)$, among them k nodes and m cusps, satisfying

$$(2.11) \quad (D - L - K_\Sigma)^2 \geq 18k + 32m + 18 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i),$$

$$(2.12) \quad (D - L - K_\Sigma).B > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1$$

for any irreducible curve B with $B^2 = 0$ and $\dim |B|_a > 0$,

$$(2.13) \quad D.L - 2g(L) > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1, \text{ and}$$

$$(2.14) \quad D - L - K_\Sigma \text{ is nef,}$$

then there is an irreducible reduced curve C in $|D|_1$ with r singular points of analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

2.6 Remark

Knowing something more about the singularity type one can achieve much better results, applying the corresponding bounds for the $e^s(\mathcal{S}_i)$ respectively $e^a(\mathcal{S}_i)$.

3. Examples

In this section we are going to examine the conditions in the existence results for various types of surfaces. $m_1, \dots, m_r \in \mathbb{N}$ are positive integers.

3.a. The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$

We note that in $\mathbb{P}_{\mathbb{C}}^2$ there are no irreducible curves of self-intersection number zero, and we take $L \in |\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^2}(1)|_l$ to be a generic line.

In the existence theorem Corollary 1.3 for ordinary fat point schemes, we, of course, find that condition (1.9) is obsolete, and so is (1.11), taking into account that (1.10) implies $d > 0$. But then conditions (1.10) becomes also redundant in view of condition (1.8) and the assumption $d \geq 3$.

Thus Corollary 1.3 and Corollary 2.5 reduce to the following versions.

1.3a Corollary

Let H be a line in $\mathbb{P}_{\mathbb{C}}^2$, and $d \geq 3$ an integer satisfying

$$(1.8a) \quad (d+2)^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2,$$

then for $z_1, \dots, z_r \in \Sigma$ in very general position there is an irreducible reduced curve $C \in |dH|_l$ with ordinary singularities of multiplicities m_i at the z_i as only singularities. Moreover, $V_{|dH|}(\underline{m})$ is T -smooth at C .

2.5a Corollary

Let H be a line in $\mathbb{P}_{\mathbb{C}}^2$, $\mathcal{S}_1, \dots, \mathcal{S}_r$ topological respectively analytical singularity types, which are neither nodes nor cusps, and $d \geq 3$ an integer such that

$$(2.7a) \quad (d+2)^2 \geq 18k + 32m + \frac{625}{24} \cdot \sum_{i=1}^r \delta(\mathcal{S}_i)$$

respectively

$$(2.11a) \quad (d+2)^2 \geq 18k + 32m + 18 \cdot \sum_{i=1}^r \mu(\mathcal{S}_i).$$

Then there is an irreducible reduced curve C in $|D|_l$ with r singular points of topological respectively analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$, k nodes and m cusps as its only singularities.

Our results for ordinary multiple points are weaker than those in [GLS98c] (see also [Los98] Proposition 4.11), where the factor 2 is replaced by $\frac{10}{9}$) which use the Vanishing Theorem of Geng Xu (cf. [Xu95] Theorem 3), particularly designed for $\mathbb{P}_{\mathbb{C}}^2$. Similarly, our general conditions are weaker than the conditions which recently have been found by Shustin applying the Castelnuovo function (cf. [Shu01]). – Using $L \in |\mathcal{O}_{\Sigma}(l)|_l$ with $l > 1$ instead of $\mathcal{O}_{\Sigma}(1)$ in Corollary 1.3 does not improve the conditions.

3.b. Geometrically Ruled Surfaces

Throughout this section we use the notation of Section G.a.

With the choice of $L = C_0 + lF$ as indicated in Remark G.4 we have $g(L) = g$, and hence the generic curve in $|L|_l$ is a smooth curve whose genus equals the genus of the base curve.

In order to obtain nice formulae we considered $D = (a - 2)C_0 + (b - 2 + 2g)F$ in the formulation of the vanishing theorem (Corollary II.1.2b). For the existence theorems it turns out that the formulae look best if we work with the divisor $D = (a - 1)C_0 + (b + l + 2g - 2 - e)F$ instead. In the case of Hirzebruch surfaces this is just $D = (a - 1)C_0 + (b - 1)F$.

1.3b Corollary

Given integers $a, b \in \mathbb{Z}$ satisfying

$$(1.8b) \quad a(b - \frac{a}{2}e) \geq \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.9b.i) \quad a > \max\{m_i \mid i = 1, \dots, r\},$$

$$(1.9b.ii) \quad b > \max\{m_i \mid i = 1, \dots, r\}, \quad \text{if } e = 0,$$

$$(1.9b.iii) \quad 2 \cdot (b - \frac{a}{2}e) > \max\{m_i \mid i = 1, \dots, r\}, \quad \text{if } e < 0, \text{ and}$$

$$(1.11b) \quad b \geq ae, \text{ if } e > 0,$$

then for $z_1, \dots, z_r \in \Sigma$ in very general position there is an irreducible reduced curve $C \in |(a - 1)C_0 + (b + l + 2g - 2 - e)F|_a$ with ordinary singularities of multiplicities m_i at the z_i as only singularities. Moreover, $V_{|C|}(\underline{m})$ is T -smooth at C .

Proof: Note that by (1.8b) and (1.9b.i) $b > \frac{a}{2}e \geq ae$, if $e \leq 0$, and thus the inequality

$$b \geq ae, \tag{3.1}$$

is fulfilled no matter what e is.

Noting that $D - L - K_\Sigma \sim_a aC_0 + bF$, it is in view of Lemma G.2 clear, that the conditions (1.8) and (1.9) take the form (1.8b) respectively (1.9b). It, therefore, remains to show that (1.10) is obsolete, and that (1.11) takes the form (1.11b), which in particular means that it is obsolete in the case $\Sigma \cong C \times \mathbb{P}_C^1$.

Step 1: (1.10) is obsolete.

If $\Sigma \not\cong \mathbb{P}_C^1 \times \mathbb{P}_C^1$, then $l \geq 2$. Since, moreover, $g(L) = g$ and $D.L = a(l - e) + b + 2g - 2$, condition (1.9b.i) and (3.1) imply (1.10), i. e. for all i, j

$$D.L - 2g(L) = a(l - e) + b - 2 \geq \begin{cases} a + b - 2 > m_i, & \text{if } \Sigma \cong \mathbb{P}_C^1 \times \mathbb{P}_C^1, \\ 2a + (b - ae) - 2 > m_i, & \text{else.} \end{cases}$$

Step 2: (1.11) takes the form (1.11b).

If $e \leq 0$, then by [Har77] V.2.20 and V.2.21 we find in view of (1.9b.i)-(1.9b.iii) that $D - L - K_\Sigma$ is even ample, while, if $e > 0$, the result follows from Lemma G.1. \square

2.5b Corollary

Given integers a and b .

(a) Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ topological singularity types with $\delta(\mathcal{S}_1) \geq \dots \geq \delta(\mathcal{S}_r)$, among them k nodes and m cusps, such that

$$(2.7b) \quad a(b - \frac{a}{2}e) \geq 9k + 16m + \frac{625}{48} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i),$$

$$(2.8b.i) \quad a > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1$$

$$(2.8b.ii) \quad b > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1 \text{ if } e = 0,$$

$$(2.8b.iii) \quad 2(b - \frac{a}{2}e) > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1 \text{ if } e < 0, \text{ and}$$

$$(2.10b) \quad b \geq ae, \text{ if } e > 0.$$

Then there is an irreducible reduced curve C in $|(a-1)C_0 + (b+l+2g-2-e)F|_a$ with r singular points of topological types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

(b) Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ analytical singularity types with $\mu(\mathcal{S}_1) \geq \dots \geq \mu(\mathcal{S}_r)$, among them k nodes and m cusps, such that

$$(2.11b) \quad a(b - \frac{a}{2}e) \geq 9k + 16m + 9 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i),$$

$$(2.12b.i) \quad a > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1$$

$$(2.12b.ii) \quad b > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1 \text{ if } e = 0,$$

$$(2.12b.iii) \quad 2(b - \frac{a}{2}e) > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1 \text{ if } e < 0, \text{ and}$$

$$(2.14b) \quad b \geq ae, \text{ if } e > 0.$$

Then there is an irreducible reduced curve C in $|(a-1)C_0 + (b+l+2g-2-e)F|_a$ with r singular points of analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

3.c. Products of Curves

As we have seen in Proposition G.12, for a generic choice of smooth projective curves of genera $g_1 \geq 1$ and $g_2 \geq 1$ respectively the surface $\Sigma = C_1 \times C_2$ has Picard number two. Furthermore, according to Remark G.11 the only irreducible curves $B \subset \Sigma$ with selfintersection $B^2 = 0$ are the fibres C_1 and C_2 , and for any irreducible curve $B \sim_a aC_1 + bC_2$ the coefficients a and b must be non-negative.

Then choosing for the existence theorem Corollary 1.3 $l \geq 3$ minimal such that $L = lC_1 + lC_2$ is very ample, we claim that the conditions (1.10) and (1.11) become obsolete and the corollary takes the following form.

1.3c Corollary

Let $\Sigma = C_1 \times C_2$ for two smooth projective curves C_1 and C_2 of genera $g_1, g_2 \geq 1$ such that the Picard number is two, and let $l > 0$ be minimal such that $lC_1 + lC_2$ is very ample. Suppose that $a, b \in \mathbb{Z}$ are two integers satisfying

$$(1.8c) \quad (a - l - 2g_2 + 2) \cdot (b - l - 2g_1 + 2) \geq \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

$$(1.9c) \quad (a - l - 2g_2 + 2), (b - l - 2g_1 + 2) > \max\{m_i \mid i = 1, \dots, r\}.$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position there is an irreducible reduced curve $C \in |aC_1 + bC_2|_a$ with ordinary singularities of multiplicities m_i at the z_i as only singularities. Moreover, $V_{|C|}(\underline{m})$ is T -smooth at C .

Proof: (1.11) becomes redundant in view of (1.9c) and since an irreducible curve $B \sim_a a'C_1 + b'C_2$ has non-negative coefficients a' and b' .

It remains to show that $D.L - 2g(L) > m_i$ for all i, j . However, by the adjunction formula $g(L) = 1 + \frac{1}{2}(L^2 + L.K_\Sigma) = 1 + l \cdot (l + g_1 + g_2 - 2)$, and by (1.9c) $D.L - 2g(L) > l \cdot ((a - l - 2g_2 + 2) + (b - l - 2g_1 + 2)) > 6m_i > m_i$. Thus the claim is proved. \square

From these considerations we at once deduce the conditions for the existence of an irreducible curve in $|D|_l$, $D \sim_a aC_1 + bC_2$, with prescribed singularities of arbitrary type, i. e. the conditions in Corollary 2.5.

2.5c Corollary

Let $\Sigma = C_1 \times C_2$ for two smooth projective curves C_1 and C_2 of genera $g_1, g_2 \geq 1$ such that the Picard number is two, and let $l > 0$ be minimal such that $lC_1 + lC_2$ is very ample.

- (a) Suppose that $a, b \in \mathbb{Z}$ are two integers and $\mathcal{S}_1, \dots, \mathcal{S}_r$ topological singularity types with $\delta(\mathcal{S}_1) \geq \dots \geq \delta(\mathcal{S}_r)$, among them k nodes and m cusps, satisfying

$$(2.7c) \quad (a - l - 2g_2 + 2) \cdot (b - l - 2g_1 + 2) \geq 9k + 16m + \frac{625}{48} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i),$$

$$(2.8c) \quad (a - l - 2g_2 + 2), (b - l - 2g_1 + 2) > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1.$$

Then there is an irreducible reduced curve C in $|aC_1 + bC_2|_a$ with r singular points of topological types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

- (b) Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ analytical singularity types with $\mu(\mathcal{S}_1) \geq \dots \geq \mu(\mathcal{S}_r)$, among them k nodes and m cusps, such that

$$(2.11c) \quad (a - l - 2g_2 + 2) \cdot (b - l - 2g_1 + 2) \geq 9k + 16m + 9 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i),$$

$$(2.12c) \quad (a - l - 2g_2 + 2), (b - l - 2g_1 + 2) > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1.$$

Then there is an irreducible reduced curve C in $|aC_1 + bC_2|_a$ with r singular points of analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

3.d. Products of Elliptic Curves

That C_1 and C_2 be “generic” in the above sense means elliptic curves for just that they are non-isogenous. Working with the very ample divisor class $L = 3C_1 + 3C_2$ the theorems in Section 3.c look as follows.

1.3d Corollary

Let $\Sigma = C_1 \times C_2$ for two non-isogenous elliptic curves C_1 and C_2 . Suppose that $a, b \in \mathbb{Z}$ are two integers satisfying

$$(1.8d) \quad (a-3) \cdot (b-3) \geq \sum_{i=1}^r (m_i + 1)^2, \text{ and}$$

$$(1.9d) \quad (a-3), (b-3) > \max\{m_i \mid i = 1, \dots, r\}.$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position there is an irreducible reduced curve $C \in |aC_1 + bC_2|_a$ with ordinary singularities of multiplicities m_i at the z_i as only singularities. Moreover, $V_{|C|}(\underline{m})$ is T -smooth at C .

2.5d Corollary

Let $\Sigma = C_1 \times C_2$ for two non-isogenous elliptic curves C_1 .

- (a) Suppose that $a, b \in \mathbb{Z}$ are two integers and $\mathcal{S}_1, \dots, \mathcal{S}_r$ topological singularity types with $\delta(\mathcal{S}_1) \geq \dots \geq \delta(\mathcal{S}_r)$, among them k nodes and m cusps, satisfying

$$(2.7d) \quad (a-3) \cdot (b-3) \geq 9k + 16m + \frac{625}{48} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i) \text{ and}$$

$$(2.8d) \quad a-3, b-3 > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1.$$

Then there is an irreducible reduced curve C in $|aC_1 + bC_2|_a$ with r singular points of topological types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

- (b) Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ analytical singularity types with $\mu(\mathcal{S}_1) \geq \dots \geq \mu(\mathcal{S}_r)$, among them k nodes and m cusps, such that

$$(2.11d) \quad (a-3) \cdot (b-3) \geq 9k + 16m + 9 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i) \text{ and}$$

$$(2.12d) \quad a-3, b-3 > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1.$$

Then there is an irreducible reduced curve C in $|aC_1 + bC_2|_a$ with r singular points of analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

3.e. Surfaces in $\mathbb{P}_\mathbb{C}^3$

Since we consider the case of rational surfaces separately the following considerations thus give a full answer for the “general case” of a surface in $\mathbb{P}_\mathbb{C}^3$.

1.3e Corollary

Let Σ be a smooth projective surface of degree n in $\mathbb{P}_\mathbb{C}^3$ whose Picard group is generated by a hyperplane section H , and let $d \geq 3$ be an integer such that

$$(1.8e) \quad n \cdot (d-n+3)^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2.$$

Then for $z_1, \dots, z_r \in \Sigma$ in very general position there is an irreducible reduced curve $C \in |dH|_a$ with ordinary singularities of multiplicities m_i at the z_i as only singularities. Moreover, $V_{|dH|}(\underline{m})$ is T -smooth at C .

Proof: The role of the very ample divisor L is filled by a hyperplane section, and thus $g(L) = 1 + \frac{L^2 + L \cdot K_\Sigma}{2} = \binom{n-1}{2}$. Therefore, (1.8e) obviously implies (1.8), and (1.10) takes the form

$$n \cdot (d - n + 3) > m_i + 2 \text{ for all } i = 1, \dots, r. \quad (3.2)$$

However, from (1.8e) we deduce for any $i \in \{1, \dots, r\}$

$$n \cdot (d - n + 3) \geq \sqrt{n} \cdot \sqrt{2} \cdot (m_i + 1) \geq m_i + 2,$$

unless $n = r = m_1 = 1$, in which case we are done by the assumption $d \geq 3$. Thus (1.10) is redundant.

Moreover, there are no curves of self-intersection zero on Σ , and it thus remains to verify (1.11), which in this situation takes the form

$$d \geq n - 3,$$

and follows at once from (3.2). \square

2.5e Corollary

Let Σ be a smooth projective surface of degree n in $\mathbb{P}_\mathbb{C}^3$ whose Picard group is generated by a hyperplane section H .

- (a) Let d be an integer and $\mathcal{S}_1, \dots, \mathcal{S}_r$ topological singularity types with $\delta(\mathcal{S}_1) \geq \dots \geq \delta(\mathcal{S}_r)$, among them k nodes and m cusps, satisfying

$$(2.7e) \quad n(d - n + 3)^2 \geq 18k + 32m + \frac{625}{24} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i).$$

Then there is an irreducible reduced curve C in $|dH|_a$ with r singular points of topological types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

- (b) Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ analytical singularity types with $\mu(\mathcal{S}_1) \geq \dots \geq \mu(\mathcal{S}_r)$, among them k nodes and m cusps, such that

$$(2.11e) \quad n \cdot (d - n + 3)^2 \geq 18k + 32m + 18 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i).$$

Then there is an irreducible reduced curve C in $|dH|_a$ with r singular points of analytical types $\mathcal{S}_1, \dots, \mathcal{S}_r$ as its only singularities.

3.f. K3-Surfaces

3.f.i. *Generic K3-Surfaces.* A generic K3-surface does not possess an elliptic fibration, and hence it does not possess any irreducible curve of self-intersection zero. (cf. [FrM94] I.1.3.7)

Therefore, the conditions in Corollary 1.3 reduce to

$$(1.8f.i) \quad (D - L)^2 \geq 2 \cdot \sum_{i=1}^r (m_i + 1)^2,$$

$$(1.10f.i) \quad D.L - 2g(L) > m_i \text{ for all } i, j, \text{ and}$$

$$(1.11f.i) \quad D - L \text{ nef,}$$

and, analogously, the conditions in Corollary 2.5 in the topological case reduce to (2.9) and

$$(2.7f.i) \quad (D - L)^2 \geq 18k + 32m + \frac{625}{24} \cdot \sum_{i=1}^{r-k-m} \delta(\mathcal{S}_i), \text{ and}$$

$$(2.10f.i) \quad D - L \text{ nef,}$$

and in the analytical case to (2.13) and

$$(2.11f.i) \quad (D - L)^2 \geq 18k + 32m + 18 \cdot \sum_{i=1}^{r-k-m} \mu(\mathcal{S}_i), \text{ and}$$

$$(2.14f.i) \quad D - L \text{ nef.}$$

3.f.ii. *K3-Surfaces with an Elliptic Structure.* The conditions in Corollary 1.3 then reduce to (1.8f.i), (1.11f.i), (1.10f.i), and

$$(1.9f.ii) \quad (D - L).B > \max\{m_i \mid i = 1, \dots, r\}$$

for any irreducible curve B with $B^2 = 0$.

Similarly, the conditions in Corollary 2.5 reduce to (2.7f.i) / (2.11f.i), (2.10f.i) / (2.14f.i), (2.5), and

$$(2.12f.ii) \quad (D - L).B > \frac{25}{4\sqrt{3}} \cdot \sqrt{\delta(\mathcal{S}_1)} - 1$$

for any irreducible curve B with $B^2 = 0$,

respectively

$$(2.12f.ii) \quad (D - L).B > 3 \cdot \sqrt{\mu(\mathcal{S}_1)} - 1$$

for any irreducible curve B with $B^2 = 0$.

CHAPTER IV

T-Smoothness

The varieties $V_{|D|}(rA_1)$ (respectively the open subvarieties $V_{|D|}^{\text{irr}}(rA_1)$) of reduced (respectively reduced and irreducible) nodal curves in a fixed linear system $|D|_1$ on a smooth projective surface Σ are also called *Severi varieties*. When $\Sigma = \mathbb{P}_c^2$ Severi showed that these varieties are smooth of the expected dimension, whenever they are non-empty – that is, nodes always impose independent conditions. It seems natural to study this question on other surfaces, but it is not surprising that the situation becomes harder.

Tannenbaum showed in [Tan82] that also on K3-surfaces $V_{|D|}(rA_1)$ is always smooth, that, however, the dimension is larger than the expected one and thus $V_{|D|}(rA_1)$ is not T-smooth in this situation. If we restrict our attention to the subvariety $V_{|D|}^{\text{irr}}(rA_1)$ of *irreducible* curves with k nodes, then we gain T-smoothness again whenever the variety is non-empty. That is, while on a K3-surface the conditions which nodes impose on irreducible curves are always independent, they impose dependent conditions on reducible curves.

On more complicated surfaces the situation becomes even worse. Chiantini and Sernesi study in [ChS97] Severi varieties on surfaces in \mathbb{P}_c^3 . They show that on a generic quintic Σ in \mathbb{P}_c^3 with hyperplane section H the variety $V_{|dH|}^{\text{irr}}(\frac{5d(d-2)}{4} \cdot A_1)$ has a non-smooth reduced component of the expected dimension, if d is even. They construct their examples by intersecting a general cone over Σ in \mathbb{P}_c^4 with a general complete intersection surface of type $(2, \frac{d}{2})$ in \mathbb{P}_c^4 and projecting the resulting curve to Σ in \mathbb{P}_c^3 . Moreover, Chiantini and Ciliberto give in [ChC99] examples showing that the Severi varieties $V_{|dH|}^{\text{irr}}(rA_1)$ on a surface in \mathbb{P}_c^3 also may have components of dimension larger than the expected one.

Hence, one can only ask for numerical conditions ensuring that $V_{|dH|}^{\text{irr}}(rA_1)$ is T-smooth, and Chiantini and Sernesi answer this question by showing that on a surface of degree $n \geq 5$ the condition

$$r < \frac{d(d - 2n + 8)n}{4} \tag{0.1}$$

implies that $V_{|dH|}^{\text{irr}}(rA_1)$ is T-smooth for $d > 2n - 8$. Note that the above example shows that this bound is even sharp. Actually Chiantini and Sernesi prove a somewhat more general result for surfaces with ample canonical divisor K_Σ and curves which are in $|pK_\Sigma|_1$ for some $p \in \mathbb{Q}$. For their proof they suppose that for some curve $C \in V_{|dH|}^{\text{irr}}(rA_1)$ the cohomology group $H^1(\Sigma, \mathcal{I}_{X^{\text{es}}(C)/\Sigma}(D))$ does not vanish and derive from this the existence of a Bogomolov unstable

rank-two bundle E . This bundle in turn provides them a divisor Δ_0 such that they are able to find a lower and an upper bound for $\Delta_0 \cdot K_\Sigma$ contradicting each other.

This is basically the same approach which we use as well. However, we allow arbitrary singularities rather than only nodes, we weaken the assumption of K_Σ being ample to $K_\Sigma^2 \geq -3$ and we study a wider range of divisors. Yet we still get the same optimal condition (0.1) for the T-smoothness of nodal curves on surfaces in $\mathbb{P}_\mathbb{C}^3$ – see Corollary 2.6. We should like to point out that our general result for the T-smoothness of $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ with the main condition

$$\left(\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1) + \frac{D \cdot K_\Sigma}{2} \right)^2 < \left(r + \frac{K_\Sigma^2}{4} \right)^2 \cdot D^2,$$

does apply to a wider range of surfaces than most of the previously known results. E. g. we deduce in Section 2.d that on a product $\Sigma = C_1 \times C_2$ of elliptic curves the Severi variety $V_{|aC_1+bC_2|}^{\text{irr}}(rA_1)$ is T-smooth as soon as

$$2r \leq ab.$$

For the plane case there are of course better results like the condition

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 \leq (d + 3)^2,$$

for T-smoothness (cf. [GLS00] or [Los98] Theorem 5.4), where d is the degree of D . Note that the a series of irreducible plane curves of degree d with r singularities of type A_μ , μ arbitrarily large, satisfying

$$r\mu^2 = \sum_{i=1}^r \tau^*(A_\mu) = 9d^2 + \text{terms of lower order}$$

constructed by Shustin (cf. [Shu97]) shows that asymptotically we cannot expect to do essentially better in general. For a survey on other known results on $\Sigma = \mathbb{P}_\mathbb{C}^2$ we refer to [Los98] and [GLS00].

In [GLS98a] nodal curves on the projective plane blown up in $m \geq 10$ points are considered. This is a case to which our theorem does not apply in general since $K_\Sigma^2 = 9 - m < -3$ for $m \geq 13$. Nodal curves on arbitrary rational surfaces were also studied by Tannenbaum in [Tan80]. There he gives for the T-smoothness a condition which basically coincides with the only previously known general condition for the T-smoothness of $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) - \epsilon^*(\mathcal{S}_i) < -K_\Sigma \cdot D,$$

where $\epsilon^*(\mathcal{S}_i) \geq 1$ is the so-called isomorphism defect of \mathcal{S}_i (cf. [GrK89, GrL96]). In [Fla01] Flamini studies the case of Severi varieties on surfaces

of general type and he receives conditions for the T-smoothness of $V_{|D|}^{\text{irr}}(rA_1)$ of the type

$$r < \frac{D \cdot (D - 4H) + \sqrt{D^2 \cdot (D - 4H)^2}}{8},$$

where H is some hyperplane section (cf. [Fla01] Theorem 5.1 and 5.3). For a generalisation of the examples of Chiantini and Sernesi of families of nodal curves which are not T-smooth to general non-degenerate complete intersection surfaces with ample canonical bundle we refer to [FIM01]. These show that the results of [Fla01] are sharp, see [Fla01] Remark 5.6.

The theorem which we here present in the following section and its proof are a slight modification of Theorem 1 in [GLS97]. We sacrifice the sophisticated examination of the zero-dimensional schemes involved in the proof for the sake of simplified conditions. A somewhat stronger version may be found in the book [GLS02].

1. T-Smoothness

1.1 Theorem

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types with $\tau^*(\mathcal{S}_1) \geq \dots \geq \tau^*(\mathcal{S}_r)$ and let $D \in \text{Div}(\Sigma)$ be a divisor such that¹

$$(1.1) \quad D + K_\Sigma \text{ nef,}$$

$$(1.2) \quad D - K_\Sigma \text{ is big and nef,}$$

$$(1.3) \quad D^2 - 2 \cdot D \cdot K_\Sigma \geq \frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4} + 4, \quad \text{if } D \cdot K_\Sigma < -4,$$

$$(1.4) \quad \sum_{i=1}^r \tau^*(\mathcal{S}_i) < \frac{1}{4} \cdot (D - K_\Sigma)^2, \text{ and}$$

$$(1.5) \quad \left(\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) + \frac{D \cdot K_\Sigma}{2} \right)^2 < \left(s + \frac{K_\Sigma^2}{4} \right) \cdot D^2 \quad \text{for all } s = 1, \dots, r.$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T-smooth.

Idea of the Proof: T-smoothness at a point $C \in V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ follows once we know $h^1(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(D)) = 0$. Supposing this is not the case we find a Bogomolov unstable rank-two bundle E which provides us with some “nice” divisor Δ_0 . This means, we get an upper and a lower bound for $D^2 \cdot (D - K_\Sigma - \Delta_0)^2$ which contradict each other due to Condition (1.5).

Proof: Let $C \in V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ be arbitrary. We set $X^*(C) = X^{\text{es}}(C)$ respectively $X^*(C) = X^{\text{ea}}(C)$, so that $\deg(X^*(C)) = \sum_{i=1}^r \tau^*(\mathcal{S}_i)$.

We are going to show that

$$h^1(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(D)) = 0,$$

¹Here $\tau^*(\mathcal{S}_i) = \tau^{\text{es}}(\mathcal{S}_i)$, the codimension of the μ -constant stratum in the base of the semi-universal deformation of \mathcal{S}_i , respectively $\tau^*(\mathcal{S}_i) = \tau(\mathcal{S}_i)$, the Tjurina number of \mathcal{S}_i .

which by Remark I.2.14 implies the T-smoothness of $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ at C .

Suppose the contrary, that is $h^1(\Sigma, \mathcal{J}_{X^*(C)/\Sigma}(D)) > 0$.

Step 1: Find a *non-empty* subscheme $X_0 \subseteq X^*(C)$ and a rank-two vector bundle E satisfying the relations (1.6) and (1.7).

Choose $X_0 \subseteq X^*(C)$ minimal such that still $h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) > 0$. By (1.2) the divisor $D - K_\Sigma$ is big and nef, and thus $h^1(\Sigma, \mathcal{O}_\Sigma(D)) = 0$ by the Kawamata–Viehweg Vanishing Theorem. Hence X_0 cannot be empty.

Due to the Grothendieck-Serre duality (cf. [Har77] III.7.6) we have $0 \neq H^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) \cong \text{Ext}^1(\mathcal{J}_{X_0/\Sigma}(D), \mathcal{O}_\Sigma(K_\Sigma))$, and thus, since $\mathcal{O}_\Sigma(K_\Sigma)$ is locally free, (cf. [Har77] III.6.7)

$$\text{Ext}^1(\mathcal{J}_{X_0/\Sigma}(D - K_\Sigma), \mathcal{O}_\Sigma) \neq 0.$$

That is, there is an extension (cf. [Har77] Ex. III.6.1)

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow E \rightarrow \mathcal{J}_{X_0/\Sigma}(D - K_\Sigma) \rightarrow 0. \quad (1.6)$$

The minimality of X_0 implies that E is locally free (cf. [Laz97] Proposition 3.9). Moreover, we have (cf. [Laz97] Exercise 4.3)

$$c_1(E) = D - K_\Sigma \quad \text{and} \quad c_2(E) = \deg(X_0). \quad (1.7)$$

Step 2: Find a divisor Δ_0 such that

- (a) $(2\Delta_0 - D + K_\Sigma)^2 \geq c_1(E)^2 - 4 \cdot c_2(E) > 0$, and
- (b) $(2\Delta_0 - D + K_\Sigma) \cdot H > 0$ for all $H \in \text{Div}(\Sigma)$ ample.

By (1.4) and (1.7) we have

$$c_1(E)^2 - 4 \cdot c_2(E) = (D - K_\Sigma)^2 - 4 \cdot \deg(X_0) > 0,$$

and thus E is Bogomolov unstable (cf. [Laz97] Theorem 4.2). This, however, implies that there exists a divisor $\Delta_0 \in \text{Div}(\Sigma)$ and a zero-dimensional scheme $Z \subset \Sigma$ such that

$$0 \rightarrow \mathcal{O}_\Sigma(\Delta_0) \rightarrow E \rightarrow \mathcal{J}_{Z/\Sigma}(D - K_\Sigma - \Delta_0) \rightarrow 0 \quad (1.8)$$

is exact (cf. [Laz97] Theorem 4.2) and such that (a) and (b) are fulfilled.

Step 3: Find a curve $\Delta \subset \Sigma$ such that

- (c) $\Delta + \Delta_0 \sim_l D - K_\Sigma$,
- (d) $D \cdot \Delta \geq \deg(X_0) + \#X_0$, and
- (e) $\deg(X_0) \geq \Delta_0 \cdot \Delta$.

Tensoring (1.8) with $\mathcal{O}_\Sigma(-\Delta_0)$ leads to the following exact sequence

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow E(-\Delta_0) \rightarrow \mathcal{J}_{Z/\Sigma}(D - K_\Sigma - 2\Delta_0) \rightarrow 0, \quad (1.9)$$

and we deduce $h^0(\Sigma, E(-\Delta_0)) \neq 0$.

Now tensoring (1.6) with $\mathcal{O}_\Sigma(-\Delta_0)$ leads to

$$0 \rightarrow \mathcal{O}_\Sigma(-\Delta_0) \rightarrow E(-\Delta_0) \rightarrow \mathcal{J}_{X_0/\Sigma}(D - K_\Sigma - \Delta_0) \rightarrow 0. \quad (1.10)$$

Let H be some ample divisor. By (b) and since $D - K_\Sigma$ is nef by (1.2):

$$-\Delta_0.H < -\frac{1}{2} \cdot (D - K_\Sigma).H \leq 0. \quad (1.11)$$

Hence $-\Delta_0$ cannot be effective, that is $H^0(\Sigma, -\Delta_0) = 0$. But the long exact cohomology sequence of (1.10) then implies

$$0 \neq H^0(\Sigma, E(-\Delta_0)) \hookrightarrow H^0(\Sigma, \mathcal{J}_{X_0/\Sigma}(D - K_\Sigma - \Delta_0)).$$

In particular we may choose a curve

$$\Delta \in |\mathcal{J}_{X_0/\Sigma}(D - K_\Sigma - \Delta_0)|_1.$$

Thus (c) is obviously fulfilled and it remains to show (d) and (e).

We note that $C \in |D|_1$ is irreducible and that Δ cannot contain C as an irreducible component: otherwise applying (b) with some ample divisor H we would get the following contradiction, since $D + K_\Sigma$ is nef by (1.1),

$$0 \leq (\Delta - C).H < -\frac{1}{2} \cdot (D + K_\Sigma).H \leq 0.$$

Since $X_0 \subseteq X^*(C)$, Lemma 1.3 applies² to the local ideals of X_0 , that is for the points $z \in \text{supp}(X_0)$ we have $i(C, \Delta; z) \geq \deg(X_0, z) + 1$. Thus, since $X_0 \subset C \cap \Delta$ the Theorem of Bézout implies

$$D.\Delta = C.\Delta = \sum_{z \in C \cap \Delta} i(C, \Delta; z) \geq \sum_{z \in \text{supp}(X_0)} (\deg(X_0, z) + 1) = \deg(X_0) + \#X_0.$$

Finally, by (a), (c) and (1.7) we get

$$(\Delta_0 - \Delta)^2 \geq c_1(E)^2 - 4 \cdot c_2(E) = (\Delta_0 + \Delta)^2 - 4 \cdot \deg(X_0),$$

and thus $\deg(X_0) \geq \Delta_0.\Delta$.

Step 4: Find a lower and an upper bound for $D^2 \cdot (\Delta_0 - \Delta)^2$, contradicting each other.

From (a) we derive

$$(\Delta_0 - \Delta)^2 \geq (D - K_\Sigma)^2 - 4 \deg(X_0). \quad (1.12)$$

²We note that $X^{\text{es}}(C) \subseteq X^{\text{ea}}(C)$, i. e. if $X_0 \subseteq X^{\text{es}}(C)$ then $X_0 \subseteq X^{\text{ea}}(C)$ as well.

(1.1) and (1.2) imply that D is nef, and since the strict inequality for ample divisors in (b) comes down to “ \geq ” for nef divisors, we get in view of (c)

$$\begin{aligned} 0 \leq D \cdot (\Delta_0 - \Delta) &= D \cdot (\Delta_0 + \Delta) - 2 \cdot D \cdot \Delta \\ &\leq D \cdot (\Delta_0 + \Delta) - 2 \deg(X_0) - 2 \cdot \#X_0 \\ &= D \cdot (D - K_\Sigma) - 2 \deg(X_0) - 2 \cdot \#X_0. \end{aligned} \quad (1.13)$$

But then applying the Hodge Index Theorem E.4 to the divisors D and $\Delta_0 - \Delta$, taking (a) into account, we get

$$D^2 \cdot (\Delta_0 - \Delta)^2 \leq (D \cdot (\Delta_0 - \Delta))^2 \leq (D \cdot (D - K_\Sigma) - 2 \deg(X_0) - 2 \cdot \#X_0)^2. \quad (1.14)$$

From (1.12) and (1.14) we deduce that

$$(D \cdot (D - K_\Sigma) - 2 \deg(X_0) - 2 \cdot \#X_0)^2 \geq D^2 \cdot ((D - K_\Sigma)^2 - 4 \deg(X_0)),$$

and hence

$$(2 \deg(X_0) + 2 \cdot \#X_0 + D \cdot K_\Sigma)^2 \geq (4 \cdot \#X_0 + K_\Sigma^2) \cdot D^2 \geq 0. \quad (1.15)$$

Suppose first that $-D \cdot K_\Sigma > 2 \deg(X_0) + 2 \cdot \#X_0 \geq 4$, then by (1.5) and (1.3)

$$(2 \deg(X_0) + 2 \cdot \#X_0 + D \cdot K_\Sigma)^2 < (4 + D \cdot K_\Sigma)^2 \leq (4 + K_\Sigma^2) \cdot D^2 \leq (4 \cdot \#X_0 + K_\Sigma^2) \cdot D^2 \quad (1.16)$$

in contradiction to (1.15).

Thus we must have $2 \deg(X_0) + 2 \cdot \#X_0 + D \cdot K_\Sigma \geq 0$, and since

$$\deg(X_0) \leq \sum_{z \in \text{supp}(X_0)} \tau^*(C, z) \leq \sum_{i=1}^{\#X_0} \tau^*(\mathcal{S}_i)$$

we find once more a contradiction to (1.15) with the aid of (1.5):

$$(2 \deg(X_0) + 2 \cdot \#X_0 + D \cdot K_\Sigma)^2 \leq \left(2 \sum_{i=1}^{\#X_0} (\tau^*(\mathcal{S}_i) + 1) + D \cdot K_\Sigma \right)^2 < (4 \cdot \#X_0 + K_\Sigma^2) \cdot D^2.$$

□

In the following remark we replace (1.5) by a number of other conditions, which partially may even be better in certain situations.

1.2 Remark

- (a) For a surface with $K_\Sigma^2 \leq -3$ condition (1.5) will never be satisfied.
- (b) If K_Σ is nef, then (1.1) and (1.3) are superfluous.
- (c) If $D \cdot K_\Sigma \leq 0$, we can replace (1.5) by

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < D^2 - 2 \cdot D \cdot K_\Sigma - \frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4}. \quad (1.17)$$

Moreover, (1.3) is always fulfilled.

(d) If $D \cdot K_\Sigma \geq 0$, we may replace (1.5) in Theorem 1.1 by

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < D^2 - D \cdot K_\Sigma \cdot (\tau^*(\mathcal{S}_1) + 1) - \frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4}. \quad (1.18)$$

(e) For divisors D of the form $D \sim_n pK_\Sigma$ for some $0 \neq p \in \mathbb{Q}$ the inequalities (1.17) and (1.18) take the simpler forms

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < D^2 - 2 \cdot D \cdot K_\Sigma \quad (1.19)$$

respectively

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < D^2 - D \cdot K_\Sigma \cdot (\tau^*(\mathcal{S}_1) + 1). \quad (1.20)$$

Moreover, (1.3) is always fulfilled.

(f) If D is of the form $D \sim_n pK_\Sigma$ for some $0 \neq p \in \mathbb{Q}$, Condition (1.5) can only be fulfilled if $\tau^*(\mathcal{S}_i) + 1 < \frac{D^2}{D \cdot K_\Sigma}$ for all $i = 1, \dots, r$. Assuming this, the inequality (1.5) may be replaced by

$$\sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)} < D^2. \quad (1.21)$$

(g) We could replace (1.5) by

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) < \sqrt{1 + \frac{K_\Sigma^2}{4}} \cdot \sqrt{D^2} - \left| \frac{K_\Sigma \cdot D}{2} + 1 \right|. \quad (1.22)$$

(h) If $-K_\Sigma$ is ample and $D \sim_n pK_\Sigma$ for some $p \in \mathbb{Q}$ with $p < 1 - \sqrt{1 + \frac{4}{K_\Sigma^2}} < 0$, we could replace (1.5) in Theorem 1.1 by

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) < \sqrt{1 + \frac{K_\Sigma^2}{4}} \cdot \sqrt{D^2} - \frac{1}{2} \cdot K_\Sigma \cdot D - 1. \quad (1.23)$$

(i) If $D^2 \geq 8 + K_\Sigma^2 + 4 \cdot \sqrt{K_\Sigma^2 + 4}$, then (1.22) respectively (1.23) replaces both (1.4) and (1.5).

(j) We note that always

$$4 \cdot \sum_{i=1}^r \tau^*(\mathcal{S}_i) \leq \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2. \quad (1.24)$$

Proof: (a) & (b) These are obvious.

(c) We note that

$$\left(\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) \right)^2 \leq s \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)^2.$$

Thus the inequalities

$$\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)^2 < D^2 - \frac{D \cdot K_\Sigma}{s} \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) - \frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4 \cdot s} \quad (1.25)$$

for $s = 1, \dots, r$ will imply (1.5). However, by (1.1) and (1.2) D is nef and thus $D^2 \geq 0$ and by the Hodge Index Theorem E.4 $\frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4} \geq 0$.

Moreover, $D \cdot K_\Sigma \leq 0$ and we therefore get the desired result noting that 2 is a lower bound for the coefficient $\frac{1}{s} \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)$ of $D \cdot K_\Sigma$.

Since $\tau^*(\mathcal{S}_i) \geq 1$, from (1.17) we have $4 < D^2 - 2 \cdot D \cdot K_\Sigma - \frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4}$, which implies (1.3).

(d) It again suffices to verify (1.25) for $s = 1, \dots, r$. However, since this time D^2 , $\frac{(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2}{4}$ and $D \cdot K_\Sigma$ are all non-negative, (1.25) follows from (1.18), once we note that $\tau^*(\mathcal{S}_1) + 1 \geq \frac{1}{s} \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)$.

(e) If $D \sim_n p \cdot K_\Sigma$, then $(D \cdot K_\Sigma)^2 - D^2 \cdot K_\Sigma^2 = 0$.

(f) Once (1.5) is satisfied, we get from $s = 1$ that $D \cdot K_\Sigma \cdot (\tau^*(\mathcal{S}_1) + 1) < D^2$. Let us therefore suppose that $\tau^*(\mathcal{S}_i) + 1 < \frac{D^2}{D \cdot K_\Sigma}$ for each $i = 1, \dots, r$. We note that then for any $s = 1, \dots, r$

$$\sum_{i=1}^s \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)} \leq \sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)}.$$

The Cauchy-Schwartz Inequality thus shows

$$\begin{aligned} \left(\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) \right)^2 &= \left(\sum_{i=1}^s \frac{\tau^*(\mathcal{S}_i) + 1}{\sqrt{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)}} \cdot \sqrt{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)} \right)^2 \\ &\leq \sum_{i=1}^s \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1)} \cdot \sum_{i=1}^s \left(1 - \frac{D \cdot K_\Sigma}{D^2} \cdot (\tau^*(\mathcal{S}_i) + 1) \right) \\ &< s \cdot D^2 - D \cdot K_\Sigma \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1), \end{aligned}$$

which is just (1.5), since $(D \cdot K_\Sigma)^2 = D^2 \cdot K_\Sigma^2$.

(g) In the proof of Theorem 1.1 we could replace $D \cdot \Delta$ in (1.13) by $\deg(X_0) + 1$ instead of $\deg(X_0) + \#X_0$. Then (1.15) would be replaced by

$$(2 \deg(X_0) + 2 + D \cdot K_\Sigma)^2 \geq (4 + K_\Sigma^2) \cdot D^2 \geq 0, \quad (1.26)$$

and hence

$$\left(2 \deg(X_0) + |2 + D \cdot K_\Sigma| \right)^2 \geq (4 + K_\Sigma^2) \cdot D^2 \geq 0$$

which contradicts (1.22).

(h) We have to derive from (1.26) a contradiction, replacing (1.22) by (1.23). For this we consider the convex function

$$\varphi : \left[1, \deg(X^*(C)) \right] \rightarrow \mathbb{R} : t \mapsto (2t + 2 + D \cdot K_\Sigma)^2.$$

Since φ is convex, it takes its maximum either in $t^* = 1$ or in $t^* = \deg(X^*(C))$.

If $t^* = 1$, then we get from (1.26)

$$(4 + p \cdot K_\Sigma^2)^2 \geq (2 \cdot \deg(X_0) + 2 + D \cdot K_\Sigma)^2 \geq (4 + K_\Sigma^2) \cdot p^2 \cdot K_\Sigma^2,$$

that is $4p^2 K_\Sigma^2 - 8p K_\Sigma^2 - 16 \leq 0$, which contradicts the assumptions on p .

If $t^* = \deg(X^*(C))$, then we get from (1.26)

$$\left(2 \cdot \deg(X^*(C)) + 2 + D \cdot K_\Sigma\right)^2 \geq (4 + K_\Sigma^2) \cdot D^2,$$

and, moreover, in this situation we must have $2 \cdot \deg(X^*(C)) + 2 + D \cdot K_\Sigma \geq 0$. But then we may take square roots on both sides and get

$$2 \cdot \deg(X^*(C)) \geq \sqrt{4 + K_\Sigma^2} \cdot \sqrt{D^2} - D \cdot K_\Sigma - 2,$$

which is in contradiction to (1.22).

(i) By the above condition we know that

$$\left(D^2 - (8 + K_\Sigma^2)\right)^2 \geq 16 \cdot K_\Sigma^2 + 64.$$

This, however, implies

$$\left(1 + \frac{K_\Sigma^2}{4}\right) \cdot D^2 \leq \frac{1}{16} \cdot (D^2 + K_\Sigma^2)^2,$$

and thus finally

$$\sqrt{1 + \frac{K_\Sigma^2}{4}} \cdot \sqrt{D^2} - \left|\frac{K_\Sigma \cdot D}{2} + 1\right| \leq \sqrt{1 + \frac{K_\Sigma^2}{4}} \cdot \sqrt{D^2} - \frac{D \cdot K_\Sigma}{2} - 1 \leq \frac{1}{4} \cdot (D - K_\Sigma)^2,$$

which finishes the proof.

$$(j) \quad 0 \leq \sum_{i=1}^r (\tau^*(\mathcal{S}_i) - 1)^2 = \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 - 4 \cdot \sum_{i=1}^r \tau^*(\mathcal{S}_i).$$

□

Throughout the proof of Theorem 1.1 we used the following lemma.

1.3 Lemma

Let (C, z) be a reduced plane curve singularity given by $f \in \mathcal{O}_{\Sigma, z}$ and let $I \subseteq \mathfrak{m}_{\Sigma, z} \subset \mathcal{O}_{\Sigma, z}$ be an ideal containing the Tjurina ideal $I^{\text{ea}}(C, z)$. Then for any $g \in I$ we have

$$\dim_{\mathbb{C}}(\mathcal{O}_{\Sigma, z}/I) < \dim_{\mathbb{C}}(\mathcal{O}_{\Sigma, z}/(f, g)) = i(f, g; z).$$

Proof: Cf. [Shu97] Lemma 4.1.

□

2. Examples

Throughout this section for a topological respectively analytical singularity type \mathcal{S} we will denote by $\tau^*(\mathcal{S}) = \tau^{\text{es}}(\mathcal{S})$, the codimension of the μ -constant stratum in the base of the semi-universal deformation of \mathcal{S} , respectively $\tau^*(\mathcal{S}) = \tau(\mathcal{S})$, the Tjurina number of \mathcal{S} .

2.a. The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$

In view of Remark 1.2 we get the following version of Theorem 1.1.

1.1a Theorem

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, let $d \geq 3$ and suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < d^2 + 6d. \quad (1.5a)$$

Then either $V_{|\mathrm{dH}|}^{\mathrm{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

Proof: (1.1) and (1.2) are fulfilled since $d \geq 3$, while by Remark 1.2 (e) (1.3) is redundant and (1.5) takes the form (1.5a). It remains to show that (1.5a) implies (1.4), which follows at once from (1.24) in Remark 1.2 (j). \square

2.1 Remark

(a) The sufficient condition

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < (d + 3)^2 \quad (2.1)$$

in [Los98] Corollary 5.5 (see also [GLS00]) is always slightly better than (1.5a).

(b) Going back to the original equations in (1.5) we could replace them by

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < \left(1 + \frac{3\sqrt{4r+9}+9}{2r}\right) \cdot d^2. \quad (2.2)$$

For a fixed r this condition is asymptotically better even than the one in (2.1) – more precisely, $d \geq 3\sqrt{r}$ suffices. However, in order to get rid of (1.4) we then need some assumptions on the singularities – e. g. $r \geq 18$ and $\tau^*(\mathcal{S}_i) \geq 6$ for all i .

(c) Working with condition (1.23) from Remark 1.2 (h) we could replace (1.5a) by

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i)^2 < \frac{10}{r} \cdot d^2, \quad (2.3)$$

if we require $d \geq 8$. This result is better than (2.1) only for $r \leq 9$.

Proof: (a) This is obvious.

(b) Since the derivative

$$\frac{\partial f}{\partial r}(r) = \frac{-3 \cdot (2r + 3\sqrt{4r+9} + 9)}{2 \cdot \sqrt{4r+9} \cdot r^2}$$

of $f : \mathbb{R} \rightarrow \mathbb{R} : r \mapsto \frac{3\sqrt{4r+9}+9}{2r}$ is negative for all $r > 0$, the function is monotonously decreasing, that is

$$\frac{3\sqrt{4r+9}+9}{2r} \leq \frac{3\sqrt{4s+9}+9}{2s}$$

for all $s = 1, \dots, r$, and therefore

$$\begin{aligned} \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)^2 &\leq \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 \\ &< \left(1 + \frac{3\sqrt{4r+9}+9}{2r}\right) \cdot d^2 \leq \left(1 + \frac{3\sqrt{4s+9}+9}{2s}\right) \cdot d^2. \end{aligned}$$

But this implies

$$\begin{aligned} \left(\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)\right)^2 &\leq s \cdot \sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1)^2 \\ &< \left(s + \frac{3}{2} \cdot \sqrt{4s+9} + \frac{9}{2}\right) \cdot d^2 = \left(\sqrt{s + \frac{9}{4}} + \frac{3}{2}\right)^2 \cdot d^2, \end{aligned}$$

and then

$$\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) - \frac{3}{2} \cdot d < \sqrt{s + \frac{9}{4}} \cdot d,$$

which in turn gives (1.5), supposed that the left hand side is non-negative. However, if it is negative, then

$$\left(\sum_{i=1}^s (\tau^*(\mathcal{S}_i) + 1) - \frac{3}{2} \cdot d\right)^2 < \frac{9d^2}{4} < \left(s + \frac{9}{4}\right) \cdot d^2,$$

that is, (1.5) is still fulfilled.

If $d \geq 3\sqrt{r}$, then $\frac{3\sqrt{4r}}{2r} \cdot d^2 \geq 9d \geq 6d + 9$. Thus (2.1) implies (2.2).

Furthermore, if $r \geq 18$, then $1 + \frac{3\sqrt{4r+9}+9}{2r} \leq 2$, and if moreover $\tau^*(\mathcal{S}_i) \geq 6$ for all i , then

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) \leq \frac{1}{8} \cdot \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < \frac{1}{4} \cdot d^2,$$

which implies (1.4).

(c) We note that (2.3) implies

$$\left(\sum_{i=1}^r \tau^*(\mathcal{S}_i)\right)^2 \leq r \cdot \sum_{i=1}^r \tau^*(\mathcal{S}_i)^2 < 10 \cdot d^2 < \left(\frac{\sqrt{13}+3}{2} \cdot d - 1\right)^2,$$

once $d \geq 8$. This however is sufficient for the T-smoothness according to Remark 1.2 (h) and (i).

□

For further results in the plane case see [Wah74a, GrK89, Lue87a, Lue87b, Shu87b, Vas90, Shu91b, Shu94, GrL96, Shu96b, Shu97, GLS98a, Los98, GLS00].

2.b. Geometrically Ruled Surfaces

Throughout this section we use the notation of Section G.a, thus in particular $\pi : \Sigma \rightarrow C$ is a geometrically ruled surface with $g = g(C)$ and $K_\Sigma^2 = 8 - 8g$. Theorem 1.1 therefore only applies for Hirzebruch surfaces and if C is elliptic. Below we state the theorem in its general version for the invariant $e \geq 0$, even though the conditions look far too nasty to be of much use. We afterwards consider the cases separately, so that the conditions take a nicer form, in particular we consider the unique geometrically ruled surface with negative invariant e and $g = 1$.

Throughout the section it is convenient to set $\kappa = 2 + e - 2g$, so that $K_\Sigma \sim -2C_0 - \kappa F$. We note that for $g \leq 1$ and $e \geq 0$ we have $\kappa \geq 0$. Thus $-K_\Sigma$ is nef in this situation and we may replace (1.5) by (1.17) in Remark 1.2.

1.1b Theorem

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with invariant $e \geq 0$ and $g \in \{0, 1\}$, let $D = aC_0 + bF \in \text{Div}(\Sigma)$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$(1.1b) \quad a \geq 2, b \geq e \cdot (a - 2) + \kappa,$$

$$(1.2b) \quad b > e \cdot (a + 2) - \kappa,$$

$$(1.5b) \quad \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < (2 + \kappa) \cdot ab + 4b + (2\kappa - 4e) \cdot a - \left(\frac{\kappa^2}{4} + e\right) \cdot a^2 - b^2.$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

Proof: The conditions (1.1) and (1.2) just come down to (1.1b) and (1.2b), and (1.17) takes the form (1.5b). Moreover, by Remark 1.2 (c) Condition (1.3) is obsolete.

It thus remains to show that (1.5b) implies (1.4), which in this situation looks like

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) < \frac{1}{4} \cdot (2 \cdot (a + 2) \cdot (b + \kappa) - e \cdot (a + 2)^2).$$

In view of (1.24) in Remark 1.2 (j) and by (1.5b) we know already that

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) < \frac{1}{4} \cdot ((2 + \kappa) \cdot ab + 4b + (2\kappa - 4e) \cdot a - \left(\frac{\kappa^2}{4} + e\right) \cdot a^2 - b^2).$$

Since $\kappa - e = 2 - 2g \geq 0$, the claim follows from

$$\begin{aligned} & (2 + \kappa) \cdot ab + 4b + (2\kappa - 4e) \cdot a - \left(\frac{\kappa^2}{4} + e\right) \cdot a^2 - b^2 \\ &= 2 \cdot (a + 2) \cdot (b + \kappa) - e \cdot (a + 2)^2 - \left(\frac{\kappa}{2} \cdot a - b\right)^2 - 4 \cdot (\kappa - e) \\ &\leq 2 \cdot (a + 2) \cdot (b + \kappa) - e \cdot (a + 2)^2. \end{aligned}$$

□

We note that (1.5b) gives an obstruction on a and b , namely

$$(2 + \kappa) \cdot ab + 4b + (2\kappa - 4e) \cdot a \left(\frac{\kappa^2}{4} + e \right) \cdot a^2 + b^2 + 4. \quad (2.4)$$

Let us first consider Hirzebruch surfaces.

2.2 Corollary

Let $\Sigma \cong \mathbb{F}_0 = \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$, let $D \in \text{Div}(\Sigma)$ of type (a, b) with $a, b \geq 2$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 4ab + 4b + 4a - a^2 - b^2. \quad (2.5)$$

Then either $V_{|\mathbb{D}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

We note that the obstruction (2.4) in this case reads

$$4ab + 4b + 4a \geq a^2 + b^2 + 4, \quad (2.6)$$

which could be replaced by

$$a \leq 2b + 3 \quad \text{and} \quad b \leq 2a + 3. \quad (2.7)$$

If we consider the case $a = b$, that is, if D is a rational multiple of the canonical divisor, then (2.5) looks like

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 2a^2 + 8a. \quad (2.8)$$

In this situation we could even use (1.21) in Remark 1.2 instead and replace (2.8) by the better inequality

$$\sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 + \frac{2}{a} \cdot (\tau^*(\mathcal{S}_i) + 1)} < 2a^2. \quad (2.9)$$

Comparing these results with the conditions for irreducibility in Section V.4.b the right hand side there differs by a factor of about $\frac{1}{200}$.

The only Hirzebruch surface, which is not minimal, is \mathbb{F}_1 . Since the obstructions for b there differ from the remaining cases we give the result separately.

2.3 Corollary

Let $\Sigma = \mathbb{F}_1$, let $D = aC_0 + bF \in \text{Div}(\Sigma)$ with $a \geq 2$ and $b \geq a + 1$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 5ab + 4b + 2a - \frac{13}{4} \cdot a^2 - b^2. \quad (2.10)$$

Then either $V_{|\mathbb{D}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

In contrast to the Hirzebruch surfaces with $e \geq 2$, the corollary here applies to divisors, which are multiples of the canonical divisor, that is when $b = \frac{3}{2} \cdot a$. Then, however, we may in view of Remark 1.2 (f) replace (2.10) by the inequality

$$\sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 + \frac{3}{a} \cdot (\tau^*(\mathcal{S}_i) + 1)} < 2a^2. \quad (2.11)$$

For the Hirzebruch surfaces with $e \geq 2$ we have to study the conditions (1.2b) and (2.4) a bit closer. We then see that they may be replaced by

$$b \geq ea + e - 1 \quad (2.12)$$

and

$$\varphi_{a,e}(b) := -\left(b - \frac{e+2}{2} \cdot a\right)^2 + (2ab - ea^2) + 4b + (4 - 2e) \cdot a - 4 > 0. \quad (2.13)$$

For $a \geq 2$ fixed we therefore may consider $\varphi_{a,e} : [ea + e - 1, \infty[\rightarrow \mathbb{R}$ as a concave function, and we find that

$$\varphi_{a,e}(b) > 0 \iff b < \left(2 + \frac{e}{2}\right) \cdot a + \sqrt{3a^2 + 12a} + 2.$$

In particular, (2.12) and (2.13) cannot be satisfied both at the same time unless $e \leq 7$! And if $e \leq 7$, then we may replace condition (1.1b) and (1.2b) by

$$a \geq 2 \quad \text{and} \quad b \in \left[ea + e - 1, \left(2 + \frac{e}{2}\right) \cdot a + \sqrt{3a^2 + 12a} + 2\right]. \quad (2.14)$$

2.4 Corollary

Let $\Sigma = \mathbb{F}_e$ with $2 \leq e \leq 7$, let $D = aC_0 + bF \in \text{Div}(\Sigma)$ with $a \geq 2$ and $b \in [ea + e - 1, \left(2 + \frac{e}{2}\right) \cdot a + \sqrt{3a^2 + 12a} + 2[$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < (2ab - ea^2) + 4b + (4 - 2e) \cdot a - \left(b - \frac{e+2}{2} \cdot a\right)^2. \quad (2.15)$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

Next we consider the case of an elliptic base curve. As in the case of Hirzebruch surfaces, we study Condition (1.2b) and Obstruction (2.4) a bit closer. We then see that they may be replaced by

$$b \geq ea + e + 1 \quad (2.16)$$

and

$$\psi_{a,e}(b) := -\left(b - \frac{e}{2} \cdot a\right)^2 + (2ab - ea^2) + 4b - 2ea - 4 > 0. \quad (2.17)$$

For $a \geq 2$ fixed we consider $\psi_{a,e} : [ea + e + 1, \infty[\rightarrow \mathbb{R}$ as a concave function, and we find that

$$\psi_{a,e}(b) > 0 \iff b < a + \frac{e}{2} \cdot a + \sqrt{a^2 + 4a} + 2.$$

Thus (2.16) and (2.17) will never be satisfied at the same time if $e \geq 4$, while if $0 \leq e \leq 3$ we may replace (1.1b) and (1.2b) by

$$a \geq 2 \quad \text{and} \quad b \in \left[ea + e + 1, a + \frac{e}{2} \cdot a + \sqrt{a^2 + 4a} + 2 \right]. \quad (2.18)$$

2.5 Corollary

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with $g(C) = 1$ and $0 \leq e \leq 3$, let $D = aC_0 + bF \in \text{Div}(\Sigma)$ with $a \geq 2$ and $b \in \left[ea + e + 1, a + \frac{e}{2} \cdot a + \sqrt{a^2 + 4a} + 2 \right]$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < (2ab - ea^2) + 4b - 2ea - \left(b - \frac{e}{2} \cdot a\right)^2. \quad (2.19)$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

We note that in the most interesting case, namely when $e = 0$, which includes the product $C \times \mathbb{P}_C^1$, we just need

$$0 < b \leq a + \sqrt{a^2 + 4a} + 2 \quad (2.20)$$

and

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 2ab + 4b - b^2. \quad (2.21)$$

Finally we have to consider the unique case of a geometrically ruled surface over an elliptic curve with negative invariant e . (1.1) and (1.2) come down to

$$a \geq 2 \quad \text{and} \quad b > -\frac{a}{2}. \quad (2.22)$$

Moreover, then $(aC_0 + bF) \cdot K_\Sigma = -2b - a < 0$, so that Remark 1.2 (c) still applies, that is, (1.3) is redundant and (1.5) may be replaced by (1.17), which takes the form

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < \frac{3}{4} \cdot a^2 + ab + 2a + 4b - b^2.$$

This, however, leads to the obstruction

$$\psi_{a,-1}(b) = \frac{3}{4} \cdot a^2 + ab + 2a + 4b - b^2 - 4 > 0.$$

Thus, considering $\psi_{a,-1}$ now as a concave function $\psi_{a,-1} :] - \frac{a}{2}, \infty[\rightarrow \mathbb{R}$ we find that

$$\psi_{a,-1}(b) > 0 \iff -\frac{a}{2} < \frac{a}{2} + 2 - \sqrt{a^2 + 4a} < b < \frac{a}{2} + 2 + \sqrt{a^2 + 4a}.$$

And we may therefore replace (1.1b)-(1.2b) by

$$a \geq 2 \quad \text{and} \quad b \in \left] \frac{a}{2} + 2 - \sqrt{a^2 + 4a}, \frac{a}{2} + 2 + \sqrt{a^2 + 4a} \right[. \quad (2.23)$$

2.6 Corollary

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with $g(C) = 1$ and $e = -1$, let $D = aC_0 + bF \in \text{Div}(\Sigma)$ with $a \geq 2$ and $b \in]\frac{a}{2} + 2 - \sqrt{a^2 + 4a}, \frac{a}{2} + 2 + \sqrt{a^2 + 4a}[$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < \frac{3}{4} \cdot a^2 + ab + 2a + 4b - b^2. \quad (2.24)$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

In the above corollary we could for instance let $b = \frac{a}{2} + 2$, in which case (2.19) reads

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < (a + 2)^2.$$

2.c. Products of Curves

Throughout this section we use the notation of Section G.b. In particular, $\Sigma = C_1 \times C_2$ where C_1 and C_2 are smooth projective curves over \mathbb{C} of genera g_1 and g_2 respectively. Since Σ is geometrically ruled if some $g_i = 0$, we may restrict our attention to the case $g_1, g_2 \geq 1$.

For a generic choice of C_1 and C_2 the Néron–Severi group $\text{NS}(\Sigma)$ is two-dimensional by Proposition G.12. Thus the following theorem answers the general case completely.

1.1c Theorem

Let C_1 and C_2 be two smooth projective curves of genera g_1 and g_2 respectively with $g_1 \geq g_2 \geq 1$.

Let $D = aC_1 + bC_2 \in \text{Div}(\Sigma)$ with $a > 2g_2 - 2$ and $b > 2g_1 - 2$, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types with $\tau^*(\mathcal{S}_1) \geq \tau^*(\mathcal{S}_i)$ for all $i = 1, \dots, r$.

Suppose that

$$(1.5c) \quad \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 2ab - 2 \cdot (\tau^*(\mathcal{S}_1) + 1) \cdot ((g_1 - 1)a + (g_2 - 1)b) - ((g_1 - 1)a - (g_2 - 1)b)^2.$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

Proof: We note that $K_\Sigma = (2g_2 - 2) \cdot C_1 + (2g_1 - 2) \cdot C_2$ is nef. Thus by Remark 1.2 (b) and (d) and due to the assumptions on a and b (1.1)-(1.3) are fulfilled, while (1.5) may be replaced by (1.18), which in this case is just (1.5c). It thus remains to show that (1.4) is satisfied, which in this case takes the form

$$\sum_{i=1}^r \tau^*(\mathcal{S}_i) < \frac{1}{2} \cdot (a + 2 - 2g_2) \cdot (b + 2 - 2g_1).$$

This, however, follows from (1.24) in Remark 1.2 since

$$\begin{aligned} & 2ab - 2 \cdot (\tau^*(\mathcal{S}_1) + 1) \cdot ((g_1 - 1)a + (g_2 - 1)b) - ((g_1 - 1)a - (g_2 - 1)b)^2 \\ & \leq 2ab - 4 \cdot ((g_1 - 1)a + (g_2 - 1)b) \leq 2 \cdot (a + 2 - 2g_2) \cdot (b + 2 - 2g_1). \end{aligned}$$

□

Of course, if $(g_1 - 1) \cdot a$ and $(g_2 - 1) \cdot b$ differ too much from each other, then Condition (1.5c) can never be satisfied, since then $((g_1 - 1) \cdot a - (g_2 - 1) \cdot b)^2$ becomes larger than $2ab$ - e. g. if $g_1, g_2 > 1$ and $a > 4 \cdot \frac{g_2 - 1}{g_1 - 1} \cdot b$. Moreover, we see that the largest $\tau^*(\mathcal{S}_i)$ should be considerably smaller than a or b .

If $g_1, g_2 > 1$ and if $a = p \cdot (g_2 - 1)$ and $b = p \cdot (g_1 - 1)$ for some $p \in \mathbb{Q}$, that is if D is a multiple of the ample divisor K_Σ then (1.5c) takes the simpler form

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < 2 \cdot (p^2 - 2p \cdot (\tau^*(\mathcal{S}_1) + 1)) \cdot (g_1 - 1) \cdot (g_2 - 1). \quad (2.25)$$

In this case, however, we could replace (2.25) in view of Remark 1.2 (f) also by

$$\sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{4}{p} \cdot (\tau^*(\mathcal{S}_i) + 1)} < 2p^2 \cdot (g_1 - 1) \cdot (g_2 - 1). \quad (2.26)$$

2.d. Products of Elliptic Curves

If in Section V.4.c the curves C_1 and C_2 are chosen to be both elliptic curves, Theorem 1.1c looks much nicer, since $K_\Sigma = 0$.

1.1d Theorem

Let C_1 and C_2 be two smooth elliptic curves, let $D \in \text{Div}(\Sigma)$ big and nef, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$(1.5d) \quad \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < D^2.$$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T-smooth.

If C_1 and C_2 are non-isogenous, then any divisor of Σ is of the form $D = aC_1 + bC_2$, and for such a divisor the condition “ D big and nef” means that $a, b > 0$.

The Severi variety $V_{|aC_1 + bC_2|}^{\text{irr}}(rA_1)$ is therefore T-smooth as soon as

$$r \leq \frac{ab}{2},$$

and in the case of cusps and nodes the condition for the T-smoothness of $V_{|aC_1 + bC_2|}^{\text{irr}}(kA_1 + mA_2)$ thus just reads

$$4k + 9m < 2ab.$$

2.e. Surfaces in $\mathbb{P}_\mathbb{C}^3$

For a generic surface of degree $n \geq 4$ in $\mathbb{P}_\mathbb{C}^3$ any curve is a hypersurface section by a Theorem of Noether (see Section G.d). Therefore the following result on hypersurface sections answers the problem completely on a generic surface in $\mathbb{P}_\mathbb{C}^3$. We note that a hypersurface section is always a rational multiple of the canonical bundle and thus we may use Remark 1.2 (b) and (1.20).

1.1e Theorem

Let $\Sigma \subset \mathbb{P}_\mathbb{C}^3$ be a smooth hypersurface of degree $n \geq 4$, $H \subset \Sigma$ be a hyperplane section, $d > n - 4$ and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types with $\tau^*(\mathcal{S}_1) \geq \tau^*(\mathcal{S}_i)$ for all $i = 1, \dots, r$. Suppose that

$$(1.4e) \quad \sum_{i=1}^r \tau^*(\mathcal{S}_i) < \frac{(d+4-n)^2 \cdot n}{4}, \text{ and}$$

$$(1.5e) \quad \sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 1)^2 < d^2 \cdot n - d \cdot (n - 4) \cdot n \cdot (\tau^*(\mathcal{S}_1) + 1).$$

Then either $V_{|dH|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or T -smooth.

If we suppose $n > 4$ and the $\tau^*(\mathcal{S}_i) + 1 < \frac{d}{n-4}$, which is necessary anyway in order to have (1.5e), then in view of Remark 1.2 (f) we may replace (1.5e) by the better inequality

$$\sum_{i=1}^r \frac{(\tau^*(\mathcal{S}_i) + 1)^2}{1 - \frac{n-4}{d} \cdot (\tau^*(\mathcal{S}_i) + 1)} < d^2 \cdot n. \quad (2.27)$$

Let us consider the two cases of nodal curves and of curves with nodes and cusps more closely.

2.6 Corollary

Let $\Sigma \subset \mathbb{P}_\mathbb{C}^3$ be a smooth hypersurface of degree $n \geq 4$, $H \subset \Sigma$ be a hyperplane section and $d > 2n - 8$. Suppose that

$$r < \frac{d \cdot (d - 2n + 8) \cdot n}{4}. \quad (2.28)$$

Then either $V_{|dH|}^{\text{irr}}(\tau A_1)$ is empty or T -smooth.

Proof: In this situation (1.5e) is just (2.28), and (1.4e) is fulfilled as well in view of

$$\frac{d \cdot (d - 2n + 8) \cdot n}{4} \leq \frac{d \cdot (d - 2n + 8) \cdot n + (4 - n)^2 n}{4} = \frac{(d + 4 - n)^2 \cdot n}{4}. \quad \square$$

2.7 Corollary

Let $\Sigma \subset \mathbb{P}_\mathbb{C}^3$ be a smooth hypersurface of degree $n \geq 4$, $H \subset \Sigma$ be a hyperplane section and $d > 3n - 12$. Suppose that

$$4k + 9m < d \cdot (d - 3n + 12) \cdot n. \quad (2.29)$$

Then either $V_{|dH|}^{\text{irr}}(kA_1, mA_2)$ is empty or T -smooth.

Proof: In this situation (1.5e) is just (2.29), and (1.4e) is fulfilled as well in view of

$$4 \cdot (k + 2m) \leq 4k + 9m < d \cdot (d - 3n + 12) \cdot n \leq (d + 4 - n)^2 \cdot n.$$

□

2.f. K3-Surfaces

By Remark 1.2 (b), (c) and (j) Theorem 1.1 for K3-surfaces takes the following form.

1.1f Theorem

Let Σ be a smooth K3-surface, D a big and nef divisor on Σ and S_1, \dots, S_r be topological or analytical singularity types.

Suppose that

$$(1.5f) \quad \sum_{i=1}^r (\tau^*(S_i) + 1)^2 < D^2.$$

Then either $V_{|D|}^{\text{irr}}(S_1, \dots, S_r)$ is empty or T -smooth.

CHAPTER V

Irreducibility

The question whether the variety $V^{\text{irr}} = V_{|\mathcal{D}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ of irreducible curves in $|\mathcal{D}|_1$ having precisely r singular points of the given types $\mathcal{S}_1, \dots, \mathcal{S}_r$ is irreducible seems to be much harder to answer suitably than that of existence or smoothness. Even in the simplest case of nodal curves in $\mathbb{P}_{\mathbb{C}}^2$ the conjecture of Severi that V^{irr} should be irreducible whenever it is non-empty resisted a full proof until 1985 (cf. [Har85b]), while already Severi himself (cf. [Sev21]) gave a complete characterisation for the existence and showed that the variety is always T-smooth. Moreover, while for the question of existence and smoothness in the plane curve case there exist sufficient criteria showing the same asymptotical behaviour as known necessary criteria, respectively as known (series of) non-smooth families, the asymptotics for the sufficient criteria for irreducibility seem to be worse (cf. [Los98] Chapter 6). To be more precise there are conditions linear in certain invariants and quadratic in the degree which guarantee the existence, while for the irreducibility (as for the T-smoothness) the conditions are quadratic in the degree and the invariants as well. Applying similar techniques our results carry the same stigma. However, apart from nodal curves on the blown up plane (cf. [Ran89, GLS98a]) we do not know at all of any criteria for the irreducibility–problem on surfaces other than the plane. (See also Section 4.f).

The main condition for irreducibility which we get in the different cases looks like

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_{\Sigma})^2, \quad (0.1)$$

where γ is some constant.

At this point we should like to point out that on more complicated surfaces than $\mathbb{P}_{\mathbb{C}}^2$ we cannot expect the results to be as nice as in the plane case. Already Harris' result for the variety of nodal curves does not extend to arbitrary surfaces Σ – more precisely, to our knowledge there is no surface known, except the plane, where the result holds. In [ChC99] it is shown that on a generic surface in $\mathbb{P}_{\mathbb{C}}^3$ of degree $n \geq 8$ the variety $V_{|\mathcal{D}|}^{\text{irr}}(rA_1)$ is reducible for all even $d > 0$ and $r = \frac{(d^2 + (4-n)d + 2) \cdot n}{2}$ as soon as

$$\frac{n^3 - 12n^2 + 11n - 6}{6} < \frac{dn - d - 2}{2},$$

where the left hand side is the expected dimension of $V_{|dH|}^{\text{irr}}(\tau A_1)$ and the right hand side is a lower bound for the actual dimension. This lower bound is found by intersecting Σ with a family of cones with fixed vertex over plane nodal curves of degree d with $\frac{r}{n}$ nodes. The right hand side is the dimension of the family of plane nodal curves, and we thus only have to note that cones over different curves are different. The arguments given so far show that $V_{|dH|}^{\text{irr}}(\tau A_1)$ has a component which is not T -smooth and works for any surface of degree $d \geq 8$ in \mathbb{P}_c^3 , however, in [ChC99] Theorem 3.1 Chiantini and Ciliberto show that on a generic surface in \mathbb{P}_c^3 the variety $V_{|dH|}^{\text{irr}}(\tau A_1)$ with $d \geq n$ also contains a T -smooth component whenever it is non-empty.

For an overview on the different approaches to the question of irreducibility in the case of plane curves we refer to [GrS99] or [Los98] Chapter 6 – see also [BrG81, Ura83, GD84, Ura84, Har85b, Dav86, Ura86, Deg87, GrK89, Kan89a, Ran89, Shu91a, Bar93a, Shu94, Shu96b, Wal96, Bar98, Mig01]. The case of \mathbb{P}_c^2 blown up in k generic points is treated in [Ran89], if $k = 1$, and in [GLS98a] in the general case. Our proof proceeds along the lines of an unpublished result of Greuel, Lossen and Shustin (cf. [GLS98b]). The basic ideas are in some respect similar to the approach utilised in [GLS00], replacing the “Castelnuovo-function” arguments by “Bogomolov unstability”. We tackle the problem in three steps:

Step 1: We first show that the open subvariety $V^{\text{irr,reg}}$ of curves in V^{irr} with $h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) = 0$ is always irreducible, and hence so is its closure in V^{irr} . (Cf. Corollary 1.2.)

Step 2: Then we find conditions which ensure that the open subvariety $V^{\text{irr,fix}}$ of curves in V^{irr} with $h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D)) = 0$ is dense in V^{irr} . (Cf. Section 2.)

Step 3: And finally, we combine these conditions with conditions which guarantee that $V^{\text{irr,reg}}$ is dense in $V^{\text{irr,fix}}$ by showing that they share some open dense subset $V_{\mathcal{U}}^{\text{gen}}$ of curves with singularities in very general position (cf. Lemma 3.9). But then $V^{\text{irr,reg}}$ is dense in V^{irr} and V^{irr} is irreducible by Step 1.

The most difficult part is Step 2. For this one we consider the restriction of the morphism (cf. Definition I.2.15)

$$\Phi : V \rightarrow \text{Sym}^{k_1 + \dots + k_r}(\Sigma) =: \mathcal{B}$$

to an irreducible component V^* of V^{irr} not contained in the closure $\overline{V^{\text{irr,fix}}}$ in V^{irr} . Knowing, that the dimension of V^* is at least the expected dimension $\dim(V^{\text{irr,fix}})$ we deduce that the codimension of $\mathcal{B}^* = \Phi(V^*)$ in \mathcal{B} is at most $h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D))$, where $C \in V^*$ (cf. Lemma 3.7). It thus suffices to find conditions which contradict this inequality, that is, we have to get our hands on $\text{codim}_{\mathcal{B}}(\mathcal{B}^*)$. However, on the surfaces, which we consider, the non-vanishing of $h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D))$ means in some sense that the zero-dimensional scheme

$X_{\text{fix}}^*(C)$ is in special position. We may thus hope to realise large parts X_i^0 of $X_{\text{fix}}^*(C)$ on curves Δ_i of “small degree” ($i = 1, \dots, m$), which would impose at least $\#X_i^0 - \dim |\Delta_i|_1$ conditions on $X_{\text{fix}}^*(C)$, giving rise to a lower bound $\sum_{i=1}^m \#X_i^0 - \dim |\Delta_i|_1$ for $\text{codim}_{\mathcal{B}}(\mathcal{B}^*)$. The X_i^0 and the Δ_i are found in Lemma 3.1 with the aid of certain Bogomolov unstable rank-two bundles. It thus finally remains (cf. Lemma 3.3, 3.4 and 3.6) to give conditions which imply

$$\sum_{i=1}^m \#X_i^0 - \dim |\Delta_i|_1 > h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D)).$$

What are the obstructions by which our approach is restricted?

First, the Bogomolov unstability does not give us much information about the curves Δ_i apart from their existence and the fact that they are in some sense “small” compared with the divisor D . It thus is obvious that we are bound to the study of surfaces where we have a good knowledge of the dimension of arbitrary complete linear systems. Second, in order to derive the above inequality many nasty calculations are necessary which depend very much on the particular structure of the Néron–Severi group of the surface, that is, we are restricted to surfaces where the Néron–Severi group is not too large and the intersection pairing is not too hard (cf. Lemma 3.3, 3.4 and 3.6). Finally, in order to ensure the Bogomolov unstability of the vector bundle considered throughout the proof of Lemma 3.1 we heavily use the fact that the surface Σ does not contain any curves of negative self-intersection, which excludes e. g. general Hirzebruch surfaces.

In Section 1 we do not only prove that $V^{\text{irr,reg}}$ is irreducible, but that this indeed remains true if we drop the requirement that the curves should be irreducible, i. e. we show that V^{reg} is irreducible. However, unfortunately our approach does not give conditions implying that V^{reg} is dense, and thus we cannot say anything about the irreducibility of the variety of possibly reducible curves in $|D|_1$ with prescribed singularities.¹ The reason for this is that in the proof of Lemma 3.1 we use the Theorem of Bézout to estimate $D \cdot \Delta_i$. Since Δ_i may be about “half” of D , we need an irreducible curve in $|D|_1$ to be sure that at least for some curve in $|D|_1$ the curve Δ_i is not a component.

1. $V^{\text{irr,reg}}$ is irreducible

We are now showing that V^{reg} and $V^{\text{irr,reg}}$ are always irreducible. We do this by showing that under $\Psi : V \rightarrow B$ (cf. Definition I.2.15) every irreducible component of these is smooth and maps dominant to the irreducible variety B with irreducible fibres.

¹Note that e. g. the variety $V_{|2H|}^{\text{irr}}(A_1)$ of irreducible plane conics with one node is empty, while $V_{|2H|}(A_1)$ is four-dimensional – even though the latter is of course irreducible.

1.1 Theorem

Let $D \in \text{Div}(\Sigma)$, $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

If $V_{|D|}^{\text{reg}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is non-empty, then it is a T -smooth, irreducible, open subset of $V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ of dimension $\dim |D|_{\mathbb{1}} - \sum_{i=1}^r k_i \cdot \tau^*(\mathcal{S}_i)$.

Proof: Let us consider the following maps from Definition I.2.15

$$\Psi = \Psi_D(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) : V = V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) \longrightarrow B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$$

and

$$\psi = \psi(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) : B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) \longrightarrow \text{Hilb}_{\Sigma}^n.$$

Step 1: Every irreducible component V^* of V^{reg} is T -smooth of dimension $\dim |D|_{\mathbb{1}} - \sum_{i=1}^r k_i \cdot \tau^*(\mathcal{S}_i)$.

By [Los98] Proposition 2.1 (c2) V^* is T -smooth at any $C \in V^*$ of dimension $\dim |D|_{\mathbb{1}} - \deg(X^*(C)) = \dim |D|_{\mathbb{1}} - \sum_{i=1}^r k_i \cdot \tau(\mathcal{S}_i)$, since $h^1(\Sigma, \mathcal{J}_{X^*/\Sigma}(D)) = 0$ according to Remark I.2.8. (See also Remark I.2.10.)

Step 2: V^{reg} is open in V .

Let $C \in V^{\text{reg}}$. By assumption $h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) = 0$, and thus by Lemma F.1 there exists an open, dense neighbourhood \mathcal{U} of $X(C)$ in Hilb_{Σ}^n such that $h^1(\Sigma, \mathcal{J}_{Y/\Sigma}(D)) = 0$ for all $Y \in \mathcal{U}$. But then $\Psi^{-1}(\psi^{-1}(\mathcal{U})) \subseteq V^{\text{reg}}$ is an open neighbourhood of $X(C)$ in V , and hence V^{reg} is open in V .

Step 3: $\Psi = \Psi_D(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ restricted to any irreducible component V^* of V^{reg} is dominant.

Let V^* be an irreducible component of V^{reg} and let $C \in V^*$. Since $\Psi^{-1}(\Psi(C))$ is an open and dense subset of $| \mathcal{J}_{X(C)/\Sigma}(D) |_{\mathbb{1}}$ and since $h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) = 0$, we have

$$\dim \Psi^{-1}(\Psi(C)) = h^0(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) - 1 = \dim |D|_{\mathbb{1}} - \deg(X(C)).$$

By Step 1 we know the dimension of V^* and by a remark in Definition I.2.15 we also know the dimension of B . Thus we conclude

$$\begin{aligned} \dim \Psi(V^*) &= \dim V^* - \dim \Psi^{-1}(\Psi(C)) \\ &= (\dim |D|_{\mathbb{1}} - \deg X^*(C)) - (\dim |D|_{\mathbb{1}} - \deg X(C)) \\ &= \deg(X(C)) - \deg(X^*(C)) = \dim B. \end{aligned}$$

Since B is irreducible, $\Psi(V^*)$ must be dense in B .

Step 4: V^{reg} is irreducible.

Let V^* and V^{**} be two irreducible components of V^{reg} . Then $\Psi(V^*) \cap \Psi(V^{**}) \neq \emptyset$, and thus some fibre F of Ψ intersects both, V^* and V^{**} . However, the fibre is irreducible and by Step 1 both V^* and V^{**} are smooth. Thus F must be

completely contained in V^* and V^{**} , which, since both are smooth of the same dimension, implies that $V^* = V^{**}$. That is V^{reg} is irreducible. \square

1.2 Corollary

Let $D \in \text{Div}(\Sigma)$, $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

If $V_{|D|}^{\text{irr,reg}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is non-empty, then it is a T -smooth, irreducible, open subset of $V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ of dimension $\dim |D|_1 - \sum_{i=1}^r k_i \cdot \tau^*(\mathcal{S}_i)$.

Proof: This follows from Theorem 1.1 since $V_{|D|}^{\text{irr,reg}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is an open subset of the irreducible set $V_{|D|}^{\text{reg}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$. \square

2. Conditions for the irreducibility of V^{irr}

Knowing that $V^{\text{irr,reg}}$, and hence its closure $\overline{V^{\text{irr,reg}}}$ in V^{irr} , is always irreducible, the search for sufficient conditions for the irreducibility of V^{irr} may be reduced to the search for conditions ensuring that $V^{\text{irr,reg}}$ is dense in V^{irr} . As indicated in the introduction (cf. p. 82) we achieve this aim by combining conditions, which ensure that $V^{\text{irr,reg}}$ and $V^{\text{irr,fix}}$ share some dense subset $V_{\mathcal{U}}^{\text{gen}}$ (cf. Step 3 on page 82), with conditions which ensure that $V^{\text{irr,fix}}$ is dense in V^{irr} (cf. Step 2 on page 82). For the latter problem we apply a reduction technique involving the Bogomolov unstability of certain rank-two vector bundles on Σ . The main part of the work is carried out in Section 3, where we prove several technical lemmata which partly are useful in their own respect.

Let us now reformulate Step 2 and Step 3 in a more precise way.

Step 2: We derive from Lemma 3.1-3.7 conditions which ensure that $V^{\text{irr,fix}}$ is dense in V^{irr} .

Step 3: Taking Lemma 3.8 into account, we deduce from Lemma 3.9 conditions which ensure that there exists a very general subset $\mathcal{U} \subset \Sigma^m$ with $m = k_1 + \dots + k_r$ such that the family $V_{\mathcal{U}}^{\text{gen}} = V_{|D|,\mathcal{U}}^{\text{gen}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$, as defined there, satisfies

- (a) $V_{\mathcal{U}}^{\text{gen}}$ is dense in $V^{\text{irr,fix}}$, and
- (b) $V_{\mathcal{U}}^{\text{gen}} \subseteq V^{\text{irr,reg}}$.

Thus in particular, $V^{\text{irr,reg}}$ is dense in $V^{\text{irr,fix}}$.

Throughout the proof of the following theorem we will heavily rely on the study of the zero-dimensional schemes $X_{\text{fix}}^*(C)$ for some curves C , and we therefore would like to remind the reader that in view of Definition I.2.10 for a topological respectively analytical singularity type \mathcal{S} we have

$$\deg(X_{\text{fix}}^*(\mathcal{S}_i)) = \tau^*(\mathcal{S}_i) + 2.$$

2.1 Theorem

Let Σ be a surface such that²

- (i) $\text{NS}(\Sigma) = L \cdot \mathbb{Z}$,
- (ii) $h^1(\Sigma, C) = 0$, whenever C is effective.

Let $D \in \text{Div}(\Sigma)$, let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

Suppose that

$$(2.1) \quad D - K_\Sigma \text{ is big and nef,}$$

$$(2.2) \quad D + K_\Sigma \text{ is nef,}$$

$$(2.3) \quad \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2) < \beta \cdot (D - K_\Sigma)^2 \text{ for some } 0 < \beta \leq \frac{1}{4}, \text{ and}$$

$$(2.4) \quad \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2, \text{ where } \gamma = \frac{(1 + \sqrt{1 - 4\beta})^2 \cdot L^2}{4 \cdot \chi(\mathcal{O}_\Sigma) + \max\{0, 2 \cdot K_\Sigma \cdot L\} + 6 \cdot L^2}.$$

Then either $V_{|D|}^{\text{irr}}(k_1 \mathcal{S}_1, \dots, k_r \mathcal{S}_r)$ is empty or it is irreducible of the expected dimension.

Proof: We may assume that $V^{\text{irr}} = V_{|D|}^{\text{irr}}(k_1 \mathcal{S}_1, \dots, k_r \mathcal{S}_r)$ is non-empty. As indicated above it suffices to show that:

Step 2: $V^{\text{irr}} = \overline{V^{\text{irr}, \text{fix}}}$, where $V^{\text{irr}, \text{fix}} = V_{|D|}^{\text{irr}, \text{fix}}(k_1 \mathcal{S}_1, \dots, k_r \mathcal{S}_r)$, and

Step 3: the conditions of Lemma 3.9 are fulfilled.

For Step 3 we note that $\nu^*(\mathcal{S}_i) \leq \tau^*(\mathcal{S}_i)$. Thus (2.4) implies that

$$\sum_{i=1}^r k_i \cdot (\nu^*(\mathcal{S}_i) + 1)^2 \leq \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2 \leq \gamma \cdot (D - K_\Sigma)^2 \leq \frac{1}{2} \cdot (D - K_\Sigma)^2,$$

which gives the first condition in Lemma 3.9. Since a surface with Picard number one has no curves of selfintersection zero, the second condition in Lemma 3.9 is void, while the last condition is satisfied by (2.1).

It remains to show Step 2, i. e. $V^{\text{irr}} = \overline{V^{\text{irr}, \text{fix}}}$. Suppose the contrary, that is, there is an irreducible curve $C_0 \in V^{\text{irr}} \setminus \overline{V^{\text{irr}, \text{fix}}}$, in particular $h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) > 0$ for $X_0 = X_{\text{fix}}^*(C_0)$. Since $\deg(X_0) = \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)$ and $\sum_{z \in \Sigma} (\deg(X_{0,z}))^2 = \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2$ the assumptions (0)-(3) of Lemma 3.1 and (4) of Lemma 3.3 are fulfilled. Thus Lemma 3.3 implies that C_0 satisfies Condition (3.28) in Lemma 3.7, which it cannot satisfy by the same Lemma. Thus we have derived a contradiction. \square

²By Lemma E.1 we may assume w. l. o. g. that L is ample.

2.2 Remark

If we set

$$\gamma = \frac{36\alpha}{(3\alpha+4)^2} \quad \text{with} \quad \alpha = \frac{4 \cdot \chi(\mathcal{O}_\Sigma) + \max\{0, 2 \cdot K_\Sigma \cdot L\} + 6 \cdot L^2}{L^2},$$

then a simple calculation shows that (2.3) becomes redundant. For this we have to take into account that $\tau^*(\mathcal{S}) \geq 1$ for any singularity type \mathcal{S} . The claim then follows with $\beta = \frac{1}{3} \cdot \gamma \leq \frac{1}{4}$.

2.3 Theorem

Let C_1 and C_2 be two smooth projective curves of genera g_1 and g_2 respectively with $g_1 \geq g_2 \geq 0$, such that for $\Sigma = C_1 \times C_2$ the Néron–Severi group is $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z}$.

Let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_1 + bC_2$ with $a > \max\{2 - 2g_2, 2g_2 - 2 + \nu^*(\mathcal{S}_i) \mid i = 1, \dots, r\}$ and $b > \max\{2 - 2g_1, 2g_1 - 2 + \nu^*(\mathcal{S}_i) \mid i = 1, \dots, r\}$, let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and let $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

Suppose that

$$\sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2, \quad (2.5)$$

where γ may be taken from the following table with $\alpha = \frac{a-2g_2+2}{b-2g_1+2} > 0$.

g_1	g_2	γ	γ , if $\alpha = 1$
0	0	$\frac{1}{24}$	$\frac{1}{24}$
1	0	$\frac{1}{\max\{32, 2\alpha\}}$	$\frac{1}{32}$
≥ 2	0	$\frac{1}{\max\{24+16g_1, 4g_1\alpha\}}$	$\frac{1}{24+16g_1}$
1	1	$\frac{1}{\max\{32, 2\alpha, \frac{2}{\alpha}\}}$	$\frac{1}{32}$
≥ 2	≥ 1	$\frac{1}{\max\{24+16g_1+16g_2, 4g_1\alpha, \frac{4g_2}{\alpha}\}}$	$\frac{1}{24+16g_1+16g_2}$

Then $V_{|D|}^{\text{irr}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is empty or it is irreducible of the expected dimension.

Proof: The assumptions on a and b ensure that $D - K_\Sigma$ is big and nef and that $D + K_\Sigma$ is nef. Thus, once we know that (2.5) implies Condition (3) in Lemma 3.1, Step 2 in the proof of Theorem 2.1 follows in the same way, just replacing Lemma 3.3 by Lemma 3.4.

For Condition (3) we note that

$$\sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2) \leq \sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2 \leq \frac{1}{24} \cdot (D - K_\Sigma)^2 < \frac{1}{4} \cdot (D - K_\Sigma)^2.$$

Also Step 3 in the proof of Theorem 2.1 follows in the same way, except that Condition (b) in Lemma 3.9 is not void this time, but is fulfilled by the assumptions on a and b , since the only irreducible curves of selfintersection zero are linearly equivalent either to C_1 or to C_2 . \square

In the following theorem we use the notation of Section G.a.

2.4 Theorem

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with $e \leq 0$ and $g = g(C)$.

Let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_0 + bF$ with $a > \max\{1, \nu^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}$, $b > 2g - 2 + \frac{ae}{2} + \max\{\nu^*(\mathcal{S}_i) \mid i = 1, \dots, r\}$, and if $g = 0$ then $b \geq 2$. Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$.

Suppose that

$$\sum_{i=1}^r k_i \cdot (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2, \quad (2.6)$$

where γ may be taken from the following table with $\alpha = \frac{a+2}{b+2-2g-\frac{ae}{2}} > 0$.

g	e	γ	γ , if $\alpha = 1$
0	0	$\frac{1}{24}$	$\frac{1}{24}$
1	0	$\frac{1}{\max\{24, 2\alpha\}}$	$\frac{1}{24}$
1	-1	$\frac{1}{\max\left\{\min\left\{30 + \frac{16}{\alpha} + 4\alpha, 40 + 9\alpha\right\}, \frac{13}{2}\alpha\right\}}$	$\frac{1}{49}$
≥ 2	0	$\frac{1}{\max\{24 + 16g, 4g\alpha\}}$	$\frac{1}{24 + 16g}$
≥ 2	< 0	$\frac{1}{\max\left\{\min\left\{24 + 16g - 9e\alpha, 18 + 16g - 9e\alpha - \frac{16}{e\alpha}\right\}, 4g\alpha - 9e\alpha\right\}}$	

Then $V_{|D|}^{\text{irr}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is empty or it is irreducible of the expected dimension.

Proof: The proof is identical to that of Theorem 2.3, just replacing Lemma 3.4 by Lemma 3.6 and applying Lemma G.2 for the irreducible curves of selfintersection zero. \square

3. The Main Technical Lemmata

The following lemma is the heart of the proof. Given a curve $C \in |D|_1$ such that the scheme $X_0 = X_{\text{fix}}^{\text{es}}(C)$ respectively $X_0 = X_{\text{fix}}^{\text{ea}}(C)$ is *special with respect to D* in the sense that $h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) > 0$, it provides a “small” curve Δ_1 through a subscheme X_1^0 of X_0 , so that we can reduce the problem by replacing X_0 and D by $X_0 : \Delta_1$ and $D - \Delta_1$ respectively. We can of course proceed inductively as long as the new zero-dimensional scheme is again special with respect to the new divisor.

In order to find Δ_1 we choose a subscheme $X_1^0 \subseteq X_0$ which is minimal among those subschemes special with respect to D . By Grothendieck-Serre duality

$$H^1(\Sigma, \mathcal{J}_{X_1^0/\Sigma}(D)) \cong \text{Ext}^1(\mathcal{J}_{X_1^0/\Sigma}(D - K_\Sigma), \mathcal{O}_\Sigma)$$

and a non-trivial element of the latter group gives rise to an extension

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow E_1 \rightarrow \mathcal{J}_{X_1^0/\Sigma}(D - K_\Sigma) \rightarrow 0.$$

We then show that the rank-two bundle E_1 is Bogomolov unstable and deduce the existence of a divisor Δ_1^0 such that

$$H^0(\Sigma, \mathcal{J}_{X_1^0/\Sigma}(D - K_\Sigma - \Delta_1^0)) \neq 0,$$

that is, we find a curve $\Delta_1 \in |\mathcal{J}_{X_1^0/\Sigma}(D - K_\Sigma - \Delta_1^0)|_1$.

3.1 Lemma

Let Σ be a surface such that

(*) any curve $C \subset \Sigma$ is nef.

Let $D \in \text{Div}(\Sigma)$ and $X_0 \subset \Sigma$ a zero-dimensional scheme satisfying

- (0) $D - K_\Sigma$ is big and nef, and $D + K_\Sigma$ is nef,
- (1) $\exists C_0 \in |D|_1$ irreducible : $X_0 \subset C_0$,
- (2) $h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) > 0$, and
- (3) $\deg(X_0) < \beta \cdot (D - K_\Sigma)^2$ for some $0 < \beta \leq \frac{1}{4}$.

Then there exist curves $\Delta_1, \dots, \Delta_m \subset \Sigma$ and zero-dimensional locally complete intersections $X_i^0 \subseteq X_{i-1} \cap \Delta_i$ for $i = 1, \dots, m$, where $X_i = X_{i-1} : \Delta_i$ for $i = 1, \dots, m$, such that

$$(a) \quad h^1(\Sigma, \mathcal{J}_{X_m/\Sigma}(D - \sum_{i=1}^m \Delta_i)) = 0,$$

and for $i = 1, \dots, m$

- (b) $h^1(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)) = 1$
- (c) $D \cdot \Delta_i \geq \deg(X_{i-1} \cap \Delta_i) \geq \deg(X_i^0) \geq (D - K_\Sigma - \sum_{k=1}^i \Delta_k) \cdot \Delta_i \geq \Delta_i^2 \geq 0$
- (d) $(D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i)^2 > 0$,
- (e) $(D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i) \cdot H > 0$ for all $H \in \text{Div}(\Sigma)$ ample, and
- (f) $D - K_\Sigma - \sum_{k=1}^i \Delta_k$ is big and nef.

Moreover, it follows

$$0 \leq \frac{1}{4}(D - K_\Sigma)^2 - \sum_{i=1}^m \deg(X_i^0) \leq \left(\frac{1}{2}(D - K_\Sigma) - \sum_{i=1}^m \Delta_i \right)^2. \quad (3.1)$$

Proof: We are going to find the Δ_i and X_i^0 recursively. Let us therefore suppose that we have already found $\Delta_1, \dots, \Delta_{i-1}$ and X_1^0, \dots, X_{i-1}^0 satisfying (b)-(f), and suppose that still $h^1\left(\Sigma, \mathcal{J}_{X_{i-1}/\Sigma}(D - \sum_{i=1}^{i-1} \Delta_i)\right) > 0$.

We choose $X_i^0 \subseteq X_{i-1}$ minimal such that $h^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) > 0$.

Step 1: $h^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) = 1$, i. e. (b) is fulfilled.

Suppose it was strictly larger than one. By (0) respectively (f), and the Kawamata–Viehweg Vanishing Theorem we have $h^1\left(\Sigma, \mathcal{O}_\Sigma(D - \sum_{k=1}^{i-1} \Delta_k)\right) = 0$.

Thus X_i^0 cannot be empty, that is $\deg(X_i^0) \geq 1$ and we may choose a subscheme $Y \subset X_i^0$ of degree $\deg(Y) = \deg(X_i^0) - 1$. The inclusion $\mathcal{J}_{X_i^0} \hookrightarrow \mathcal{J}_Y$ implies $h^0\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) \leq h^0\left(\Sigma, \mathcal{J}_{Y/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right)$ and the structure sequences of Y and X_i^0 thus lead to

$$\begin{aligned} & h^1\left(\Sigma, \mathcal{J}_{Y/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) \\ &= h^0\left(\Sigma, \mathcal{J}_{Y/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) - h^0\left(\Sigma, \mathcal{O}_\Sigma(D - \sum_{k=1}^{i-1} \Delta_k)\right) + \deg(Y) \\ &\geq h^0\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) - h^0\left(\Sigma, \mathcal{O}_\Sigma(D - \sum_{k=1}^{i-1} \Delta_k)\right) + \deg(X_i^0) - 1 \\ &= h^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) - 1 > 0 \end{aligned}$$

in contradiction to the minimality of X_i^0 .

Step 2: $\deg(X_i^0) \leq \deg(X_0) - \sum_{k=1}^{i-1} \deg(X_{k-1} \cap \Delta_k)$.

The case $i = 1$ follows from the fact that $X_1^0 \subseteq X_0$, and for $i > 1$ the inclusion $X_i^0 \subseteq X_{i-1} = X_{i-2} : \Delta_{i-1}$ implies

$$\deg(X_i^0) \leq \deg(X_{i-2} : \Delta_{i-1}) = \deg(X_{i-2}) - \deg(X_{i-2} \cap \Delta_{i-1}).$$

It thus suffices to show, that

$$\deg(X_{i-2}) - \deg(X_{i-2} \cap \Delta_{i-1}) = \deg(X_0) - \sum_{k=1}^{i-1} \deg(X_{k-1} \cap \Delta_k).$$

If $i = 2$, there is nothing to show. Otherwise $X_{i-2} = X_{i-3} : \Delta_{i-2}$ implies

$$\begin{aligned} & \deg(X_{i-2}) - \deg(X_{i-2} \cap \Delta_{i-1}) \\ &= \deg(X_{i-3} : \Delta_{i-2}) - \deg(X_{i-2} \cap \Delta_{i-1}) \\ &= \deg(X_{i-3}) - \deg(X_{i-3} \cap \Delta_{i-2}) - \deg(X_{i-2} \cap \Delta_{i-1}) \end{aligned}$$

and we are done by induction.

Step 3: There exists a “suitable” locally free rank-two vector bundle E_i .

By the Grothendieck-Serre duality (cf. [Har77] III.7.6) we have $0 \neq H^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) \cong \text{Ext}^1\left(\mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k), \mathcal{O}_\Sigma(K_\Sigma)\right)$, and thus, since $\mathcal{O}_\Sigma(K_\Sigma)$ is locally free, (cf. [Har77] III.6.7)

$$\text{Ext}^1\left(\mathcal{J}_{X_i^0/\Sigma}(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k), \mathcal{O}_\Sigma\right) \neq 0.$$

That is, there is an extension (cf. [Har77] Ex. III.6.1)

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow E_i \rightarrow \mathcal{J}_{X_i^0/\Sigma} \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k \right) \rightarrow 0. \quad (3.2)$$

The minimality of X_i^0 implies that E_i is locally free (cf. [Laz97] Proposition 3.9) and hence that X_i^0 is a locally complete intersection (cf. [Laz97] p. 175). Moreover, we have (cf. [Laz97] Exercise 4.3)

$$c_1(E_i) = D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k \quad \text{and} \quad c_2(E_i) = \deg(X_i^0). \quad (3.3)$$

Step 4: E_i is Bogomolov unstable.

According to the Theorem of Bogomolov we only have to show $c_1(E_i)^2 > 4c_2(E_i)$ (cf. [Bog79] or [Laz97] Theorem 4.2). In order to show this we note that

$$\begin{aligned} & -2 \sum_{k=1}^{i-1} \Delta_k \cdot (D - K_\Sigma - \sum_{j=1}^k \Delta_j) \\ &= -2 \left(\sum_{k=1}^{i-1} \Delta_k \right) \cdot (D - K_\Sigma) + 2 \sum_{k=1}^{i-1} \sum_{j=1}^k \Delta_k \cdot \Delta_j \\ &= -2 \left(\sum_{k=1}^{i-1} \Delta_k \right) \cdot (D - K_\Sigma) + \sum_{k=1}^{i-1} \Delta_k^2 + \left(\sum_{k=1}^{i-1} \Delta_k \right)^2. \end{aligned} \quad (3.4)$$

Since $(4\beta - 1) \cdot (D - K_\Sigma)^2 \leq 0$ by (3) and since $\Delta_k^2 \geq 0$ by (*) we deduce:

$$\begin{aligned} 4c_2(E_i) &= 4 \deg(X_i^0) \leq_{\text{Step 2}} 4 \deg(X_0) - 4 \sum_{k=1}^{i-1} \deg(X_{k-1} \cap \Delta_k) \\ &<_{(3)/(c)} 4\beta(D - K_\Sigma)^2 - 2 \sum_{k=1}^{i-1} \Delta_k \cdot (D - K_\Sigma - \sum_{j=1}^k \Delta_j) - 2 \sum_{k=1}^{i-1} \Delta_k^2 \\ &=_{(3.4)} 4\beta(D - K_\Sigma)^2 - 2 \left(\sum_{k=1}^{i-1} \Delta_k \right) \cdot (D - K_\Sigma) + \left(\sum_{k=1}^{i-1} \Delta_k \right)^2 - \sum_{k=1}^{i-1} \Delta_k^2 \\ &= \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k \right)^2 + (4\beta - 1) \cdot (D - K_\Sigma)^2 - \sum_{k=1}^{i-1} \Delta_k^2 \\ &\leq \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k \right)^2 = c_1(E_i)^2. \end{aligned}$$

Step 5: Find Δ_i .

Since E_i is Bogomolov unstable there exists a zero-dimensional scheme $Z_i \subset \Sigma$ and a divisor $\Delta_i^0 \in \text{Div}(\Sigma)$ such that

$$0 \rightarrow \mathcal{O}_\Sigma(\Delta_i^0) \rightarrow E_i \rightarrow \mathcal{J}_{Z_i/\Sigma} \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - \Delta_i^0 \right) \rightarrow 0 \quad (3.5)$$

is exact (cf. [Laz97] Theorem 4.2) and that

- (d') $(2\Delta_i^0 - D + K_\Sigma + \sum_{k=1}^{i-1} \Delta_k)^2 \geq c_1(E_i)^2 - 4 \cdot c_2(E_i) > 0$, and
- (e') $(2\Delta_i^0 - D + K_\Sigma + \sum_{k=1}^{i-1} \Delta_k) \cdot H > 0$ for all $H \in \text{Div}(\Sigma)$ ample.

We note that (e') implies $h^0\left(\Sigma, \mathcal{O}_\Sigma(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - 2\Delta_i^0)\right) = 0$ and the inclusion $\mathcal{J}_{Z_i/\Sigma} \hookrightarrow \mathcal{O}_\Sigma$ thus gives

$$h^0\left(\Sigma, \mathcal{J}_{Z_i/\Sigma}(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - 2\Delta_i^0)\right) = 0. \quad (3.6)$$

Tensoring (3.5) with $\mathcal{O}_\Sigma(-\Delta_i^0)$ leads to the following exact sequence

$$0 \rightarrow \mathcal{O}_\Sigma \rightarrow E_i(-\Delta_i^0) \rightarrow \mathcal{J}_{Z_i/\Sigma} \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - 2\Delta_i^0 \right) \rightarrow 0, \quad (3.7)$$

and we deduce with (3.6) $h^0(\Sigma, E_i(-\Delta_i^0)) = h^0(\Sigma, \mathcal{O}_\Sigma) = 1$.

Now tensoring (3.2) with $\mathcal{O}_\Sigma(-\Delta_i^0)$ leads to

$$0 \rightarrow \mathcal{O}_\Sigma(-\Delta_i^0) \rightarrow E_i(-\Delta_i^0) \rightarrow \mathcal{J}_{X_i^0/\Sigma} \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - \Delta_i^0 \right) \rightarrow 0. \quad (3.8)$$

By (e'), and (0) respectively (f)

$$-\Delta_i^0.H < -\frac{1}{2}(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k).H \leq 0$$

for an ample divisor H , hence $-\Delta_i^0$ cannot be effective, that is $H^0(\Sigma, -\Delta_i^0) = 0$. But the long exact cohomology sequence of (3.8) then implies

$$0 \neq H^0(\Sigma, E_i(-\Delta_i^0)) \hookrightarrow H^0\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma} \left(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - \Delta_i^0 \right)\right).$$

In particular the linear system $|\mathcal{J}_{X_i^0/\Sigma}(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - \Delta_i^0)|_1$ is non-empty, and we may choose

$$\Delta_i \in \left| \mathcal{J}_{X_i^0/\Sigma}(D - K_\Sigma - \sum_{k=1}^{i-1} \Delta_k - \Delta_i^0) \right|_1.$$

Step 6: Δ_i satisfies (d)-(f).

We note that by the choice of Δ_i we have the following linear equivalences

$$\Delta_i^0 \sim_l D - K_\Sigma - \sum_{k=1}^i \Delta_k \quad (3.9)$$

$$\Delta_i^0 - \Delta_i \sim_l 2\Delta_i^0 - D + K_\Sigma + \sum_{k=1}^{i-1} \Delta_k \sim_l D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i. \quad (3.10)$$

Thus (d) and (e) is a reformulation of (d') and (e').

Moreover, $(\Delta_i^0 - \Delta_i).H > 0$ for any ample H , implies $(\Delta_i^0 - \Delta_i).H \geq 0$ for any H in the closure of the ample cone, in particular

$$(\Delta_i^0 - \Delta_i).H \geq 0 \quad \text{for all } H \text{ nef.} \quad (3.11)$$

But then

$$\Delta_i^0.H \geq \Delta_i.H \geq 0 \quad \text{for all } H \text{ nef,} \quad (3.12)$$

since Δ_i is effective. And finally, since by assumption (*) any effective divisor is nef, we deduce that $\Delta_i^0.C \geq 0$ for any curve C , that is, Δ_i^0 is nef. In view of (3.9) for (f) it remains to show that $(\Delta_i^0)^2 > 0$. Taking once more into account that Δ_i is nef by (*) we have by (d'), (3.10), (3.11) and (3.12)

$$(\Delta_i^0)^2 = (\Delta_i^0 - \Delta_i)^2 + (\Delta_i^0 - \Delta_i).\Delta_i + \Delta_i^0.\Delta_i > 0.$$

Step 7: Δ_i satisfies (c).

We would like to apply the Theorem of Bézout to C_0 and Δ_i . Thus suppose that the irreducible curve C_0 is a component of Δ_i and let H be any ample divisor. Applying (d) and the fact that $D + K_\Sigma$ is nef by (0), we derive the contradiction

$$0 \leq (\Delta_i - C_0).H < -\frac{1}{2} \cdot \left(D + K_\Sigma + \sum_{k=1}^{i-1} \Delta_k \right).H \leq -\frac{1}{2} \cdot (D + K_\Sigma).H \leq 0.$$

Since $X_{i-1} \subseteq X_0 \subset C_0$ the Theorem of Bézout therefore implies

$$D.\Delta_i = C_0.\Delta_i \geq \deg(X_{i-1} \cap \Delta_i).$$

By definition $X_i^0 \subseteq X_{i-1}$ and $X_i^0 \subset \Delta_i$, thus

$$\deg(X_{i-1} \cap \Delta_i) \geq \deg(X_i^0).$$

By assumption (*) the curve Δ_i is nef and thus (3.12) gives

$$(D - K_\Sigma - \sum_{k=1}^i \Delta_k).\Delta_i = \Delta_i^0.\Delta_i \geq \Delta_i^2 \geq 0.$$

Finally from (d') and by (3.3) it follows that

$$(\Delta_i^0 - \Delta_i)^2 \geq c_1(E_i)^2 - 4 \cdot c_2(E_i) = (\Delta_i^0 + \Delta_i)^2 - 4 \cdot \deg(X_i^0),$$

and thus

$$\deg(X_i^0) \geq \Delta_i^0.\Delta_i.$$

Step 8: After a finite number m of steps $h^1\left(\Sigma, \mathcal{J}_{X_m/\Sigma}(D - \sum_{i=1}^m \Delta_i)\right) = 0$, i. e. (a) is fulfilled.

As we have mentioned in Step 1 $\deg(X_i^0) > 0$. This ensures that

$$\deg(X_i) = \deg(X_{i-1}) - \deg(X_{i-1} \cap \Delta_i) \leq \deg(X_{i-1}) - \deg(X_i^0) < \deg(X_{i-1}),$$

i. e. the degree of X_i is strictly diminished each time. Thus the procedure must stop after a finite number m of steps, which is equivalent to the fact that $h^1\left(\Sigma, \mathcal{J}_{X_m/\Sigma}(D - \sum_{i=1}^m \Delta_i)\right) = 0$.

Step 9: It remains to show (3.1).

By assumption (*) the curves Δ_i are nef, in particular $\Delta_i.\Delta_j \geq 0$ for all i, j . Thus (c) implies

$$\begin{aligned} \sum_{i=1}^m \deg(X_i^0) &\geq \sum_{i=1}^m (D - K_\Sigma - \sum_{k=1}^i \Delta_k).\Delta_i \\ &= (D - K_\Sigma).\sum_{i=1}^m \Delta_i - \frac{1}{2} \left(\left(\sum_{i=1}^m \Delta_i \right)^2 + \sum_{i=1}^m \Delta_i^2 \right) \\ &\geq (D - K_\Sigma).\sum_{i=1}^m \Delta_i - \left(\sum_{i=1}^m \Delta_i \right)^2. \end{aligned}$$

But then, taking condition (3) into account,

$$\begin{aligned} 0 \leq \frac{1}{4}(D - K_\Sigma)^2 - \deg(X_0) &\leq \frac{1}{4}(D - K_\Sigma)^2 - \sum_{i=1}^m \deg(X_i^0) \\ &\leq \frac{1}{4}(D - K_\Sigma)^2 - (D - K_\Sigma).\sum_{i=1}^m \Delta_i + \left(\sum_{i=1}^m \Delta_i \right)^2 \\ &= \left(\frac{1}{2}(D - K_\Sigma) - \sum_{i=1}^m \Delta_i \right)^2. \end{aligned}$$

□

It is our overall aim to compare the dimension of a cohomology group of the form $H^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D))$ with some invariants of the X_i^0 and Δ_i . The following lemma will be vital for the necessary estimations.

3.2 Lemma

Let $D \in \text{Div}(\Sigma)$ and let $X_0 \subset \Sigma$ be a zero-dimensional scheme such that there exist curves $\Delta_1, \dots, \Delta_m \subset \Sigma$ and zero-dimensional schemes $X_i^0 \subseteq X_{i-1}$ for $i = 1, \dots, m$, where $X_i = X_{i-1} : \Delta_i$ for $i = 1, \dots, m$, such that (a)-(f) in Lemma 3.1 are fulfilled.

Then:

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) &\leq \sum_{i=1}^m h^1(\Delta_i, \mathcal{J}_{X_{i-1} \cap \Delta_i / \Delta_i}(D - \sum_{k=1}^{i-1} \Delta_k)) \\ &\leq \sum_{i=1}^m \left(1 + \deg(X_{i-1} \cap \Delta_i) - \deg(X_i^0)\right) \\ &\leq \sum_{i=1}^m \left(\Delta_i \cdot (K_\Sigma + \sum_{k=1}^i \Delta_k) + 1\right). \end{aligned}$$

Proof: Throughout the proof we use the following notation

$$\mathcal{G}_i = \mathcal{J}_{X_{i-1} \cap \Delta_i / \Delta_i}(D - \sum_{k=1}^{i-1} \Delta_k) \quad \text{and} \quad \mathcal{G}_i^0 = \mathcal{J}_{X_i^0 / \Delta_i}(D - \sum_{k=1}^{i-1} \Delta_k)$$

for $i = 1, \dots, m$, and

$$\mathcal{F}_i = \mathcal{J}_{X_i / \Sigma}(D - \sum_{k=1}^i \Delta_k),$$

for $i = 0, \dots, m$.

Since $X_{i+1} = X_i : \Delta_{i+1}$ we have the following short exact sequence

$$0 \longrightarrow \mathcal{F}_{i+1} \xrightarrow{\cdot \Delta_{i+1}} \mathcal{F}_i \longrightarrow \mathcal{G}_{i+1} \longrightarrow 0 \quad (3.13)$$

for $i = 0, \dots, m-1$ and the corresponding long exact cohomology sequence

$$\begin{aligned} 0 \longrightarrow H^0(\Sigma, \mathcal{F}_{i+1}) \longrightarrow H^0(\Sigma, \mathcal{F}_i) \longrightarrow H^0(\Sigma, \mathcal{G}_{i+1}) \longrightarrow H^1(\Sigma, \mathcal{F}_{i+1}) \\ \downarrow \\ 0 = H^2(\Sigma, \mathcal{G}_{i+1}) \longleftarrow H^2(\Sigma, \mathcal{F}_i) \longleftarrow H^2(\Sigma, \mathcal{F}_{i+1}) \longleftarrow H^1(\Sigma, \mathcal{G}_{i+1}) \longleftarrow H^1(\Sigma, \mathcal{F}_i) \end{aligned} \quad (3.14)$$

Step 1: $h^1(\Sigma, \mathcal{F}_i) \leq \sum_{j=i+1}^m h^1(\Sigma, \mathcal{G}_j)$ for $i = 0, \dots, m-1$.

We prove the claim by descending induction on i . From (3.14) we deduce

$$0 = H^1(\Sigma, \mathcal{F}_m) \longrightarrow H^1(\Sigma, \mathcal{F}_{m-1}) \longrightarrow H^1(\Sigma, \mathcal{G}_m),$$

which implies $h^1(\Sigma, \mathcal{F}_{m-1}) \leq h^1(\Sigma, \mathcal{G}_m)$ and thus proves the case $i = m-1$.

We may therefore assume that $1 \leq i \leq m-2$. Once more from (3.14) we deduce

$$a = h^0(\Sigma, \mathcal{F}_{i+1}) - h^0(\Sigma, \mathcal{F}_i) + h^0(\Sigma, \mathcal{G}_{i+1}) \geq 0,$$

and

$$b = h^2(\Sigma, \mathcal{F}_{i+1}) - h^2(\Sigma, \mathcal{F}_i) \geq 0,$$

and finally

$$\begin{aligned}
h^1(\Sigma, \mathcal{F}_i) &= h^1(\Sigma, \mathcal{G}_{i+1}) + h^1(\Sigma, \mathcal{F}_{i+1}) - a - b \\
&\leq h^1(\Sigma, \mathcal{G}_{i+1}) + h^1(\Sigma, \mathcal{F}_{i+1}) \\
&\leq_{\text{Ind.}} h^1(\Sigma, \mathcal{G}_{i+1}) + \sum_{j=i+2}^m h^1(\Sigma, \mathcal{G}_j) \\
&= \sum_{j=i+1}^m h^1(\Sigma, \mathcal{G}_j).
\end{aligned}$$

Step 2: $h^1(\Delta_i, \mathcal{G}_i) = h^0(\Delta_i, \mathcal{G}_i) - \chi\left(\mathcal{O}_{\Delta_i}(D - \sum_{k=1}^{i-1} \Delta_k)\right) + \deg(X_{i-1} \cap \Delta_i)$.

We consider the exact sequence

$$0 \longrightarrow \mathcal{G}_i \longrightarrow \mathcal{O}_{\Delta_i}\left(D - \sum_{k=1}^{i-1} \Delta_k\right) \longrightarrow \mathcal{O}_{X_{i-1} \cap \Delta_i / \Delta_i}\left(D - \sum_{k=1}^{i-1} \Delta_k\right) \longrightarrow 0.$$

The result then follows from the long exact cohomology sequence.

Step 3: $h^0(\Delta_i, \mathcal{G}_i^0) - \chi\left(\mathcal{O}_{\Delta_i}(D - \sum_{k=1}^{i-1} \Delta_k)\right) = h^1(\Delta_i, \mathcal{G}_i^0) - \deg(X_i^0)$.

This follows analogously, replacing X_{i-1} by X_i^0 , since $X_i^0 = X_i^0 \cap \Delta_i$.

Step 4: $h^1(\Delta_i, \mathcal{G}_i^0) \leq h^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) = 1$.

Note that $X_i^0 : \Delta_i = \emptyset$, and hence $\mathcal{J}_{X_i^0/\Sigma} = \mathcal{O}_{\Sigma}$. We thus have the following short exact sequence

$$0 \longrightarrow \mathcal{O}_{\Sigma}\left(D - \sum_{k=1}^i \Delta_k\right) \xrightarrow{\cdot \Delta_i} \mathcal{J}_{X_i^0/\Sigma}\left(D - \sum_{k=1}^{i-1} \Delta_k\right) \longrightarrow \mathcal{G}_i^0 \longrightarrow 0. \quad (3.15)$$

By assumption (f) the divisor $D - K_{\Sigma} - \sum_{k=1}^i \Delta_k$ is big and nef and hence

$$0 = h^0\left(\Sigma, \mathcal{O}_{\Sigma}(-D + K_{\Sigma} + \sum_{k=1}^i \Delta_k)\right) = h^2\left(\Sigma, \mathcal{O}_{\Sigma}(D - \sum_{k=1}^i \Delta_k)\right).$$

Thus the long exact cohomology sequence of (3.15) gives

$$H^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right) \longrightarrow H^1(\Delta_i, \mathcal{G}_i^0) \longrightarrow 0,$$

and

$$h^1(\Delta_i, \mathcal{G}_i^0) \leq h^1\left(\Sigma, \mathcal{J}_{X_i^0/\Sigma}(D - \sum_{k=1}^{i-1} \Delta_k)\right).$$

However, by assumption (b) the latter is just one.

Step 5: $h^1(\Delta_i, \mathcal{G}_i) \leq 1 + \deg(X_{i-1} \cap \Delta_i) - \deg(X_i^0)$.

We note that $\mathcal{G}_i \hookrightarrow \mathcal{G}_i^0$, and thus

$$h^0(\Delta_i, \mathcal{G}_i) \leq h^0(\Delta_i, \mathcal{G}_i^0).$$

But then

$$\begin{aligned}
h^1(\Delta_i, \mathcal{G}_i) &\leq_{\text{Step 2/3}} h^1(\Delta_i, \mathcal{G}_i^0) - \deg(X_i^0) + \deg(X_{i-1} \cap \Delta_i) \\
&\leq_{\text{Step 4}} 1 - \deg(X_i^0) + \deg(X_{i-1} \cap \Delta_i).
\end{aligned}$$

Step 6: Finish the proof.

Taking into account, that $h^1(\Sigma, \mathcal{G}_i) = h^1(\Delta_i, \mathcal{G}_i)$, since \mathcal{G}_i is concentrated on Δ_i (cf. [Har77] III.2.10), the first inequality follows from Step 1, while the second inequality is a consequence of Step 5 and the last inequality follows from assumption (c). \square

In the Lemmata 3.3, 3.4 and 3.6 we consider special classes of surfaces which allow us to do the necessary estimations in order to finally derive

$$\sum_{i=1}^m (\#X_i^0 - \dim |\Delta_i|_l) > h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)).$$

We first consider surfaces with Picard number one.

3.3 Lemma

Let Σ be a surface such that³

- (i) $\text{NS}(\Sigma) = L \cdot \mathbb{Z}$,
- (ii) $h^1(\Sigma, C) = 0$, whenever C is effective.

Let $D \in \text{Div}(\Sigma)$ and $X_0 \subset \Sigma$ a zero-dimensional scheme satisfying (0)–(3) from Lemma 3.1 and

$$(4) \quad \sum_{z \in \Sigma} (\deg(X_{0,z}))^2 < \gamma \cdot (D - K_\Sigma)^2, \quad \text{where } \gamma = \frac{(1 + \sqrt{1 - 4\beta})^2 \cdot L^2}{4 \cdot \chi(\mathcal{O}_\Sigma) + \max\{0, 2 \cdot K_\Sigma \cdot L\} + 6 \cdot L^2}.$$

Then, using the notation of Lemma 3.1 and setting $X_S = \bigcup_{i=1}^m X_i^0$,

$$h^1(\Sigma, \mathcal{J}_{X_0/\Sigma}(D)) + \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1) < \#X_S.$$

Proof: We note that by Lemma E.1 we may assume that L is ample. We fix the following notation:

$$D \sim_a d \cdot L, \quad K_\Sigma \sim_a \kappa \cdot L, \quad \Delta_i \sim_a \delta_i \cdot L, \quad \text{and } l = \sqrt{L^2} > 0.$$

Furthermore, we set

$$\alpha = \frac{4 \cdot \chi(\mathcal{O}_\Sigma) + \max\{0, 2 \cdot K_\Sigma \cdot L\} + 6 \cdot L^2}{4 \cdot L^2} = \begin{cases} \frac{\chi(\mathcal{O}_\Sigma)}{L^2} + \frac{\kappa + 3}{2}, & \text{if } \kappa \geq 0, \\ \frac{\chi(\mathcal{O}_\Sigma)}{L^2} + \frac{3}{2}, & \text{if } \kappa < 0, \end{cases}$$

$$\text{and thus } \gamma = \frac{(1 + \sqrt{1 - 4\beta})^2}{4\alpha}.$$

Step 1: Σ satisfies the assumption (*) of Lemma 3.1.

If $c \cdot L \sim_a C \subset \Sigma$ is effective, then in particular $c = \frac{1}{l^2} \cdot C \cdot L > 0$, and thus C is ample, in particular nef. Hence (*) in Lemma 3.1 is fulfilled.

Step 2: $\sum_{i=1}^m \delta_i \cdot l \leq \frac{(d - \kappa) \cdot l}{2} - \sqrt{\frac{(d - \kappa)^2 \cdot l^2}{4} - \deg(X_S)}$, by (3.1).

³By Lemma E.1 we may assume w. l. o. g. that L is ample. – Remember that $\#X_S$ is the number of points in the support of X_S .

Step 3: $h^1(\Sigma, \mathcal{J}_{X_0}(D)) \leq (\kappa \cdot \sum_{i=1}^m \delta_i) \cdot l^2 + \frac{1}{2} \left(\left(\sum_{i=1}^m \delta_i \right)^2 + \sum_{i=1}^m \delta_i^2 \right) \cdot l^2 + m.$

By Lemma 3.2 we know:

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_0}(D)) &\leq \sum_{i=1}^m \left(\Delta_i \cdot (K_\Sigma + \sum_{k=1}^i \Delta_k) + 1 \right) \\ &= \sum_{i=1}^m \delta_i \cdot (\kappa + \sum_{k=1}^i \delta_k) \cdot l^2 + m \\ &= (\kappa \cdot \sum_{i=1}^m \delta_i) \cdot l^2 + \left(\sum_{i=1}^m \sum_{k=1}^i \delta_i \cdot \delta_k \right) \cdot l^2 + m \\ &= (\kappa \cdot \sum_{i=1}^m \delta_i) \cdot l^2 + \frac{1}{2} \left(\left(\sum_{i=1}^m \delta_i \right)^2 + \sum_{i=1}^m \delta_i^2 \right) \cdot l^2 + m. \end{aligned}$$

Step 4: $\sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) \leq m \cdot (\chi(\mathcal{O}_\Sigma) - 1) + \frac{l^2}{2} \cdot \sum_{i=1}^m \delta_i^2 - \frac{\kappa \cdot l^2}{2} \cdot \sum_{i=1}^m \delta_i.$

Since Δ_i is effective by (ii), $h^1(\Sigma, \Delta_i) = 0$. Hence by Riemann-Roch

$$h^0(\Sigma, \Delta_i) \leq \chi(\mathcal{O}_\Sigma(\Delta_i)) = \frac{\Delta_i^2 - K_\Sigma \cdot \Delta_i}{2} + \chi(\mathcal{O}_\Sigma).$$

This implies

$$\begin{aligned} \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) &\leq -m + m \cdot \chi(\mathcal{O}_\Sigma) + \frac{1}{2} \sum_{i=1}^m (\Delta_i^2 - K_\Sigma \cdot \Delta_i) \\ &= m \cdot (\chi(\mathcal{O}_\Sigma) - 1) + \frac{l^2}{2} \cdot \sum_{i=1}^m \delta_i^2 - \frac{\kappa \cdot l^2}{2} \cdot \sum_{i=1}^m \delta_i. \end{aligned}$$

Step 5: Finish the proof.

In the following consideration we use the facts that

$$m \leq \left(\sum_{i=1}^m \delta_i \right)^2, \quad \sum_{i=1}^m \delta_i \leq \left(\sum_{i=1}^m \delta_i \right)^2, \quad \sum_{i=1}^m \delta_i^2 \leq \left(\sum_{i=1}^m \delta_i \right)^2, \quad (3.16)$$

and that

$$\deg(X_S) \leq \deg(X_0) \leq \beta \cdot (d - \kappa)^2 \cdot l^2. \quad (3.17)$$

We thus get:

$$\begin{aligned} &h^1(\Sigma, \mathcal{J}_{X_0}(D)) + \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) \\ &\leq_{\text{Step 3/4}} m \cdot \chi(\mathcal{O}_\Sigma) + l^2 \cdot \sum_{i=1}^m \delta_i^2 + \frac{\kappa \cdot l^2}{2} \cdot \sum_{i=1}^m \delta_i + \frac{l^2}{2} \cdot \left(\sum_{i=1}^m \delta_i \right)^2 \\ &\leq_{(3.16)} \alpha \cdot \left(l \cdot \sum_{i=1}^m \delta_i \right)^2 \leq_{\text{Step 2}} \alpha \cdot \left(\frac{(d-\kappa) \cdot l}{2} - \sqrt{\frac{(d-\kappa)^2 \cdot l^2}{4} - \deg(X_S)} \right)^2 \\ &\leq \alpha \cdot \left(\frac{\frac{(d-\kappa)^2 \cdot l^2}{4} - \left(\frac{(d-\kappa)^2 \cdot l^2}{4} - \deg(X_S) \right)}{\frac{(d-\kappa) \cdot l}{2} + \sqrt{\frac{(d-\kappa)^2 \cdot l^2}{4} - \deg(X_S)}} \right)^2 = \alpha \cdot \left(\frac{2 \cdot \deg(X_S)}{(d-\kappa) \cdot l + \sqrt{(d-\kappa)^2 \cdot l^2 - 4 \cdot \deg(X_S)}} \right)^2 \\ &\leq_{(3.17)} \frac{4\alpha}{(1 + \sqrt{1 - 4\beta})^2 \cdot (d-\kappa)^2 \cdot l^2} \cdot (\deg(X_S))^2 = \frac{1}{\gamma \cdot (D - K_\Sigma)^2} \cdot \left(\sum_{z \in \Sigma} \deg(X_{S,z}) \right)^2 \\ &\leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{S,z})^2 \leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{0,z})^2 <_{(4)} \#X_S. \end{aligned}$$

□

The second class of surfaces which we consider, are products of curves. We use the notation of Section G.b.

3.4 Lemma

Let C_1 and C_2 be two smooth projective curves of genera g_1 and g_2 respectively with $g_1 \geq g_2 \geq 0$, such that for $\Sigma = C_1 \times C_2$ the Néron–Severi group is $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z}$, and let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_1 + bC_2$ with $a > \max\{2g_2 - 2, 2 - 2g_2\}$ and $b > \max\{2g_1 - 2, 2 - 2g_1\}$. Suppose moreover that $X_0 \subset \Sigma$ is a zero-dimensional scheme satisfying (1)–(3) from Lemma 3.1 and

$$(4) \quad \sum_{z \in \Sigma} (\deg(X_{0,z}))^2 < \gamma \cdot (D - K_\Sigma)^2,$$

where γ may be taken from the following table with $\alpha = \frac{a-2g_2+2}{b-2g_1+2} > 0$.

g_1	g_2	γ
0	0	$\frac{1}{24}$
1	0	$\frac{1}{\max\{32, 2\alpha\}}$
≥ 2	0	$\frac{1}{\max\{24+16g_1, 4g_1, \alpha\}}$
1	1	$\frac{1}{\max\{32, 2\alpha, \frac{2}{\alpha}\}}$
≥ 2	≥ 1	$\frac{1}{\max\{24+16g_1+16g_2, 4g_1, \frac{4g_2}{\alpha}\}}$

Then, using the notation of Lemma 3.1 and setting⁴ $X_S = \bigcup_{i=1}^m X_i^0$,

$$h^1(\Sigma, \mathcal{I}_{X_0}(D)) + \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) < \#X_S.$$

Proof: Then $K_\Sigma \sim_a (2g_2 - 2) \cdot C_1 + (2g_1 - 2) \cdot C_2$ and we fix the notation:

$$\Delta_i \sim_a a_i C_1 + b_i C_2.$$

Step 1: Σ satisfies the assumption (*) of Lemma 3.1 by Lemma G.7. Moreover, due to the assumptions on a and b we know that $D - K_\Sigma$ is ample and $D + K_\Sigma$ is nef, i. e. (0) in Lemma 3.1 is fulfilled as well.

Step 2a: $\left(\frac{a-2g_2+2}{4}\right) \cdot \sum_{i=1}^m b_i + \left(\frac{b-2g_1+2}{4}\right) \cdot \sum_{i=1}^m a_i \leq \deg(X_S)$.

Let us notice first that the strict inequality “ $<$ ” in Lemma 3.1 (e) for ample divisors H comes down to “ \leq ” for nef divisors H . We may apply this for $H = C_1$ and $H = C_2$ and deduce the following inequalities:

$$0 \leq \left(D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i \right) \cdot C_1 = b - 2g_1 + 2 - \sum_{k=1}^i b_k - b_i, \quad (3.18)$$

⁴Remember that $\#X_S$ is the number of points in the support of X_S .

and

$$0 \leq \left(D - K_{\Sigma} - \sum_{k=1}^i \Delta_k - \Delta_i \right) \cdot C_2 = a - 2g_2 + 2 - \sum_{k=1}^i a_k - a_i. \quad (3.19)$$

For the following consideration we choose $i_0, j_0 \in \{1, \dots, m\}$ such that $a_{i_0} \geq a_i$ for all $i = 1, \dots, m$ and $b_{j_0} \geq b_j$ for all $j = 1, \dots, m$. Then

$$a - 2g_2 + 2 \geq \sum_{k=1}^{i_0} a_k + a_{i_0} \geq 2a_{i_0} \geq 2a_i \quad (3.20)$$

and

$$b - 2g_1 + 2 \geq \sum_{k=1}^{j_0} b_k + b_{j_0} \geq 2b_{j_0} \geq 2b_j \quad (3.21)$$

for all $i, j = 1, \dots, m$, and we finally get

$$\begin{aligned} \deg(X_S) &= \sum_{i=1}^m \deg(X_i^0) \stackrel{\text{Lemma 3.1 (c)}}{\geq} \sum_{i=1}^m (D - K_{\Sigma} - \sum_{k=1}^i \Delta_k) \cdot \Delta_i \\ &= (D - K_{\Sigma}) \cdot \sum_{i=1}^m \Delta_i - \sum_{1 \leq k \leq i \leq 1} \Delta_k \cdot \Delta_i \\ &= (D - K_{\Sigma}) \cdot \sum_{i=1}^m \Delta_i - \frac{1}{2} \sum_{i=1}^m \Delta_i^2 - \frac{1}{2} \left(\sum_{i=1}^m \Delta_i \right)^2 \\ &= (a - 2g_2 + 2) \sum_{i=1}^m b_i + (b - 2g_1 + 2) \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b_i - \sum_{i=1}^m a_i \sum_{i=1}^m b_i \\ &\stackrel{(3.18)/(3.19)}{\geq} \frac{a-2g_2+2}{2} \sum_{i=1}^m b_i + \frac{b-2g_1+2}{2} \sum_{i=1}^m a_i + \frac{1}{2} \left(\sum_{i=1}^m a_i + a_m \right) \cdot \sum_{i=1}^m b_i \\ &\quad + \frac{1}{2} \left(\sum_{i=1}^m b_i + b_m \right) \cdot \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b_i - \sum_{i=1}^m a_i \sum_{i=1}^m b_i \\ &= \frac{a-2g_2+2}{2} \sum_{i=1}^m b_i + \frac{b-2g_1+2}{2} \sum_{i=1}^m a_i + \frac{a_m}{2} \sum_{i=1}^m b_i + \frac{b_m}{2} \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b_i \\ &\stackrel{(3.20)/(3.21)}{\geq} \frac{a-2g_2+2}{4} \sum_{i=1}^m b_i + \frac{1}{4} \sum_{i=1}^m 2a_i b_i + \frac{b-2g_1+2}{4} \sum_{i=1}^m a_i + \frac{1}{4} \sum_{i=1}^m 2a_i b_i \\ &\quad + \frac{a_m}{2} \sum_{i=1}^m b_i + \frac{b_m}{2} \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b_i \\ &\geq \frac{a-2g_2+2}{4} \sum_{i=1}^m b_i + \frac{b-2g_1+2}{4} \sum_{i=1}^m a_i. \end{aligned}$$

Step 2b: $\sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i \leq \frac{8}{(D-K_{\Sigma})^2} \cdot (\deg(X_S))^2$.

Using Step 2a we deduce

$$\begin{aligned} (\deg(X_S))^2 &> \left(\frac{b-2g_1+2}{4} \cdot \sum_{i=1}^m a_i + \frac{a-2g_2+2}{4} \cdot \sum_{i=1}^m b_i \right)^2 \\ &\geq \frac{4 \cdot (a-2g_2+2) \cdot (b-2g_1+2)}{16} \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i \\ &= \frac{(D-K_{\Sigma})^2}{8} \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i. \end{aligned}$$

Step 2c: $\sum_{i=1}^m a_i \leq \begin{cases} \frac{2\alpha}{(D-K_{\Sigma})^2} \cdot (\deg(X_S))^2, & \text{if } \sum_{i=1}^m b_i = 0, \\ \frac{8}{(D-K_{\Sigma})^2} \cdot (\deg(X_S))^2, & \text{else.} \end{cases}$

If $\sum_{i=1}^m b_i = 0$, then the same consideration as in Step 2a shows

$$\deg(X_S) \geq (b - 2g_1 + 2) \cdot \sum_{i=1}^m a_i > 0,$$

and thus

$$\frac{(D-K_\Sigma)^2}{2\alpha} \cdot \sum_{i=1}^m a_i \leq (b - 2g_1 + 2)^2 \cdot \left(\sum_{i=1}^m a_i \right)^2 \leq (\deg(X_S))^2.$$

If $\sum_{i=1}^m b_i \neq 0$, then by Step 2b

$$\sum_{i=1}^m a_i \leq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i \leq \frac{8}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2.$$

Step 2d: $\sum_{i=1}^m b_i \leq \begin{cases} \frac{2}{\alpha \cdot (D-K_\Sigma)^2} \cdot (\deg(X_S))^2, & \text{if } \sum_{i=1}^m a_i = 0, \\ \frac{8}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2, & \text{else.} \end{cases}$

This is proved in the same way as Step 2c.

Step 3: $h^1(\Sigma, \mathcal{J}_{X_0}(D)) \leq 2 \sum_{i=1}^m a_i \sum_{i=1}^m b_i + (2g_1 - 2) \sum_{i=1}^m a_i + (2g_2 - 2) \sum_{i=1}^m b_i + m.$

The following sequence of inequalities is due to Lemma 3.2 and the fact that $\Delta_i \cdot \Delta_j \geq 0$ for any $i, j \in \{1, \dots, m\}$:

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_0}(D)) &\leq \sum_{i=1}^m \left(\Delta_i \cdot (K_\Sigma + \sum_{k=1}^i \Delta_k) + 1 \right) \\ &= K_\Sigma \cdot \sum_{i=1}^m \Delta_i + \sum_{1 \leq k \leq i \leq m} \Delta_i \cdot \Delta_k + m \\ &\leq K_\Sigma \cdot \sum_{i=1}^m \Delta_i + \left(\sum_{i=1}^m \Delta_i \right)^2 + m \\ &= (2g_1 - 2) \cdot \sum_{i=1}^m a_i + (2g_2 - 2) \cdot \sum_{i=1}^m b_i + 2 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + m. \end{aligned}$$

Step 4: We find the estimation $\sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) \leq \beta$, where

$$\beta = \begin{cases} \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + \sum_{i=1}^m b_i, & \text{if } g_1 = 1, g_2 = 0, \\ \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i - m, & \text{if } g_1 = 1, g_2 = 1, \exists i_0 : a_{i_0} b_{i_0} > 0, \\ \sum_{i=1}^m a_i + \sum_{i=1}^m b_i - m, & \text{if } g_1 = 1, g_2 = 1, \forall i : a_i b_i = 0, \\ \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + \sum_{i=1}^m a_i + \sum_{i=1}^m b_i, & \text{else.} \end{cases}$$

In general by Corollary G.9 $h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) \leq a_i b_i + a_i + b_i + 1$, while if $g_1 = 1, g_2 = 0$ by Lemma G.5 we have $h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) = a_i b_i + b_i + 1$. It thus only remains to consider the case $g_1 = g_2 = 1$.

Applying Lemma G.15 we get

$$\sum_{i=1}^m h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) = \sum_{a_i, b_i > 0} a_i b_i + \sum_{a_i=0} b_i + \sum_{b_i=0} a_i.$$

If always either a_i or b_i is zero, we are done. Otherwise there exists some $i_0 \in \{1, \dots, m\}$ such that $a_{i_0} \neq 0 \neq b_{i_0}$. Then looking at the right hand side we see

$$\sum_{i=1}^m h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) \leq \sum_{a_i, b_i > 0} a_i b_i + a_{i_0} \cdot \sum_{a_i=0} b_i + b_{i_0} \cdot \sum_{b_i=0} a_i \leq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i.$$

Step 5: Finish the proof.

Using Step 3 and Step 4, and taking $m \leq \sum_{i=1}^m a_i + b_i$ into account, we get $h^1(\Sigma, \mathcal{J}_{X_0}(D)) + \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1) \leq \beta'$, where β' may be chosen as

$$\beta' = \begin{cases} 3 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i, & \text{if } g_1 = 0, g_2 = 0, \\ 3 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + \sum_{i=1}^m a_i, & \text{if } g_1 = 1, g_2 = 0, \\ 3 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + 2g_1 \cdot \sum_{i=1}^m a_i + 2g_2 \cdot \sum_{i=1}^m b_i, & \text{if } g_1 \geq 2, g_2 \geq 0. \end{cases}$$

For the case $g_1 = g_2 = 1$ we take a closer look. We find at once the following upper bounds β'' for $h^1(\Sigma, \mathcal{J}_{X_0}(D)) + \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1)$

$$\beta'' = \begin{cases} 3 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i, & \text{if } \exists i_0 : a_{i_0} b_{i_0} \neq 0, \\ 2 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i + \sum_{i=1}^m a_i + \sum_{i=1}^m b_i, & \text{if } \forall i : a_i b_i = 0. \end{cases}$$

Considering now the cases $\sum_{i=1}^m a_i \neq 0 \neq \sum_{i=1}^m b_i$, $\sum_{i=1}^m a_i = 0$ and $\sum_{i=1}^m b_i = 0$, we can replace these by

$$\beta'' \leq \beta' = \begin{cases} 4 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i, & \text{if } \sum_{i=1}^m a_i \neq 0 \neq \sum_{i=1}^m b_i, \\ \sum_{i=1}^m a_i, & \text{if } \sum_{i=1}^m b_i = 0, \\ \sum_{i=1}^m b_i, & \text{if } \sum_{i=1}^m a_i = 0. \end{cases}$$

Applying now the results of Step 2 in all cases we get

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_0}(D)) + \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1) &\leq \beta' \leq \frac{1}{\gamma \cdot (D - K_\Sigma)^2} \cdot (\deg(X_S))^2 \\ &= \frac{1}{\gamma \cdot (D - K_\Sigma)^2} \cdot \left(\sum_{z \in \Sigma} \deg(X_{S,z}) \right)^2 \leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{S,z})^2 \\ &\leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{0,z})^2 <_{(4)} \#X_S. \end{aligned}$$

□

3.5 Remark

Lemma 3.4, and hence Theorem 2.3 could easily be generalised to other surfaces Σ with irreducible curves $C_1, C_2 \subset \Sigma$ such that

$$\text{NS}(\Sigma) = C_1 \mathbb{Z} \oplus C_2 \mathbb{Z} \text{ with intersection matrix } \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (3.22)$$

once we have an estimation similar to

$$h^0(\Sigma, aC_1 + bC_2) \leq ab + a + b + 1$$

for an effective divisor $aC_1 + bC_2$.

With a number of small modifications we are even able to adapt it in the following lemma in the case of geometrically ruled surfaces with non-positive invariant e although the intersection pairing looks more complicated.

The problem with arbitrary geometrically ruled surfaces is the existence of the section with negative self-intersection, once the invariant $e > 0$, since then the proof of Lemma 3.1 no longer works.

In the following lemma we use the notation of Section G.a.

3.6 Lemma

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with invariant $e \leq 0$ and $g = g(C)$, and let $D \in \text{Div}(\Sigma)$ such that $D \sim_{\alpha} aC_0 + bF$ with $a \geq 2$, $b > 2g - 2 + \frac{ae}{2}$, and if $g = 0$ then $b \geq 2$. Suppose moreover that $X_0 \subset \Sigma$ is a zero-dimensional scheme satisfying (1)–(3) from Lemma 3.1 and

$$(4) \quad \sum_{z \in \Sigma} (\deg(X_{0,z}))^2 < \gamma \cdot (D - K_{\Sigma})^2,$$

where γ may be taken from the following table with $\alpha = \frac{a+2}{b+2-2g-\frac{ae}{2}} > 0$.

g	e	γ
0	0	$\frac{1}{24}$
1	0	$\frac{1}{\max\{24, 2\alpha\}}$
1	-1	$\frac{1}{\max\left\{\min\left\{30 + \frac{16}{\alpha} + 4\alpha, 40 + 9\alpha\right\}, \frac{13}{2}\alpha\right\}}$
≥ 2	0	$\frac{1}{\max\{24 + 16g, 4g\alpha\}}$
≥ 2	< 0	$\frac{1}{\max\left\{\min\left\{24 + 16g - 9e\alpha, 18 + 16g - 9e\alpha - \frac{16}{e\alpha}\right\}, 4g\alpha - 9e\alpha\right\}}$

Then, using the notation of Lemma 3.1 and setting⁵ $X_S = \bigcup_{i=1}^m X_i^0$,

$$h^1(\Sigma, \mathcal{I}_{X_0}(D)) + \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_{\Sigma}(\Delta_i)) - 1 \right) < \#X_S.$$

Proof: Remember that the Néron–Severi group of Σ is generated by a section C_0 of π and a fibre F with intersection pairing given by $\begin{pmatrix} -e & 1 \\ 1 & 0 \end{pmatrix}$. Then $K_{\Sigma} \sim_{\alpha} -2C_0 + (2g - 2 - e) \cdot F$ and we fix the notation:

$$\Delta_i \sim_{\alpha} a_i C_0 + b_i F.$$

Note that by Lemma G.1 we have

$$a_i \geq 0 \quad \text{and} \quad b'_i := b_i - \frac{e}{2} a_i \geq 0.$$

Finally we set $\kappa_1 = a + 2$ and $\kappa_2 = b + 2 - 2g - \frac{ae}{2}$ and get

$$(D - K_{\Sigma})^2 = -e \cdot (a + 2)^2 + 2 \cdot (a + 2) \cdot (b + 2 + e - 2g) = 2 \cdot \kappa_1 \cdot \kappa_2. \quad (3.23)$$

⁵Remember that $\#X_S$ is the number of points in the support of X_S .

Step 1: By Lemma G.1 Σ satisfies the assumption (*) of Lemma 3.1, and by the assumptions on a and b we know that $D - K_\Sigma$ is ample and $D + K_\Sigma$ is nef, that is, (0) in Lemma 3.1 is fulfilled.

Step 2a: $\frac{\kappa_1}{4} \cdot \sum_{i=1}^m b'_i + \frac{\kappa_2}{4} \cdot \sum_{i=1}^m a_i \leq \deg(X_S)$.

Let us notice first that the strict inequality “ $<$ ” in Lemma 3.1 (e) for ample divisors H comes down to “ \leq ” for nef divisors H . We may apply this for $H = C_0 + \frac{\epsilon}{2} \cdot F$ and $H = F$ and deduce the following inequalities:

$$0 \leq \left(D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i \right) \cdot (C_0 + \frac{\epsilon}{2} F) = \kappa_2 - \sum_{k=1}^i b'_k - b'_i, \quad (3.24)$$

and

$$0 \leq \left(D - K_\Sigma - \sum_{k=1}^i \Delta_k - \Delta_i \right) \cdot F = \kappa_1 - \sum_{k=1}^i a_k - a_i. \quad (3.25)$$

For the following consideration we choose $i_0, j_0 \in \{1, \dots, m\}$ such that $a_{i_0} \geq a_i$ for all $i = 1, \dots, m$ and $b'_{j_0} \geq b'_j$ for all $j = 1, \dots, m$. Then

$$\kappa_1 \geq \sum_{k=1}^{i_0} a_k + a_{i_0} \geq 2a_{i_0} \geq 2a_i \quad (3.26)$$

and

$$\kappa_2 \geq \sum_{k=1}^{j_0} b'_k + b'_{j_0} \geq 2b'_{j_0} \geq 2b'_j \quad (3.27)$$

for all $i, j = 1, \dots, m$, and we finally get

$$\begin{aligned} \deg(X_S) &= \sum_{i=1}^m \deg(X_i^0) \stackrel{\geq_{3.1(c)}}{\geq} \sum_{i=1}^m (D - K_\Sigma - \sum_{k=1}^i \Delta_k) \cdot \Delta_i \\ &= (D - K_\Sigma) \cdot \sum_{i=1}^m \Delta_i - \sum_{1 \leq k < i \leq m} \Delta_k \cdot \Delta_i \\ &= (D - K_\Sigma) \cdot \sum_{i=1}^m \Delta_i - \frac{1}{2} \sum_{i=1}^m \Delta_i^2 - \frac{1}{2} \left(\sum_{i=1}^m \Delta_i \right)^2 \\ &= \kappa_1 \cdot \sum_{i=1}^m b'_i + \kappa_2 \cdot \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b'_i - \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i \\ &\stackrel{\geq_{(3.24)/(3.25)}}{\geq} \frac{\kappa_1}{2} \cdot \sum_{i=1}^m b'_i + \frac{\kappa_2}{2} \cdot \sum_{i=1}^m a_i + \frac{1}{2} \cdot \left(\sum_{i=1}^m a_i + a_m \right) \cdot \sum_{i=1}^m b'_i \\ &\quad + \frac{1}{2} \cdot \left(\sum_{i=1}^m b'_i + b'_m \right) \cdot \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b'_i - \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i \\ &= \frac{\kappa_1}{2} \cdot \sum_{i=1}^m b'_i + \frac{\kappa_2}{2} \cdot \sum_{i=1}^m a_i + \frac{a_m}{2} \cdot \sum_{i=1}^m b'_i + \frac{b'_m}{2} \cdot \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b'_i \\ &\stackrel{\geq_{(3.26)/(3.27)}}{\geq} \frac{\kappa_1}{4} \cdot \sum_{i=1}^m b'_i + \frac{1}{4} \sum_{i=1}^m 2a_i b'_i + \frac{\kappa_2}{4} \cdot \sum_{i=1}^m a_i + \frac{1}{4} \sum_{i=1}^m 2a_i b'_i \\ &\quad + \frac{a_m}{2} \cdot \sum_{i=1}^m b'_i + \frac{b'_m}{2} \cdot \sum_{i=1}^m a_i - \sum_{i=1}^m a_i b'_i \\ &\geq \frac{\kappa_1}{4} \cdot \sum_{i=1}^m b'_i + \frac{\kappa_2}{4} \cdot \sum_{i=1}^m a_i. \end{aligned}$$

Step 2b: $\sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i \leq \frac{8}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2$.

Using Step 2a and taking (3.23) into account, we deduce

$$(\deg(X_S))^2 \geq \left(\frac{\kappa_2}{4} \cdot \sum_{i=1}^m a_i + \frac{\kappa_1}{4} \cdot \sum_{i=1}^m b'_i \right)^2 \geq \frac{4 \cdot \kappa_1 \cdot \kappa_2}{16} \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i.$$

Step 2c: $\left(\sum_{i=1}^m a_i \right)^2 \leq \frac{32\alpha}{(D-K_\Sigma)^2} (\deg(X_S))^2$ and $\left(\sum_{i=1}^m b'_i \right)^2 \leq \frac{32}{\alpha \cdot (D-K_\Sigma)^2} (\deg(X_S))^2$.

This follows from the following inequality with the aid of Step 2a and (3.23),

$$\begin{aligned} (\deg(X_S))^2 &\geq \left(\frac{\kappa_2}{4} \cdot \sum_{i=1}^m a_i \right)^2 + \left(\frac{\kappa_1}{4} \cdot \sum_{i=1}^m b'_i \right)^2 \\ &\geq \frac{2 \cdot \kappa_1 \cdot \kappa_2}{32\alpha} \cdot \left(\sum_{i=1}^m a_i \right)^2 + \frac{2 \cdot \kappa_1 \cdot \kappa_2 \cdot \alpha}{32} \cdot \left(\sum_{i=1}^m b'_i \right)^2. \end{aligned}$$

Step 2d: $\sum_{i=1}^m a_i \leq \begin{cases} \frac{2\alpha}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2, & \text{if } \sum_{i=1}^m b'_i = 0, \\ \frac{8}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2, & \text{else.} \end{cases}$

If $\sum_{i=1}^m b'_i = 0$, i. e. $b'_i = 0$ for all $i = 1, \dots, m$, then the same consideration as in Step 2a shows

$$\deg(X_S) \geq \kappa_2 \cdot \sum_{i=1}^m a_i,$$

and thus

$$\frac{(D-K_\Sigma)^2}{2\alpha} \cdot \sum_{i=1}^m a_i \leq \kappa_2^2 \cdot \left(\sum_{i=1}^m a_i \right)^2 \leq (\deg(X_S))^2.$$

If $\sum_{i=1}^m b'_i \neq 0$, then by Step 2b

$$\sum_{i=1}^m a_i \leq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i \leq \frac{8}{(D-K_\Sigma)^2} \cdot (\deg(X_S))^2.$$

Step 3: $h^1(\Sigma, \mathcal{J}_{X_0}(D)) \leq 2 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + (2g-2) \cdot \sum_{i=1}^m a_i - 2 \cdot \sum_{i=1}^m b'_i + m$.

By Lemma 3.2 and since $\Delta_i \cdot \Delta_j \geq 0$ for any $i, j \in \{1, \dots, m\}$ we have:

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_0}(D)) &\leq \sum_{i=1}^m \left(\Delta_i \cdot (K_\Sigma + \sum_{k=1}^i \Delta_k) + 1 \right) \\ &\leq K_\Sigma \cdot \sum_{i=1}^m \Delta_i + \left(\sum_{i=1}^m \Delta_i \right)^2 + m \\ &= (2g-2) \cdot \sum_{i=1}^m a_i - 2 \cdot \sum_{i=1}^m b'_i + 2 \cdot \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + m. \end{aligned}$$

Step 4a: If $e = 0$, we find the estimation

$$\sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) \leq \begin{cases} \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + \sum_{i=1}^m b'_i - m, & \text{if } g = 1, \sum_{i=1}^m b'_i \neq 0, \\ \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + \sum_{i=1}^m b'_i = 0, & \text{if } g = 1, \sum_{i=1}^m b'_i = 0, \\ \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + \sum_{i=1}^m a_i + \sum_{i=1}^m b'_i, & \text{for } g \text{ arbitrary.} \end{cases}$$

We note that in this case $b_i = b'_i$ and that $b'_i = 0$ thus implies $a_i > 0$. By Corollary G.5 we have

$$h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) \leq \begin{cases} a_i b'_i + b'_i, & \text{if } g = 1, b'_i > 0, \\ a_i b'_i + b'_i + 1 = 1, & \text{if } g = 1, b'_i = 0, \\ a_i b'_i + a_i + b'_i + 1, & \text{else.} \end{cases}$$

The results for g arbitrary respectively $g = 1$ and $\sum_{i=1}^m b'_i = 0$ thus follow right away. If, however, some $b'_{i_0} > 0$, then

$$\sum_{i \neq j} a_i b_j \geq b_{i_0} \sum_{i \neq i_0} a_i \geq \#\{b'_i \mid b'_i = 0\}$$

and hence

$$\begin{aligned} h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) &\leq \sum_{i=1}^m a_i b'_i + \sum_{i=1}^m b'_i + \#\{b'_i \mid b'_i = 0\} \\ &= \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + \sum_{i=1}^m b'_i + \#\{b'_i \mid b'_i = 0\} - \sum_{i \neq j} a_i b'_j \leq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b'_i + \sum_{i=1}^m b'_i. \end{aligned}$$

Step 4b: If $e < 0$, we give several upper bounds for $\beta = \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1)$:

$$\beta \leq \begin{cases} \frac{1}{2} \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + \frac{1}{2} \left(\sum_{i=1}^m b'_i \right)^2 + \frac{1}{8} \left(\sum_{i=1}^m a_i \right)^2 + \frac{1}{4} \sum_{i=1}^m a_i + \frac{1}{2} \sum_{i=1}^m b'_i, & \text{if } g = 1, \\ \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + \sum_{i=1}^m a_i + \sum_{i=1}^m b'_i - \frac{9e}{32} \left(\sum_{i=1}^m a_i \right)^2, & \text{for } g \text{ arbitrary.} \\ \frac{1}{4} \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + \sum_{i=1}^m a_i + \sum_{i=1}^m b'_i - \frac{9e}{32} \left(\sum_{i=1}^m a_i \right)^2 - \frac{1}{2e} \left(\sum_{i=1}^m b'_i \right)^2, & g \text{ arbitrary.} \end{cases}$$

If g is arbitrary, the claim follows since by Corollary G.5 we have

$$h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) \leq a_i b'_i + a_i + b'_i + 1 - \frac{9e}{32} \cdot a_i^2$$

and

$$h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) \leq \frac{1}{4} \cdot a_i b'_i + a_i + b'_i + 1 - \frac{9e}{32} \cdot a_i^2 - \frac{1}{2e} \cdot b_i'^2.$$

If $g = 1$, then $e = -1$ and $b' = b + \frac{a}{2}$. We may once more apply Corollary G.5 and see that in any case

$$\begin{aligned} h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) &\leq a_i b_i + b_i + 1 + \frac{a_i(a_i+1)}{2} + \frac{b_i(b_i-1)}{2} \\ &= \frac{1}{2} \cdot a_i b'_i + \frac{1}{2} \cdot b_i'^2 + \frac{1}{8} \cdot a_i^2 + \frac{1}{4} \cdot a_i + \frac{1}{2} \cdot b'_i + 1, \end{aligned}$$

which finishes the case $g = 1$.

Step 5: In this last step we gather the information from the previous investigations and finish the proof considering a bunch of different cases.

Using Step 3 and Step 4 and taking $\sum_{i=1}^m a_i + b'_i \leq m$ into account, we get the following upper bounds for $\beta' = h^1(\Sigma, \mathcal{J}_{X_0}(D)) + \sum_{i=1}^m (h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1)$

$$\beta' \leq \begin{cases} 3 \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + 2g \sum_{i=1}^m a_i, & \text{if } e = 0, \\ 3 \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + 2g \sum_{i=1}^m a_i - \frac{9e}{32} \left(\sum_{i=1}^m a_i \right)^2, & \text{if } e < 0, \\ \frac{9}{4} \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + 2g \sum_{i=1}^m a_i - \frac{9e}{32} \left(\sum_{i=1}^m a_i \right)^2 - \frac{1}{2e} \left(\sum_{i=1}^m b'_i \right)^2, & \text{if } e < 0, \\ 3 \sum_{i=1}^m a_i \sum_{i=1}^m b'_i, & \text{if } e = 0, g = 1, \sum_{i=1}^m b'_i \neq 0, \\ m \leq \sum_{i=1}^m a_i, & \text{if } e = 0, g = 1, \sum_{i=1}^m b'_i = 0, \\ \frac{5}{2} \sum_{i=1}^m a_i \sum_{i=1}^m b'_i + \frac{1}{2} \left(\sum_{i=1}^m b'_i \right)^2 + \frac{1}{8} \left(\sum_{i=1}^m a_i \right)^2 + \frac{5}{4} \sum_{i=1}^m a_i, & \text{if } e < 0, g = 1. \end{cases}$$

Applying now the results of Step 2b-2d we get

$$\frac{\beta' \cdot (D - K_\Sigma)^2}{(\deg(X_S))^2} \leq \begin{cases} 24, & \text{if } g = 0, \\ 24, & \text{if } g = 1, e = 0, \sum_{i=1}^m b'_i \neq 0, \\ 2\alpha, & \text{if } g = 1, e = 0, \sum_{i=1}^m b'_i = 0, \\ \min \left\{ 30 + \frac{16}{\alpha} + 4\alpha, 40 + 9\alpha \right\}, & \text{if } g = 1, e < 0, \sum_{i=1}^m b'_i \neq 0, \\ \frac{13}{2}\alpha, & \text{if } g = 1, e < 0, \sum_{i=1}^m b'_i = 0, \\ 24 + 16g, & \text{if } g \geq 2, e = 0, \sum_{i=1}^m b'_i \neq 0, \\ 4g\alpha, & \text{if } g \geq 2, e = 0, \sum_{i=1}^m b'_i = 0, \\ \min \left\{ 24 + 16g - 9e\alpha, 18 + 16g - 9e\alpha - \frac{16}{e\alpha} \right\}, & \\ 4g\alpha - 9e\alpha, & \text{if } g \geq 2, e < 0, \sum_{i=1}^m b'_i \neq 0, \\ & \text{if } g \geq 2, e < 0, \sum_{i=1}^m b'_i = 0. \end{cases}$$

Hence we have

$$\frac{\beta' \cdot (D - K_\Sigma)^2}{(\deg(X_S))^2} \leq \begin{cases} 24, & \text{if } g = 0, \\ \max\{24, 2\alpha\}, & \text{if } g = 1, e = 0, \\ \max \left\{ \min \left\{ 30 + \frac{16}{\alpha} + 4\alpha, 40 + 9\alpha \right\}, \frac{13}{2}\alpha \right\}, & \text{if } g = 1, e < 0, \\ \max\{24 + 16g, 4g\alpha\} & \text{if } g \geq 2, e = 0, \\ \max \left\{ \min \left\{ 24 + 16g - 9e\alpha, 18 + 16g - 9e\alpha - \frac{16}{e\alpha} \right\}, 4g\alpha - 9e\alpha \right\}, & \\ & \text{if } g \geq 2, e < 0. \end{cases}$$

We thus finally get

$$\begin{aligned} h^1(\Sigma, \mathcal{I}_{X_0}(D)) + \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) &= \beta' \leq \frac{1}{\gamma \cdot (D - K_\Sigma)^2} \cdot (\deg(X_S))^2 \\ &= \frac{1}{\gamma \cdot (D - K_\Sigma)^2} \cdot \left(\sum_{z \in \Sigma} \deg(X_{S,z}) \right)^2 \leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{S,z})^2 \\ &\leq \frac{\#X_S}{\gamma \cdot (D - K_\Sigma)^2} \cdot \sum_{z \in \Sigma} \deg(X_{0,z})^2 \stackrel{(4)}{<} \#X_S. \end{aligned}$$

□

It remains to show, that the inequality which we derived in the above cases cannot hold.

3.7 Lemma

Let $D \in \text{Div}(\Sigma)$, $\mathcal{S}_1, \dots, \mathcal{S}_r$ be pairwise distinct topological or analytical singularity types and $k_1, \dots, k_r \in \mathbb{N} \setminus \{0\}$. Suppose that $V_{|D|}^{\text{irr,fix}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ is non-empty.

Then there exists no curve⁶ $C \in V_{|D|}^{\text{irr}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) \setminus \overline{V_{|D|}^{\text{irr,fix}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)}$ such that for the zero-dimensional scheme $X_0 = X_{\text{fix}}^{\text{es}}(C)$ respectively $X_0 = X_{\text{fix}}^{\text{ea}}(C)$ there exist curves $\Delta_1, \dots, \Delta_m \subset \Sigma$ and zero-dimensional locally complete intersections $X_i^0 \subseteq X_{i-1}$ for $i = 1, \dots, m$, where $X_i = X_{i-1} : \Delta_i$ for $i = 1, \dots, m$ such that $X_S = \bigcup_{i=1}^m X_i^0$ satisfies

$$h^1(\Sigma, \mathcal{I}_{X_0}(D)) + \sum_{i=1}^m \left(h^0(\Sigma, \mathcal{O}_\Sigma(\Delta_i)) - 1 \right) < \#X_S. \quad (3.28)$$

Proof: Throughout the proof we use the notation $V^{\text{irr}} = V_{|D|}^{\text{irr}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$ and $V^{\text{irr,fix}} = V_{|D|}^{\text{irr,fix}}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$.

Suppose there exists a curve $C \in V^{\text{irr}} \setminus \overline{V^{\text{irr,fix}}}$ satisfying the assumption of the Lemma, and let V^* be the irreducible component of V^{irr} containing C . Moreover, let $C_0 \in V^{\text{irr,fix}}$.

We consider in the following the morphism

$$\Phi = \Phi_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) : V_{|D|}(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r) \rightarrow \text{Sym}^{k_1+\dots+k_r}(\Sigma) =: \mathcal{B}$$

from Definition I.2.15.

Step 1: $h^0(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C_0)}/\Sigma(D)) = h^0(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C)}/\Sigma(D)) - h^1(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C)}/\Sigma(D))$.

By the choice of C_0 we have

$$0 = H^1(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C_0)}/\Sigma(D)) \rightarrow H^1(\Sigma, \mathcal{O}_\Sigma(D)) \rightarrow H^1(\Sigma, \mathcal{O}_{X_{\text{fix}}^*(C_0)}(D)) = 0,$$

and thus D is non-special, i. e. $h^1(\Sigma, \mathcal{O}_\Sigma(D)) = 0$. But then

$$\begin{aligned} h^0(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C_0)}/\Sigma(D)) &= h^0(\Sigma, \mathcal{O}_\Sigma(D)) - \deg(X_{\text{fix}}^*(C_0)) \\ &= h^0(\Sigma, \mathcal{O}_\Sigma(D)) - \deg(X_{\text{fix}}^*(C)) \\ &= h^0(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C)}/\Sigma(D)) - h^1(\Sigma, \mathcal{I}_{X_{\text{fix}}^*(C)}/\Sigma(D)). \end{aligned}$$

⁶For a subset $U \subseteq V$ of a topological space V we denote by \overline{U} the closure of U in V .

Step 2: $h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)}(D)) \geq \text{codim}_{\mathcal{B}}(\Phi(V^*))$.

Suppose the contrary, that is $\dim(\Phi(V^*)) < \dim(\mathcal{B}) - h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D))$, then by Step 1, Remark I.2.14 and Equation (I.2.3) in Remark I.2.16

$$\begin{aligned} \dim(V^*) &\leq \dim(\Phi(V^*)) + \dim(\Phi^{-1}(\Phi(C))) \\ &< \dim(\mathcal{B}) - h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D)) + h^0(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)/\Sigma}(D)) - 1 \\ &= 2 \cdot (k_1 + \dots + k_r) + h^0(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C_0)/\Sigma}(D)) - 1 = \dim(V^{\text{irr,fix}}). \end{aligned}$$

However, any irreducible component of V^{irr} has at least the expected dimension $\dim(V^{\text{irr,fix}})$, which gives a contradiction.

Step 3: $\text{codim}_{\mathcal{B}}(\Phi(V^*)) \geq \#X_S - \sum_{i=1}^m \dim|\Delta_i|_1$.

The existence of the subschemes $X_i^0 \subseteq X_{\text{fix}}^*(C) \cap \Delta_i$ imposes at least $\#X_i^0 - \dim|\Delta_i|_1$ conditions on $X_{\text{fix}}^*(C)$ and increases thus the codimension of $\Phi(V^*)$ by the same number.

Step 4: Derive a contradiction.

Collecting the results we derive the following contradiction:

$$\begin{aligned} h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)}(D)) &\geq_{\text{Step 2}} \text{codim}_{\mathcal{B}}(\Phi(V^*)) \\ &\geq_{\text{Step 3}} \#X_S - \sum_{i=1}^m \dim|\Delta_i|_1 >_{(3.28)} h^1(\Sigma, \mathcal{J}_{X_{\text{fix}}^*(C)}(D)). \end{aligned}$$

□

The following two lemmata provide conditions which ensure that $V^{\text{irr,reg}}$ and $V^{\text{irr,fix}}$ share some dense subset $V_{\mathcal{U}}^{\text{gen}}$, and thus that $V^{\text{irr,reg}}$ is dense in $V^{\text{irr,fix}}$.

3.8 Lemma

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, let $D \in \text{Div}(\Sigma)$ and let $V^{\text{irr}} = V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$.

There exists a very general subset $\mathcal{U} \subset \Sigma^r$ such that⁷ $V_{\mathcal{U}}^{\text{gen}} = V_{|D|, \mathcal{U}}^{\text{gen}}(\mathcal{S}_1, \dots, \mathcal{S}_r) = \{C \in V^{\text{irr}} \mid \underline{z} \in \mathcal{U}, (C, z_i) \sim \mathcal{S}_i, i = 1, \dots, r\}$ is dense in $V_{|D|}^{\text{irr,fix}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$.

Proof: This follows from Remark I.2.16 (b). □

3.9 Lemma

With the notation of Lemma 3.8 we assume that

- (a) $(D - K_{\Sigma})^2 \geq 2 \cdot \sum_{i=1}^k (\nu^*(\mathcal{S}_i) + 1)^2$,
- (b) $(D - K_{\Sigma}) \cdot B > \max\{\nu^*(\mathcal{S}_i) \mid i = 1, \dots, r\}$, and
- (c) $D - K_{\Sigma}$ is nef.

Then there exists a very general subset $\mathcal{U} \subset \Sigma^r$ such that $V_{\mathcal{U}}^{\text{gen}} \subseteq V_{|D|}^{\text{irr,reg}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$.

⁷Here \sim means either topological equivalence \sim_t or contact equivalence \sim_c .

Proof: By Theorem II.1.1 we know that there is a very general subset $\mathcal{U} \subset \Sigma^r$ such that for $\underline{z} \in \mathcal{U}$ and $\underline{\nu} = (\nu^*(\mathcal{S}_1), \dots, \nu^*(\mathcal{S}_r))$ we have

$$h^1(\Sigma, \mathcal{J}_{X(\underline{\nu}; \underline{z})/\Sigma}(D)) = 0.$$

However, if $C \in V^{\text{irr}}$ and $\underline{z} \in \mathcal{U}$ with $(C, z_i) \sim \mathcal{S}_i$, then by the definition of $\nu^*(\mathcal{S}_i)$ we have

$$\mathcal{J}_{X(\underline{\nu}; \underline{z})/\Sigma} \hookrightarrow \mathcal{J}_{X(C)/\Sigma},$$

and hence the vanishing of $H^1(\Sigma, \mathcal{J}_{X(\underline{\nu}; \underline{z})/\Sigma}(D))$ implies $h^1(\Sigma, \mathcal{J}_{X(C)/\Sigma}(D)) = 0$, i. e. $C \in V^{\text{irr, reg}}$. \square

In Section 4 we would like to combine the irreducibility results with the existence results from Chapter III. Due to the following lemma this basically comes down to show that for a suitable very ample line bundle L we have (cf. Corollary III.2.4)

$$\gamma \cdot (D - K_\Sigma)^2 \leq \frac{1}{2} \cdot (D - K_\Sigma - L)^2. \quad (3.29)$$

3.10 Lemma

Let \mathcal{S} be a topological or analytical singularity type, then

$$(e^*(\mathcal{S}) + 1)^2 \leq (\tau^*(\mathcal{S}) + 2)^2.$$

Proof: Let us first consider the case that \mathcal{S} is a simple singularity. Then

$$e^*(\mathcal{S}) = e^s(\mathcal{S}) = e^a(\mathcal{S}), \quad \mu(\mathcal{S}) = \tau(\mathcal{S}) = \tau^{es}(\mathcal{S}) = \tau^*(\mathcal{S}), \quad r(\mathcal{S}) \leq \text{mult}(\mathcal{S}) \leq 3$$

and thus

$$2\delta(\mathcal{S}) = \mu(\mathcal{S}) + r(\mathcal{S}) - 1 \leq \mu(\mathcal{S}) + 2 = \tau^*(\mathcal{S}) + 2.$$

But then, once $\tau^*(\mathcal{S}) \geq 5$, we have

$$(e^*(\mathcal{S}) + 1)^2 \leq \frac{625}{48} \cdot \delta(\mathcal{S}) \leq \frac{625}{48 \cdot 2.7} \cdot (\tau^*(\mathcal{S}) + 2)^2 \leq (\tau^*(\mathcal{S}) + 2)^2,$$

while for $\tau^*(\mathcal{S}) \leq 4$, that is for $\mathcal{S} \in \{A_1, A_2, A_3, A_4, D_4\}$ we know $e^*(\mathcal{S})$ precisely and the inequality is fulfilled as well. (Cf. Remark III.2.2.)

We may thus suppose that \mathcal{S} is not simple, i. e. $\text{mod}(\mathcal{S}) \geq 1$.

Let us first consider the case of a topological singularity type. A simple calculation shows that for $\tau^{es}(\mathcal{S}_i) \geq 9$ we always have

$$\frac{625}{48} \cdot \tau^{es}(\mathcal{S}_i) \leq (\tau^{es}(\mathcal{S}) + 2)^2.$$

Moreover, from Remark III.2.2 and Remark I.2.3 we thus deduce

$$(e^s(\mathcal{S}) + 1)^2 \leq \frac{625}{48} \cdot \delta(\mathcal{S}) \leq \frac{625}{48} \cdot \tau^{es}(\mathcal{S}_i) \leq (\tau^{es}(\mathcal{S}) + 2)^2.$$

However, there is only one non-simple topological singularity type with $\tau^{es}(\mathcal{S}_i) \leq 8$, namely X_9 , that is four lines through one point (cf. Remark I.2.4).

But then

$$\delta(\mathcal{S}_i) = \frac{\mu(\mathcal{S}_i) + r(\mathcal{S}_i) - 1}{2} = 6,$$

and thus again

$$(e^s(\mathcal{S}) + 1)^2 \leq \frac{625}{48} \cdot \delta(\mathcal{S}) \leq \frac{625 \cdot 6}{48} \leq 100 = (\tau^{es}(\mathcal{S}) + 2)^2.$$

We, therefore may turn to the case of non-simple analytical singularity types. Since \mathcal{S} is not simple, we know that $\text{mult}(\mathcal{S}) \geq 3$ and $\mu(\mathcal{S}) \geq \tau(\mathcal{S}) \geq 9$. Moreover, we have

$$2\tau(\mathcal{S}) \geq 2\tau^*(\mathcal{S}) \geq \kappa(\mathcal{S}) = \mu(\mathcal{S}) + \text{mult}(\mathcal{S}) - 1,$$

and thus

$$\mu(\mathcal{S}) \leq 2\tau(\mathcal{S}) - \text{mult}(\mathcal{S}) + 1 \leq 2\tau(\mathcal{S}) - 2.$$

Thus, once $\tau(\mathcal{S}) \geq 13$, we have due to Remark III.2.2

$$(e^a(\mathcal{S}) + 1)^2 \leq 9\mu(\mathcal{S}) \leq 18\tau(\mathcal{S}) - 18 \leq (\tau(\mathcal{S}) + 2)^2,$$

and we may suppose that $9 \leq \tau(\mathcal{S}) \leq 12$.

Let us first consider the case that $\text{mod}(\mathcal{S}) \in \{1, 2\}$. Then

$$\mu(\mathcal{S}) \leq \tau(\mathcal{S}) + \text{mod}(\mathcal{S}) \leq \tau(\mathcal{S}) + 2,$$

and hence

$$(e^a(\mathcal{S}) + 1)^2 \leq 9\mu(\mathcal{S}) \leq 9 \cdot (\tau(\mathcal{S}) + 2) \leq (\tau(\mathcal{S}) + 2)^2.$$

It thus remains to consider the case $\text{mod}(\mathcal{S}) \geq 3$, and therefore we also have $\mu(\mathcal{S}) \geq 16$ (cf. Remark I.2.4). If $\text{mod}(\mathcal{S}) = 3$, then $\tau(\mathcal{S}) \geq \mu(\mathcal{S}) - \text{mod}(\mathcal{S}) \geq 16 - 3 = 13$ and we are done. Thus indeed $\text{mod}(\mathcal{S}) \geq 4$.

We do the remaining part by considering the cases $\text{mult}(\mathcal{S}) \geq 5$, $\text{mult}(\mathcal{S}) = 4$ and $\text{mult}(\mathcal{S}) = 3$ separately.

If $\text{mult}(\mathcal{S}) \geq 5$, then

$$\tau(\mathcal{S}) \geq \frac{\mu(\mathcal{S}) + \text{mult}(\mathcal{S}) - 1}{2} \geq \frac{16+5-1}{2} = 10,$$

and therefore

$$(e^a(\mathcal{S}) + 1)^2 \leq 9\mu(\mathcal{S}) \leq 9 \cdot (2\tau(\mathcal{S}) - 4) \leq (\tau(\mathcal{S}) + 2)^2.$$

If $\text{mult}(\mathcal{S}) = 4$, then by Remark I.2.4 $\mu(\mathcal{S}) \geq 22$ and therefore we are done by the following inequality:

$$\tau(\mathcal{S}) \geq \frac{\mu(\mathcal{S}) + \text{mult}(\mathcal{S}) - 1}{2} \geq \frac{22+4-1}{2} = 12\frac{1}{2}.$$

If, finally $\text{mult}(\mathcal{S}) = 3$, then $\mu(\mathcal{S}) \geq 28$ and by the same reasoning we see that $\tau(\mathcal{S}) \geq 15$ and this finishes the proof. \square

4. Examples

4.a. The Classical Case - $\Sigma = \mathbb{P}_{\mathbb{C}}^2$

Since any curve in $\mathbb{P}_{\mathbb{C}}^2$ is non-special and since the Picard number is of course one, the assumptions of Theorem 2.1 are fulfilled. In view of Remark 2.2 the theorem thus reads in this situation.

2.1a Theorem

Let $d \geq 3$, $L \subset \mathbb{P}_{\mathbb{C}}^2$ a line, and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \frac{90}{289} \cdot (d + 3)^2. \quad (2.4a)$$

Then $V_{|dL|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty, irreducible and T-smooth.

Proof: It remains to show that $V_{|dL|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is not empty and T-smooth. However, since $d \geq 3$ we have

$$\frac{90}{289} \cdot (d + 3) \leq \frac{1}{2} \cdot (d + 2) \leq d^2 + 6d$$

and, in view of Lemma 3.10, Corollary III.2.4 applies, as does Theorem IV.1.1a. \square

Many authors were concerned with the question in the case of nodes and cusps, or of nodes and one more complicated singularity, or simply of ordinary multiple points – cf. e. g. [Sev21, BrG81, ArC83, Har85b, Kan89a, Kan89b, Ran89, Shu91b, Shu91a, Bar93a, Shu94, Shu96b, Shu96a, Wal96, GLS98a, GLS98b, Los98, Bru99, GLS00]. Using particularly designed techniques they get of course better results than we may expect to.

The best general results in this case can be found in [GLS00] (see also [Los98] Corollary 6.1). Given a plane curve of degree d , omitting nodes and cusps, they get

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 \leq \frac{9}{10} \cdot d^2$$

as main irreducibility condition. The coefficients differ by a factor of about 3.

4.b. Geometrically Ruled Surfaces

Throughout this section we use the notation of Section G.a. In particular, $\pi: \Sigma \rightarrow C$ is geometrically ruled surface with $g = g(C)$.

In [Ran89] and in [GLS98a] the case of nodal curves on the Hirzebruch surface \mathbb{F}_1 is treated, since this is just $\mathbb{P}_{\mathbb{C}}^2$ blown up in one point. \mathbb{F}_1 is an example of a geometrically ruled surface with invariant $e = 1 > 0$, a case which we cannot treat with the above methods, due to the section with self-intersection -1 .

So far, we are only able to give a general result for geometrically ruled surfaces with invariant $e \leq 0$ – see Theorem 2.4. In the case where $g = 0$, i. e. where Σ is rational, $\mathbb{F}_0 = \mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^1$ is the only surface for which we get any result; and if $g = 1$, by [Har77] V.2.15 we get apart from the product $C \times \mathbb{P}_{\mathbb{C}}^1$ precisely two other surfaces – one with invariant $e = 0$ and one with $e = -1$. For $g \geq 2$ the classes become larger.

2.4b Theorem

Let $\pi : \Sigma \rightarrow C$ be a geometrically ruled surface with $e \leq 0$.

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, and let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_0 + bF$ with $a > \max\{1, \nu^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}$, and, if $g = 0$, $b > \max\{1, \nu^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}$, while if $g > 0$, $b > 2g - 2 + \frac{ae}{2}$.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2, \quad (2.6b)$$

where γ may be taken from the following table with $\alpha = \frac{a+2}{b+2-2g-\frac{ae}{2}} > 0$.

g	e	γ	γ , if $\alpha = 1$
0	0	$\frac{1}{24}$	$\frac{1}{24}$
1	0	$\frac{1}{\max\{24, 2\alpha\}}$	$\frac{1}{24}$
1	-1	$\frac{1}{\max\left\{\min\left\{30 + \frac{16}{\alpha} + 4\alpha, 40 + 9\alpha\right\}, \frac{13}{2}\alpha\right\}}$	$\frac{1}{49}$
≥ 2	0	$\frac{1}{\max\{24 + 16g, 4g\alpha\}}$	$\frac{1}{24 + 16g}$
≥ 2	< 0	$\frac{1}{\max\left\{\min\left\{24 + 16g - 9e\alpha, 18 + 16g - 9e\alpha - \frac{16}{e\alpha}\right\}, 4g\alpha - 9e\alpha\right\}}$	

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or irreducible of the expected dimension.

Addendum 1: If we moreover suppose that $a \geq e^*(\mathcal{S}_i)$ and $b \geq \max\left\{2g - 2 + \frac{ae}{2} + \frac{3}{2} \cdot \left(1 - \frac{e}{2}\right), 2g - 2 + \frac{ae}{2} + \left(1 + \frac{ae}{2}\right) + e^*(\mathcal{S}_i) + 1\right\}$, $l = 1$, if $g = 0$, and else $l \geq 2$ such that $L = C_0 + lF$ is very ample, then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty.

Addendum 2: If $g = 0$, $e = 0$, $\frac{24}{47} \cdot a \leq b \leq \frac{47}{24} \cdot a$, or if $g = 1$, $e = 0$, $b \leq \frac{23}{12} \cdot a + \frac{23}{6}$, then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is also T -smooth.

Proof: First of all we note that for $g > 0$ we have

$$(\tau^*(\mathcal{S}_i) + 2) \leq \frac{1}{2\alpha} \cdot (D - K_\Sigma)^2 = \left(b + 2 - 2g - \frac{ae}{2}\right)^2,$$

and thus

$$b + 2 - 2g - \frac{ae}{2} > \tau^*(\mathcal{S}_i) + 2 > \nu^*(\mathcal{S}_i).$$

Therefore, the conditions of Theorem 2.4 are fulfilled.

Addendum 2 follows from Corollary IV.2.2 and Corollary IV.2.5, since by the assumptions on a and b we have for $g = 0 = e$

$$\frac{1}{24} \cdot (D - K_\Sigma) = \frac{1}{12} \cdot (a + 2) \cdot (b + 2) \leq 4ab + 4a + 4b - a^2 - b^2$$

respectively for $g = 1$ and $e = 0$ we have $b \in [1, a + \sqrt{a^2 + 4a} + 2[$ and

$$\frac{1}{24} \cdot (D - K_\Sigma) = \frac{1}{12} \cdot (ab + 2b) \leq 2ab + 4b - b^2.$$

It remains to prove Addendum 1 with the aid of Corollary III.2.4.

The assumptions on \mathbf{b} can be reformulated as

$$\mathbf{b} - 2g + 2 - \frac{ae}{2} \geq \frac{3}{2} \cdot \left(1 - \frac{\epsilon}{2}\right) \quad (4.1)$$

and

$$(D - L - K_\Sigma).C_0 = \mathbf{b} - 2g + 2 - ae - l \geq e^*(\mathcal{S}_i) + 1.$$

Since moreover $(D - L - K_\Sigma).F = a + 1 > e^*(\mathcal{S}_i)$, Condition (III.2.4) is satisfied, and at the same time

$$\begin{aligned} (D - L - K_\Sigma).L &= (D - L - K_\Sigma).C_0 + l \cdot (D - L - K_\Sigma).F \\ &\geq e^*(\mathcal{S}_i) + 1 + l \cdot (e^*(\mathcal{S}_i) + 1) \geq 2 \cdot e^*(\mathcal{S}_i) + 2 > e^*(\mathcal{S}_i) + 2, \end{aligned}$$

which implies Condition (III.2.5). From (4.1) we deduce that

$$\mathbf{b} - 2g + 2 + e - l \geq \frac{a+1}{2} \cdot e,$$

and hence, since also $a + 1 \geq 0$, $D - L - K_\Sigma$ is nef. It remains to verify Condition (III.2.3), which in view of Lemma 3.10 and $\gamma \leq \frac{1}{24}$ comes down to

$$\frac{1}{24} \cdot (D - K_\Sigma)^2 \leq \frac{1}{2} (D - K_\Sigma - L)^2.$$

We note that due to $a \geq 2$ we have

$$\mathbf{b} - 2g + 2 - \frac{ae}{2} \leq \frac{a+2}{4} \cdot \left(\mathbf{b} - 2g + 2 - \frac{ae}{2}\right) = \frac{1}{8} \cdot (D - K_\Sigma)^2,$$

and by (4.1) we get

$$(a + 1) \cdot \left(1 - \frac{\epsilon}{2}\right) \leq \frac{2}{3} \cdot (a + 1) \cdot \left(\mathbf{b} - 2g + 2 - \frac{ae}{2}\right) \leq \frac{1}{3} \cdot (D - K_\Sigma)^2.$$

These two results then give

$$\frac{1}{2} (D - K_\Sigma - L)^2 = \frac{1}{2} (D - K_\Sigma)^2 - \left(\mathbf{b} - 2g + 2 - \frac{ae}{2} + (a + 1) \cdot \left(1 - \frac{\epsilon}{2}\right)\right) \geq \frac{1}{24} \cdot (D - K_\Sigma)^2$$

which finishes the proof. \square

In the case $g = 0$, that is when $\Sigma \cong \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$, we are in the lucky situation that the constant γ does not at all depend on the chosen divisor D , while in the case $g \geq 1$ the ratio of a and b is involved in γ . This means that an asymptotical behaviour can only be examined if the ratio remains unchanged.

If Σ is a product $C \times \mathbb{P}_\mathbb{C}^1$ the constant γ here is the same as in Section 4.c.

Because of the importance of the case $\mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$ we would like to formulate the result for this surface separately once more in a slightly weaker form, replacing $\nu^*(\mathcal{S}_i)$ and $e^*(\mathcal{S}_i)$ by $\tau^*(\mathcal{S}_i) + 1$, which is an upper bound for both of them.

4.1 Corollary

Let $\Sigma \cong \mathbb{F}_0 = \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$, $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types and $D \in \text{Div}(\Sigma)$ such that $D \sim_l aC_0 + bF$ with $a, b \geq \tau^*(\mathcal{S}_i) + 1$ for $i = 1, \dots, r$.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \frac{1}{24} \cdot (D - K_\Sigma)^2 = \frac{1}{12} \cdot (a + 2) \cdot (b + 2). \quad (2.6b)$$

Then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty, irreducible and of the expected dimension.

Addendum: If, moreover, $\frac{24}{47} \cdot a \leq b \leq \frac{47}{24} \cdot a$, then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is T -smooth.

4.c. Products of Curves

Throughout this section we use the notation of Section G.b. In particular, $\Sigma = C_1 \times C_2$ where C_1 and C_2 are smooth projective curves over \mathbb{C} of genera g_1 and g_2 respectively.

For a generic choice of C_1 and C_2 the Néron–Severi group $\text{NS}(\Sigma)$ is two-dimensional by Proposition G.12. Thus Theorem 2.3 gives the following result for a generic product surface.

2.3c Theorem

Let C_1 and C_2 be two smooth projective curves of genera g_1 and g_2 respectively with $g_1 \geq g_2 \geq 0$, such that for $\Sigma = C_1 \times C_2$ the Néron–Severi group is $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z}$.

Let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, and let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_1 + bC_2$ with

$$a > \begin{cases} \max \{1, \nu^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}, & \text{if } g_2 = 0 \\ 2g_2 - 2, & \text{else,} \end{cases}$$

and

$$b > \begin{cases} \max \{1, \nu^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}, & \text{if } g_1 = 0 \\ 2g_1 - 2, & \text{else.} \end{cases}$$

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \gamma \cdot (D - K_\Sigma)^2, \quad (2.5c)$$

where γ may be taken from the following table with $\alpha = \frac{a-2g_2+2}{b-2g_1+2} > 0$.

g_1	g_2	γ	$\gamma, \text{ if } \alpha = 1$
0	0	$\frac{1}{24}$	$\frac{1}{24}$
1	0	$\frac{1}{\max\{32, 2\alpha\}}$	$\frac{1}{32}$
≥ 2	0	$\frac{1}{\max\{24+16g_1, 4g_1\alpha\}}$	$\frac{1}{24+16g_1}$
1	1	$\frac{1}{\max\{32, 2\alpha, \frac{2}{\alpha}\}}$	$\frac{1}{32}$
≥ 2	≥ 1	$\frac{1}{\max\{24+16g_1+16g_2, 4g_1\alpha, \frac{4g_2}{\alpha}\}}$	$\frac{1}{24+16g_1+16g_2}$

Then either $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or irreducible of the expected dimension.

Addendum 1: *If we moreover suppose that*

$$a > \begin{cases} \max \{e^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}, & \text{if } g_2 = 0 \\ 2g_2 - 2 + \frac{3l}{2}, & \text{else,} \end{cases}$$

and

$$b > \begin{cases} \max \{e^*(\mathcal{S}_i) - 2 \mid i = 1, \dots, r\}, & \text{if } g_1 = 0 \\ 2g_1 - 2 + \frac{3l}{2}, & \text{else,} \end{cases}$$

where l is a positive integer such that $L = lC_1 + lC_2$ respectively $L = C_1 + lC_2$, if $g_2 = 0$, is very ample, then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty.

Addendum 2: *If $g_2 \geq 2$, $a = p \cdot (g_2 - 1)$, $b = p \cdot (g_1 - 1)$, i. e. $D \sim_a \frac{p}{2} \cdot K_\Sigma$, and if we suppose that $p > \frac{87}{176} \cdot (\tau^*(\mathcal{S}_i) + 1)$, then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is also T -smooth.*

Proof: First we note that for $g_1 > 0$ we have

$$(\tau^*(\mathcal{S}_i) + 2) \leq \frac{1}{2\alpha} \cdot (D - K_\Sigma)^2 = (b + 2 - 2g_1)^2,$$

and thus

$$b + 2 - 2g_1 > \tau^*(\mathcal{S}_i) + 2 > \nu^*(\mathcal{S}_i).$$

Similarly, if $g_2 > 0$, then $a + 2 - 2g_2 > \nu^*(\mathcal{S}_i)$. Therefore, the conditions of Theorem 2.4 are fulfilled.

For Addendum 2 we may apply Theorem IV.1.1c and the fact that

$$\gamma \cdot \frac{(D - K_\Sigma)^2}{(g_1 - 1) \cdot (g_2 - 1)} \leq \frac{1}{44} \cdot (p - 2)^2 \leq \frac{1}{44} \cdot p^2 \leq 2 \cdot (p^2 - 2 \cdot p \cdot (\tau^*(\mathcal{S}_i) + 1)).$$

It remains to prove Addendum 1 with the aid of Corollary III.2.4. For this we note that by Lemma 3.10 we have $e^*(\mathcal{S}_i) < \tau^*(\mathcal{S}_i) + 2$. But then Due to the assumptions on a and b and the above considerations, we have that $D - L - K_\Sigma$ is nef and that the intersection of $D - L - K_\Sigma$ with any irreducible curve of self-intersection zero, that is with the fibres C_1 and C_2 , is greater than $e^*(\mathcal{S}_i)$. Moreover,

$$D \cdot L - 2g(L) + 2 = (D - L - K_\Sigma) \cdot L \geq \frac{1}{3} \cdot (a - 2g_2 + 2 + b - 2g_1 + 2) > e^*(\mathcal{S}_i) + 2.$$

It therefore remains to show that Condition (III.2.3) is satisfied, which in view of Lemma 3.10 and $\gamma \leq \frac{1}{24}$ amounts to showing

$$\frac{1}{12} \cdot (a - 2g_2 + 2) \cdot (b - 2g_1 + 2) \leq (a - 2g_2 + 2 - l) \cdot (b - 2g_1 + 2 - l).$$

This, however, is again fulfilled due to the assumptions on a and b . \square

Once more, only in the case $\Sigma \cong \mathbb{P}_c^1 \times \mathbb{P}_c^1$ we get a constant γ which does not depend on the chosen divisor D , while in the remaining cases the ratio of a and b is involved in γ . This means that an asymptotical behaviour can only be examined if the ratio remains unchanged.

4.d. Products of Elliptic Curves

If in Section 4.c the curves C_1 and C_2 are chosen to be both elliptic curves, then “ C_1 and C_2 are generic” means precisely that they are not isogenous (cf. p. 154). We thus get the following theorem.

2.3d Theorem

Let $\Sigma = C_1 \times C_2$, where C_1 and C_2 are two non-isogenous elliptic curves, let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types, and let $D \in \text{Div}(\Sigma)$ such that $D \sim_a aC_1 + bC_2$ with $a, b \geq 5$.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \min \left\{ \frac{1}{32}, \frac{a}{2b}, \frac{b}{2a} \right\} \cdot D^2. \quad (2.5d)$$

Then $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty, irreducible and T -smooth.

Proof: We note that due to Theorem IV.1.1d $V_{|D|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is T -smooth, and thus we are done by Theorem 2.3c. \square

4.e. Complete Intersection Surfaces

Throughout this section we use the notations of Section G.d.

In Proposition G.20 we show that a smooth complete intersection surface, always satisfies Condition (ii) of Theorem 2.1 for hypersurface-sections. Thus, a complete intersection with Picard number one satisfies the assumptions of Theorem 2.1.

2.1e Theorem

Let $\mathbb{P}_\mathbb{C}^2 \not\cong \Sigma \subset \mathbb{P}_\mathbb{C}^N$ be a smooth complete intersection of type (d_1, \dots, d_{N-2}) , let $H \subset \Sigma$ be a hyperplane section and suppose that the Picard number of Σ is one.

Let $d > \kappa = \sum_{i=1}^{N-2} d_i - N - 1 \geq 0$, $n = H^2 = d_1 \cdots d_{N-2}$ and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \frac{18 \cdot ((\kappa+3) \cdot n + 2\chi(\mathcal{O}_\Sigma)) \cdot n^2}{((3\kappa+11) \cdot n + 6\chi(\mathcal{O}_\Sigma))^2} \cdot (d - \kappa)^2. \quad (2.4e)$$

Then either $V_{|dH|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is empty or irreducible of the expected dimension.

Proof: It remains to show $\kappa = -N - 1 + \sum_{i=1}^{N-2} d_i \geq 0$, then in particular $dH - K_\Sigma = (d - \kappa) \cdot H$ is ample and $dH + K_\Sigma = (d + \kappa) \cdot H$ is nef.

Since $\kappa \geq 2N - 4 - N - 1 = N - 5$ anyway, the critical situations are $N = 3$ with $d_1 \leq 3$, and $N = 4$ with $d_1 = d_2 = 2$. In these cases the surface Σ is either $\mathbb{P}_\mathbb{C}^2$ or rational with a Picard number larger than one (see p. 160 and [Har77] Ex. V.4.13). This finishes the claim. \square

By Proposition G.20 we even do know a formula for $\chi(\mathcal{O}_\Sigma)$, however, the formula would just become nastier.

4.f. Surfaces in $\mathbb{P}_\mathbb{C}^3$

For a generic surface in $\mathbb{P}_\mathbb{C}^3$ we know that the Picard number is one, since then any curve is a hypersurface section by a Theorem of Noether (see Section G.d). Being a complete intersection, the results of Section 4.e apply for generic hypersurfaces in $\mathbb{P}_\mathbb{C}^3$. However, in this case the invariants involved in the formulation are much simpler.

2.1f Theorem

Let $\Sigma \subset \mathbb{P}_\mathbb{C}^3$ be a smooth hypersurface of degree $n \geq 4$, let $H \subset \Sigma$ be a hyperplane section, and suppose that the Picard number of Σ is one.

Let $d \geq n + 6$ and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \frac{6 \cdot (n^3 - 3n^2 + 8n - 6) \cdot n^2}{(n^3 - 3n^2 + 10n - 6)^2} \cdot (d + 4 - n)^2. \quad (2.4f)$$

Then $V_{|dH|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty and irreducible of the expected dimension.

Addendum: If we, moreover, assume that $d \geq n \cdot (\tau^*(\mathcal{S}_i) + 1)$, then $V_{|dH|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is T -smooth.

Proof: For the Addendum we note that $\frac{6 \cdot (n^3 - 3n^2 + 8n - 6) \cdot n^2}{(n^3 - 3n^2 + 10n - 6)^2} \leq 2$ for $n \geq 4$, and hence

$$nd^2 - d \cdot (n - 4) \cdot n \cdot (\tau^*(\mathcal{S}_i) + 1) \geq 4d^2 \geq \frac{6 \cdot (n^3 - 3n^2 + 8n - 6) \cdot n^2}{(n^3 - 3n^2 + 10n - 6)^2} \cdot (d + 4 - n)^2,$$

and the result thus follows from Theorem IV.1.1e.

It remains to show that $V_{|dH|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty, using Corollary III.2.4. With $L = H$ we have $D - L - K_\Sigma = (d - n + 3) \cdot H$, and since $d \geq n + 6$ the divisor $D - L - K_\Sigma$ is nef. We thus have to verify the Conditions (III.2.3) and (III.2.5) in Corollary III.2.4. However, once (III.2.3) is fulfilled, then

$e^*(\mathcal{S}_i) + 1 \leq \sqrt{n} \cdot (d - n + 4) < 2 \cdot \sqrt{n} \cdot (d - n + 3) - 1 \leq n \cdot (d - n + 3) - 1 = D \cdot L - 2g(L) + 1$ for $i = 1, \dots, r$, that is, (III.2.5) is satisfied as well.

Taking Lemma 3.10 into account (III.2.3) comes down to showing that

$$\frac{6 \cdot (n^3 - 3n^2 + 8n - 6) \cdot n}{(n^3 - 3n^2 + 10n - 6)^2} \cdot n \cdot (d - n + 4)^2 \leq \frac{1}{2} \cdot n \cdot (d - n + 3)^2.$$

We claim that the function

$$\gamma : [4, \infty) \rightarrow \mathbb{R} : x \mapsto \frac{6 \cdot (x^3 - 3x^2 + 8x - 6) \cdot x}{(x^3 - 3x^2 + 10x - 6)^2}$$

is monotonously decreasing. In order to see this, it suffices to show that the derivative

$$\gamma'(x) = \frac{-6 \cdot (2x^6 - 9x^5 + 21x^4 - 24x^3 + 36x - 36)}{(x^3 - 3x^2 + 10x - 6)^3} < 0$$

for all $x \in [4, \infty)$, or equivalently that $2x^6 - 9x^5 + 21x^4 - 24x^3 + 36x - 36$ is strictly positive in the interval $[4, \infty)$. However, since $x \geq 4$ we get

$$(2x^6 - 9x^5 + 6x^4) + (15x^4 - 24x^3) + (36x - 36) > 0$$

since each summand is strictly positive.

It follows that $\gamma(n) \leq \gamma(4) = \frac{252}{625}$ for all $n \geq 4$, and it remains to show that

$$\frac{252}{625} \cdot (d - n + 4)^2 \leq \frac{1}{2} \cdot (d - n + 3)^2,$$

which is a fairly easy calculation, since $d - n + 3 \geq 9$. \square

Note that a slightly more sophisticated investigation would have allowed to replace the restriction $d \geq n + 6$ by $d \geq \max\{n - 1, 10\}$.

Calculating the invariants in (2.4f) for the examples of reducible families of irreducible nodal curves on surfaces in $\mathbb{P}_\mathbb{C}^3$ given in the introduction on page 81 we end up with

$$\begin{aligned} & \frac{6 \cdot (n^3 - 3n^2 + 8n - 6) \cdot n^2}{(n^3 - 3n^2 + 10n - 6)^2} \cdot (d + 4 - n)^2 \\ & < \frac{6}{n-3} \cdot (d + 4 - n)^2 < \frac{6}{n-3} \cdot (d^2 + (4 - n) \cdot d + 2) \\ & < \frac{9n}{2} \cdot (d^2 + (4 - n) \cdot d + 2) = 9r = \sum_{i=1}^r (\tau^*(A_i) + 2)^2, \end{aligned}$$

that is, our result fits with these families.

4.g. Generic K3-Surfaces

A generic K3-surface has Picard number one (cf. Section G.e). Moreover, if Σ is a K3-surface with $\text{NS}(\Sigma) = \mathbb{L} \cdot \Sigma$ and if $C \sim_{\mathbb{L}} d\mathbb{L}$ is a curve in Σ , then by Lemma E.1 C is ample. Thus by the Kodaira Vanishing Theorem we have $h^1(\Sigma, C) = h^1(\Sigma, K_\Sigma + C) = 0$. Hence the surface satisfies the assumptions of Theorem 2.1.

2.1g Theorem

Let Σ be a smooth K3-surface with $\text{NS}(\Sigma) = \mathbb{L} \cdot \mathbb{Z}$ and set $n = \mathbb{L}^2$.

Let $d \geq 19$ and let $\mathcal{S}_1, \dots, \mathcal{S}_r$ be topological or analytical singularity types.

Suppose that

$$\sum_{i=1}^r (\tau^*(\mathcal{S}_i) + 2)^2 < \frac{54n^2 + 72n}{(11n + 12)^2} \cdot d^2 \cdot n. \quad (2.4g)$$

Then $V_{|d\mathbb{L}|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is irreducible and T -smooth.

Proof: Apart from T-smoothness, it remains to prove that $V_{|dL|}^{\text{irr}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$ is non-empty and as in the case of surfaces in \mathbb{P}_c^3 for this it suffices to show that (III.2.3) and (III.2.5) in Corollary III.2.4 are satisfied, where once again the first condition implies the latter one due to

$$e^*(\mathcal{S}_i) + 1 \leq \sqrt{n} \cdot d < 2 \cdot \sqrt{n} \cdot (d - 1) - 1 \leq n \cdot (d - 1) - 1 = D.L - 2g(L) + 1.$$

Taking Lemma 3.10 into account it thus remains to show

$$\frac{54n^2 + 72n}{(11n + 12)^2} \cdot d^2 \cdot n \leq \frac{1}{2} \cdot (d - 1)^2 \cdot n.$$

We consider the function

$$\gamma : [0, \infty) \rightarrow \mathbb{R} : x \mapsto \frac{54x^2 + 72x}{(11x + 12)^2},$$

which converges to $\frac{54}{121}$ for $x \mapsto \infty$ and is actually bounded from above by this number. It thus suffices to show

$$\frac{54}{121} \cdot d^2 \leq \frac{1}{2} \cdot (d - 1)^2,$$

which is satisfied, since $d \geq 19$.

T-smoothness follows from Theorem IV.1.1f, since $\frac{54n^2 + 72n}{(11n + 12)^2} < 1$. □

Appendix

A. Very General Position

It is our first aim to show that if there is a curve passing through points $z_1, \dots, z_r \in \Sigma$ in very general position with multiplicities n_1, \dots, n_r , then it can be equimultiply deformed in its algebraic system in a good way - i. e. suitable for Lemma II.2.3.

For the convenience of the reader we recall the definition of a *very general subset* from the Introduction.

A.1 Definition

Let X be any Zariski topological space.

We say a subset $U \subseteq X$ is *very general* if it is an at most countable intersection of open dense subsets of X .

Some statement is said to hold for points $z_1, \dots, z_r \in X$ (or $\underline{z} \in X^r$) *in very general position* if there is a suitable very general subset $U \subseteq X^r$, contained in the complement of the closed subvariety $\bigcup_{i \neq j} \{z \in X^r \mid z_i = z_j\}$ of X^r , such that the statement holds for all $\underline{z} \in U$.

In the following we suppose that $\Sigma \subseteq \mathbb{P}_c^n$ is an embedded smooth projective surface. We will denote by Hilb_Σ the Hilbert scheme of curves on Σ , and for $h \in \mathbb{Q}[x]$ by Hilb_Σ^h the Hilbert scheme of curves on Σ with Hilbert polynomial h . The latter is a projective variety, and has in particular only finitely many connected components. If $B \subset \Sigma$ is a curve with Hilbert polynomial h , then we denote $|B|_a$ the connected component of the reduction of Hilb_Σ^h containing B . (Cf. [Mum66] Chapter 15.)

A.2 Lemma

Let $B \subset \Sigma$ be a curve, and $\underline{n} \in \mathbb{N}_0^r$. Then

$$V_{B, \underline{n}} = \{ \underline{z} \in \Sigma^r \mid \exists C \in |B|_a : \text{mult}_{z_i}(C) \geq n_i \forall i = 1, \dots, r \}$$

is a closed subset of Σ^r .

Proof:

Step 1: Show first that for $n \in \mathbb{N}_0$

$$X_{B, n} := \{ (C, z) \in H \times \Sigma \mid \text{mult}_z(C) \geq n \}$$

is a closed subset of $H \times \Sigma$, where $H := |B|_a$.

Being the reduction of a connected component of the Hilbert scheme Hilb_Σ , H is a projective variety endowed with a universal family of curves, giving rise

to the following diagram of morphisms

$$\begin{array}{ccc} \mathcal{C} = \bigcup_{C \in H} \{C\} \times C & \hookrightarrow & H \times \Sigma \xrightarrow{\text{pr}_\Sigma} \Sigma \\ & \searrow & \downarrow \text{pr}_H \\ & & H, \end{array}$$

where \mathcal{C} is an effective Cartier divisor on $H \times \Sigma$ with $\mathcal{C}_{\{C\} \times \Sigma} = C$.⁸

Let $s \in H^0(H \times \Sigma, \mathcal{O}_{H \times \Sigma}(\mathcal{C}))$ be a global section defining \mathcal{C} . Then⁹

$$X_{B,n} = \{\eta = (C, z) \in H \times \Sigma \mid s_\eta \in (\mathfrak{m}_{\Sigma,z}^n + \mathfrak{m}_{H,C}) \cdot \mathcal{O}_{H \times \Sigma, \eta}\}.$$

We may consider a finite open affine covering of $H \times \Sigma$ of the form $\{H_i \times U_j \mid i \in I, j \in J\}$, $H_i \subset H$ and $U_j \subset \Sigma$ open, such that \mathcal{C} is locally on $H_i \times U_j$ given by one polynomial equation, say¹⁰

$$s_{i,j}(\underline{a}, \underline{b}) = 0, \text{ for } \underline{a} \in H_i, \underline{b} \in U_j.$$

It suffices to show that $X_{B,n} \cap (H_i \times U_j)$ is closed in $H_i \times U_j$ for all i, j .

However, for $\eta = (C, z) = (\underline{a}, \underline{b}) \in H_i \times U_j$ we have

$$s_\eta \in (\mathfrak{m}_{\Sigma,z}^n + \mathfrak{m}_{H,C}) \cdot \mathcal{O}_{H \times \Sigma, \eta}$$

if and only if

$$s_{i,j}(\underline{a}, \underline{b}) = 0 \text{ and}$$

$$\frac{\partial^\alpha s_{i,j}}{\partial \underline{b}^\alpha}(\underline{a}, \underline{b}) = 0, \text{ for all } |\alpha| \leq n-1,$$

where α is a multiindex. Thus,

$$X_{B,n} \cap (H_i \times U_j) = \left\{ (\underline{a}, \underline{b}) \in H_i \times U_j \mid s_{i,j}(\underline{a}, \underline{b}) = 0 = \frac{\partial^\alpha s_{i,j}}{\partial \underline{b}^\alpha}(\underline{a}, \underline{b}), \forall |\alpha| \leq n-1 \right\}$$

is a closed subvariety of $H_i \times U_j$, since $s_{i,j}$ and the $\frac{\partial^\alpha s_{i,j}}{\partial \underline{b}^\alpha}(\underline{a}, \underline{b})$ are polynomial expressions in \underline{a} and \underline{b} .

Step 2: $V_{B,\underline{n}}$ is a closed subset of Σ^r .

By Step 1 for $i = 1, \dots, r$ the set

$$X_{B,\underline{n},i} := \{(\underline{z}, C) \in \Sigma^r \times H \mid \text{mult}_{z_i}(C) \geq n_i\} \cong \Sigma^{r-1} \times X_{B,n_i}$$

is a closed subset of $\Sigma^r \times H \cong \Sigma^{r-1} \times H \times \Sigma$. Considering now

$$\begin{array}{ccc} X_{B,\underline{n}} := \bigcap_{i=1}^r X_{B,\underline{n},i} & \hookrightarrow & \Sigma^r \times H \\ & \searrow \rho & \downarrow \\ & & \Sigma^r, \end{array}$$

⁸For the definition of an algebraic family of curves see [Har77] Ex. V.1.7.

⁹ $\mathfrak{m}_{\Sigma,z}^n = (\mathfrak{m}_{\Sigma,z}^n + \mathfrak{m}_{H,C}) \cdot \mathcal{O}_{H \times \Sigma, \eta} / \mathfrak{m}_{H,C} \cdot \mathcal{O}_{H \times \Sigma, \eta}$ and $C = \mathcal{C}_{\{C\} \times \Sigma}$ is locally in z given by the image of s_η in $\mathcal{O}_{\Sigma,z} = \mathcal{O}_{H \times \Sigma, \eta} / \mathfrak{m}_{H,C} \cdot \mathcal{O}_{H \times \Sigma, \eta}$.

¹⁰The \underline{a} and \underline{b} denote coordinates on the affine ambient spaces of $H_i \subseteq \mathbb{A}^{N_i}$ respectively $U_j \subseteq \mathbb{A}^{M_j}$.

we find that $V_{B, \underline{n}} = \rho(X_{B, \underline{n}})$, being the image of a closed subset under a morphism between projective varieties, is a closed subset of Σ^r (cf. [Har77] Ex. II.4.4). \square

A.3 Corollary

Then the complement of the set

$$V = \bigcup_{B \in \text{Hilb}_\Sigma} \bigcup_{\underline{n} \in \mathbb{N}_0^r} \bigcup_{V_{B, \underline{n}} \neq \Sigma^r} V_{B, \underline{n}}$$

in Σ^r is very general, where Hilb_Σ is the Hilbert scheme of curves on Σ .

In particular, there is a very general subset $U \subseteq \Sigma^r$ such that if for some $\underline{z} \in U$ there is a curve $B \subset \Sigma$ with $\text{mult}_{z_i}(B) = n_i$ for $i = 1, \dots, r$, then for any $\underline{z}' \in U$ there is a curve $B' \in |B|_a$ with $\text{mult}_{z'_i}(B') \geq n_i$.

Proof: Fixing some embedding $\Sigma \subseteq \mathbb{P}_\mathbb{C}^n$ and $h \in \mathbb{Q}[x]$, Hilb_Σ^h is a projective variety and has thus only finitely many connected components. Thus the Hilbert scheme Hilb_Σ has only a countable number of connected components, and we have only a countable number of different $V_{B, \underline{n}}$, where B runs through Hilb_Σ and \underline{n} through \mathbb{N}_0^r . By Lemma A.2 the sets $V_{B, \underline{n}}$ are closed, hence their complements $\Sigma^r \setminus V_{B, \underline{n}}$ are open. But then

$$U = \Sigma^r \setminus V = \bigcap_{B \in \text{Hilb}_\Sigma} \bigcap_{\underline{n} \in \mathbb{N}_0^r} \bigcap_{V_{B, \underline{n}} \neq \Sigma^r} (\Sigma^r \setminus V_{B, \underline{n}})$$

is an at most countable intersection of open dense subsets of Σ^r , and is hence very general. \square

If Σ is regular, i. e. the irregularity $q(\Sigma) = h^1(\Sigma, \mathcal{O}_\Sigma) = 0$, algebraic and linear equivalence coincide, and thus $\text{Hilb}_\Sigma^h = |D|_h$, if D is any divisor with Hilbert polynomial h . This makes the proofs given above a bit simpler.

A.4 Lemma

Given $\underline{n} \in \mathbb{N}_0^r$ and $D \in \text{Pic}(\Sigma)$, the set

$$V_{D, \underline{n}} = \{ \underline{z} \in \Sigma^r \mid \exists C \in |D|_1 : \text{mult}_{z_i}(C) \geq n_i \}$$

is a closed subset of Σ^r .

Proof: Fix an affine covering $\Sigma = U_1 \cup \dots \cup U_k$ of Σ , and a basis s_0, \dots, s_n of $H^0(\Sigma, \mathcal{O}_\Sigma(D))$.

It suffices to show that $V_{D, \underline{n}} \cap (U_{j_1} \times \dots \times U_{j_r})$ is closed in $U_{j_1} \times \dots \times U_{j_r}$ for all $\underline{j} = (j_1, \dots, j_r) \in \{1, \dots, k\}^r$, since those sets form an open covering of $V_{D, \underline{n}}$.

Consider the set

$$V_{\alpha, i, \underline{j}} = \{ (\underline{z}, \underline{a}) \in U_{j_1} \times \dots \times U_{j_r} \times \mathbb{P}_\mathbb{C}^n \mid a_0(D^\alpha s_0)(z_i) + \dots + a_n(D^\alpha s_n)(z_i) = 0 \},$$

where α is a multiindex and D^α denotes the corresponding differential operator. This is a closed subvariety of $U_{j_1} \times \dots \times U_{j_r} \times \mathbb{P}_\mathbb{C}^n$.¹¹ Thus, also the set

$$V_{\underline{j}} = \bigcap_{i=1, \dots, r} \bigcap_{|\alpha| \leq n_i - 1} V_{\alpha, i, \underline{j}}$$

is a closed subset of $U_{j_1} \times \dots \times U_{j_r} \times \mathbb{P}_\mathbb{C}^n$. Considering now the projection to $U_{j_1} \times \dots \times U_{j_r}$,

$$\begin{array}{ccc} V_{\underline{j}} & \hookrightarrow & U_{j_1} \times \dots \times U_{j_r} \times \mathbb{P}_\mathbb{C}^n \\ & \searrow \rho & \downarrow \\ & & U_{j_1} \times \dots \times U_{j_r}, \end{array}$$

we find that $\text{Im}(\rho) = V \cap (U_{j_1} \times \dots \times U_{j_r})$ is closed in $U_{j_1} \times \dots \times U_{j_r}$ (cf. [Har92] Theorem 3.12). \square

A.5 Corollary

Let Σ be regular. Then the complement of the set

$$V = \bigcup_{D \in \text{Pic}(\Sigma)} \bigcup_{\underline{n} \in \mathbb{N}_0^r} \bigcup_{V_{D, \underline{n}} \neq \Sigma^r} V_{D, \underline{n}}$$

in Σ^r is very general.

In particular, there is a very general subset $U \subseteq \Sigma^r$ such that if for some $\underline{z} \in U$ there is a curve $B \subset \Sigma$ with $\text{mult}_{z_i}(B) = n_i$ for $i = 1, \dots, r$, then for any $\underline{z}' \in U$ there is a curve $B' \in |B|_1$ with $\text{mult}_{z'_i}(B') \geq n_i$.

Proof: By Lemma A.4 the sets $V_{D, \underline{n}}$ are closed, hence their complement $\Sigma^r \setminus V_{D, \underline{n}}$ is open. Since Σ is regular, $\text{Pic}(\Sigma) \cong \text{NS}(\Sigma)$ is a finitely generated abelian group, hence countable.¹² But then

$$U = \Sigma^r \setminus V = \bigcap_{D \in \text{Pic}(\Sigma)} \bigcap_{\underline{n} \in \mathbb{N}_0^r} \bigcap_{V_{D, \underline{n}} \neq \Sigma^r} (\Sigma^r \setminus V_{D, \underline{n}})$$

is an at most countable intersection of open dense subsets of Σ^r and is hence very general. \square

In the proof of Theorem II.1.1 we use at some place the result of Corollary A.6. We could instead use Corollary A.3. However, since the results are quite nice and simple to prove we just give them.

A.6 Corollary

- (a) The number of curves B in Σ with $\dim |B|_a = 0$ is at most countable.

¹¹On U_j the s_i are given as quotients of polynomials where the denominator is not vanishing on U_j . Thus we may assume w. l. o. g. that s_i is represented by a polynomial, and hence the above equation is polynomial in z_i and even linear in the a_i .

¹²See [IsS96] Chapters 3.2 and 3.3. From the exponential sequence it follows that $\text{NS}(\Sigma) \cong \text{Pic}(\Sigma)/\text{Pic}^0(\Sigma)$ with $\text{Pic}^0(\Sigma) \cong \mathbb{C}^q/\mathbb{Z}^{2q}$, $q = q(\Sigma)$. Thus, $\text{Pic}(\Sigma)$ is countable if and only if Σ is regular.

- (b) The number of exceptional curves in Σ (i. e. curves with negative self intersection) is at most countable.
- (c) There is a very general subset U of Σ^r , $r \geq 1$, such that for $\underline{z} \in U$ no z_i belongs to a curve $B \subset \Sigma$ with $\dim |B|_{\alpha} = 0$, in particular to no exceptional curve.

Proof:

- (a) By definition $|B|_{\alpha}$ is a connected component of Hilb_{Σ} , whose number is at most countable (see proof of Lemma A.3). If in addition $\dim |B|_{\alpha} = 0$, then $|B|_{\alpha} = \{B\}$ which proves the claim.
- (b) Curves of negative self-intersection are not algebraically equivalent to any other curve (cf. [IsS96] . p. 153)
- (c) Follows from (a).

□

A.7 Example (Kodaira)

Let $z_1, \dots, z_9 \in \mathbb{P}_{\mathbb{C}}^2$ be in very general position¹³ and let $\Sigma = \text{Bl}_{\underline{z}}(\mathbb{P}_{\mathbb{C}}^2)$ be the blow up of $\mathbb{P}_{\mathbb{C}}^2$ in $\underline{z} = (z_1, \dots, z_9)$. Then Σ contains infinitely many irreducible smooth rational -1 -curves, i. e. exceptional curves of the first kind.

Proof: It suffices to find an infinite number of irreducible curves C in $\mathbb{P}_{\mathbb{C}}^2$ such that

$$d^2 - \sum_{i=1}^9 m_i^2 = -1, \tag{A.1}$$

and

$$p_a(C) - \sum_{i=1}^9 \frac{m_i(m_i - 1)}{2} = 0, \tag{A.2}$$

where $m_i = \text{mult}_{z_i}(C)$ and $d = \text{deg}(C)$, since the expression in (A.1) is the self intersection of the strict transform $\tilde{C} = \text{Bl}_{\underline{z}}(C)$ of C and (A.2) gives its arithmetical genus.¹⁴ In particular \tilde{C} cannot contain any singularities, since they would contribute to the arithmetical genus, and, being irreducible anyway, \tilde{C} is an exceptional curve of the first kind.

¹³To be precise, no three of the nine points should be collinear, and after any finite number of quadratic Cremona transformations centred at the z_i (respectively the newly obtained centres) still no three should be collinear. Thus the admissible tuples in $(\mathbb{P}_{\mathbb{C}}^2)^9$ form a very general set, cf. [Har77] Ex. V.4.15.

¹⁴

$$\begin{aligned} p_a(\tilde{C}) &= 1 + \frac{K_{\Sigma} \cdot \tilde{C} + \tilde{C}^2}{2} = 1 + \frac{(K_{\mathbb{P}^2} + \sum E_i) \cdot (C - \sum m_i E_i) + (C - \sum m_i E_i)^2}{2} \\ &= 1 + \frac{K_{\mathbb{P}^2} \cdot C + C^2}{2} - \sum \frac{m_i(m_i - 1)}{2} = p_a(C) - \sum \frac{m_i(m_i - 1)}{2}. \end{aligned}$$

We are going to deduce the existence of these curves with the aid of quadratic Cremona transformations.

Claim: If for some $d > 0$ and $m_1, \dots, m_9 \geq 0$ with $3d - \sum_{i=1}^9 m_i = 1$ there is an irreducible curve $C \in |\mathcal{J}_{X(\underline{m}; \underline{z})}(d)|_1$, then $T(C) \in |\mathcal{J}_{X(\underline{m}'; \underline{z}')}(d+a)|_1$ is an irreducible curve, where

- $\{i, j, k\} \subset \{1, \dots, 9\}$ are such that $m_i + m_j + m_k < d$,
- $T : \mathbb{P}_{\mathbb{C}}^2 \dashrightarrow \mathbb{P}_{\mathbb{C}}^2$ is the quadratic Cremona transformation at z_i, z_j, z_k ,
- $z'_\nu = \begin{cases} z_\nu, & \text{if } \nu \neq i, j, k, \\ T(\overline{z_\lambda z_\mu}), & \text{if } \{\nu, \lambda, \mu\} = \{i, j, k\}, \end{cases}$
- $m'_\nu = \begin{cases} m_\nu, & \text{if } \nu \neq i, j, k, \\ m_\nu + a, & \text{else, and} \end{cases}$
- $a = d - (m_i + m_j + m_k)$.

Note that, $3 \cdot (d+a) - \sum_{i=1}^9 m'_i = 1$, i. e. we may iterate the process since the hypothesis of the claim will be preserved.

Since $3d > \sum_{i=1}^9 m_i$, there must be a triple (i, j, k) such that $d > m_i + m_j + m_k$.

Let us now consider the following diagram

$$\begin{array}{ccc} \Sigma = \text{Bl}_{z_i, z_j, z_k}(\mathbb{P}_{\mathbb{C}}^2) = \text{Bl}_{z'_i, z'_j, z'_k}(\mathbb{P}_{\mathbb{C}}^2) & & \\ \swarrow \pi & & \searrow \pi' \\ \mathbb{P}_{\mathbb{C}}^2 & \xleftarrow{\quad T \quad} & \mathbb{P}_{\mathbb{C}}^2 \end{array}$$

and let us denote the exceptional divisors of π by E_i and those of π' by E'_i . Moreover, let $\tilde{C} = \text{Bl}_{z_i, z_j, z_k}(C)$ be the strict transform of C under π and let $\widetilde{T(C)} = \text{Bl}_{z'_i, z'_j, z'_k}(T(C))$ be the strict transform of $T(C)$ under π' . Then of course $\tilde{C} = \widetilde{T(C)}$, and $T(C)$, being the projection $\pi'(\tilde{C})$ of the strict transform \tilde{C} of the irreducible curve C , is of course an irreducible curve. Note that the condition $d > m_i + m_j + m_k$ ensures that \tilde{C} is not one of the curves which are contracted. It thus suffices to verify

$$\deg(T(C)) = d + a,$$

and

$$m'_i = \text{mult}_{z'_i}(T(C)) = \begin{cases} m_\nu, & \text{if } \nu \neq i, j, k, \\ m_\nu + a, & \text{else.} \end{cases}$$

Since outside the lines $\overline{z_i z_j}$, $\overline{z_i z_k}$, and $\overline{z_k z_j}$ the transformation T is an isomorphism and since by hypothesis none of the remaining z_ν belongs to one of

these lines we clearly have $m'_\nu = m_\nu$ for $\nu \neq i, j, k$. Moreover, we have

$$\begin{aligned} m'_i &= \widetilde{T(C)}.E'_i = \widetilde{C}.Bl_{z_i, z_j, z_k}(\overline{z_j z_k}) \\ &= (\pi^*C - \sum_{\nu=i, j, k} m_\nu E_\nu).(\pi^*\overline{z_j z_k} - E_j - E_k) \\ &= C.\overline{z_j z_k} - m_j - m_k = d - m_j - m_k = m_i + a. \end{aligned}$$

Analogously for m'_j and m'_k .

Finally we find

$$\begin{aligned} \deg(T(C)) &= T(C).\overline{z'_i z'_j} = \pi'^*T(C).\pi'^*\overline{z'_i z'_j} \\ &= (\widetilde{T(C)} + \sum_{\nu=i, j, k} m'_\nu E'_\nu).(E_k + E'_i + E'_j) \\ &= \widetilde{C}.E_k + \sum_{\nu=i, j, k} m'_\nu E'_\nu.E_k \\ &= m_k + m'_i + m'_j = d + a. \end{aligned}$$

This proves the claim.

Let us now show by induction that for any $d > 0$ there is an irreducible curve C of degree $d' \geq d$ satisfying (A.1) and (A.2). For $d = 1$ the line $C = \overline{z_1 z_2}$ through z_1 and z_2 gives the induction start. Given some suitable curve of degree $d' \geq d$ the above claim then ensures that through points in very general position there is an irreducible curve of higher degree satisfying (A.1) and (A.2), since $a = d - (m_1 + m_2 + m_3) > 0$. Thus the induction step is done. \square

The example shows that a smooth projective surface Σ may indeed carry an infinite number of exceptional curves - even of the first kind. According to Nagata ([Nag60] Theorem 4a, p. 283) the example is due to Kodaira. For further references on the example see [Har77] Ex. V.4.15, [BeS95] Example 4.2.7, or [Fra41]. [IsS96] p. 198 Example 3 shows that also \mathbb{P}_c^2 blown up in the nine intersection points of two plane cubics carries infinitely many exceptional curves of the first kind.

B. Curves of Self-Intersection Zero

B.1 Proposition

Suppose that $B \subset \Sigma$ is an irreducible curve with $B^2 = 0$ and $\dim |B|_a \geq 1$, then

(B.1) $|B|_a$ is an irreducible reduced projective curve, and

(B.2) there is a fibration $f : \Sigma \rightarrow H$ whose fibres are just the elements of $|B|_a$, and H is the normalisation of $|B|_a$.

We are proving the proposition in several steps.

B.2 Proposition

Let $f : Y' \rightarrow Y$ be a finite flat morphism of noetherian schemes with Y irreducible such that for some point $y_0 \in Y$ the fibre $Y'_{y_0} = f^{-1}(y_0) = Y' \times_Y \text{Spec}(k(y_0))$ is a single reduced point.¹⁵

Then the structure map $f^\# : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_{Y'}$ is an isomorphism, and hence so is f .

Proof: Since there is at least one connected reduced fibre Y'_{y_0} , semicontinuity of flat, proper¹⁶ morphisms in the version [GrD67] IV.12.2.4 (vi) implies that there is an open dense subset $U \subseteq Y$ such that Y'_y is connected and reduced, hence a single reduced point, $\forall y \in U$. (U dense in Y is due to the fact that Y is irreducible.)

Thus the assumptions are stable under restriction to open subschemes of Y , and since the claim that we have to show is local on Y , we may assume that $Y = \text{Spec}(A)$ is affine. Moreover, f being finite, thus affine, we have $Y' = \text{Spec}(B)$ is also affine.

Since f is flat it is open and hence dominates the irreducible affine variety Y and, therefore, induces an inclusion of rings $A \hookrightarrow B$. It now suffices to show:

Claim: $A \hookrightarrow B$ is an isomorphism.

By assumption there exists a $y = P \in \text{Spec}(A) = Y$ such that $Y'_y = f^{-1}(y) = \text{Spec}(B_P/PB_P)$ is a single point with reduced structure. In particular we have for the multiplicity of $Y'_y = \text{Spec}(B_P/PB_P)$ over $\{y\} = \text{Spec}(A_P/PA_P)$

$$1 = \mu(Y'_y) = \text{length}_{A_P/PA_P}(B_P/PB_P),$$

(cf. [Har77] p. 51 for the definition of the multiplicity) which implies that

$$A_P/PA_P \hookrightarrow B_P/PB_P$$

is an isomorphism. Hence by Nakayama's Lemma also

$$A_P \hookrightarrow B_P$$

is an isomorphism, that is, B_P is free of rank 1 over A_P . B being locally free¹⁷ over A , with $A/\sqrt{0}$ an integral domain, thus fulfils

$$A_Q \hookrightarrow B_Q$$

is an isomorphism for all $Q \in \text{Spec}(A)$, and hence the claim follows (cf. [AtM69] Proposition 3.9). \square

B.3 Remark

Some comments from Bernd Kreuzler.

¹⁵The assumption "reduced" is necessary, since finite flat morphisms may very well have only non-reduced fibres. Consider the ring homomorphism $\varphi : A = \mathbb{C}[x] \rightarrow \mathbb{C}[x, y]/(y^2) = B : x \mapsto x$, making $B \cong A \oplus y \cdot A$ into a free and hence flat A -module with only non-reduced fibres (= double-points).

¹⁶By [Har77] Ex. II.4.1 f is proper, since f is finite.

¹⁷Since $A \hookrightarrow B$ is flat!

- (a) Let Y' be the standard example of a non-separated variety, the affine line over k with the origin doubled, and let $f : Y' \rightarrow Y = \mathbb{A}_k^1$ be the projection to Y . Then f is quasi-finite, but NOT finite, since the preimage of the affine Y is not affine. However, $f_*\mathcal{O}_{Y'} = \mathcal{O}_Y$ (see [Har77] II.2.3.5/6), but f is certainly not an isomorphism.
- (b) By [Har77] III.11.3 for $f : Y' \rightarrow Y$ projective between noetherian schemes the condition $f_*\mathcal{O}_{Y'} = \mathcal{O}_Y$ implies that the fibres of f are connected. The converse is also true, if the fibres are connected, the $f_*\mathcal{O}_{Y'} = \mathcal{O}_Y$.
- (c) See (a) for an example, that the condition “projective” for the morphism is not obsolete.
- (d) Let $f : Y' \rightarrow Y$ be the blow up of a point of \mathbb{P}_k^2 , then the fibres are connected, and hence $f_*\mathcal{O}_{Y'} = \mathcal{O}_Y$. f is generically finite, but not finite and certainly no isomorphism.
- (e) If $f : X \rightarrow \text{Spec}(A)$ is a morphism such that $f^\# : A \rightarrow \Gamma(X, \mathcal{O}_X)$ is an isomorphism, then f need not be an isomorphism - despite the fact that $\text{Hom}(X, \text{Spec}(A))$ and $\text{Hom}(A, \Gamma(X, \mathcal{O}_X))$ are naturally bijective. E. g. \mathbb{P}_k^1 and f the constant map to $\text{Spec}(k)$. Then $\Gamma(X, \mathcal{O}_X) = k$ and thus $f^\# = \text{id}$ is an isomorphism, but f is certainly not! If X is also affine, then however f is an isomorphism if and only if $f^\#$ is so.
- (f) If $A \rightarrow B$ is a ring homomorphism and B is locally free of rank 1 as an A -module, and if for some $P \in \text{Spec}(A)$ the localisation $A_P \rightarrow B_P$ is an isomorphism, then the homomorphism itself is an isomorphism. If, however, $A \rightarrow B$ is only an A -module homomorphism, this need not be true. E. g. let X be a smooth curve, e. g. \mathbb{P}^1 , and $x \in X$ a point; let $\mathcal{J}_{x/X} := \mathcal{O}_X(-x)$ be the ideal sheaf of x , in the case of \mathbb{P}^1 it's just $\mathcal{O}_{\mathbb{P}^1}(-1)$. This is a locally free sheaf, and x is a divisor on the smooth curve X . This gives an exact sequence $0 \rightarrow \mathcal{J}_{x/X} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_x \rightarrow 0$. For the stalks over $y \neq x$ the map from $L \rightarrow \mathcal{O}_X$ is of course an isomorphism, but not for $y = x$.
- (g) Let $T = \mathbb{A}_k^1$, $X_t = V(x^2 - txy) \subset \mathbb{P}_k^1$ with homogeneous coordinates x, y on \mathbb{P}_k^1 . Then for $t_0 = 0$ the fibre X_{t_0} is connected, namely a non-reduced fat point, while for $t \neq 0$ the fibre X_t consists of two single points, and is thus not connected. The condition X_{t_0} reduced in the Corollary B.5 is hence vital.

B.4 Proposition (Principle of Connectedness)

Let X and Y be noetherian schemes, Y connected, and let $\pi : X \rightarrow Y$ be a flat projective morphism such that for some $y_0 \in Y$ the fibre $X_{y_0} = \pi^{-1}(y_0)$ is reduced and connected.

Then for all $y \in Y$ the fibre $X_y = \pi^{-1}(y)$ is connected.

Proof: Stein Factorisation (cf. [GrD67] III.4.3.3 or [Har77] III.11.5) gives a factorisation of π of the form

$$\pi : X \xrightarrow{\pi'} Y' = \text{Spec}(\pi_* \mathcal{O}_X) \xrightarrow{f} Y,$$

with

- (1) π' connected (i. e. its fibres are connected),
- (2) f finite,
- (3) $f_* \mathcal{O}_{Y'} = \pi_* \mathcal{O}_X$ locally free over \mathcal{O}_Y , since π is flat, and
- (4) $Y'_{y_0} = f^{-1}(y_0)$ is connected and reduced, i. e. a single reduced point.

Because of (1) it suffices to show that f is connected; and we claim, moreover, that the fibres of f are reduced as well.

Considering points in the intersections of the finite number of irreducible components of Y we can reduce to the case Y irreducible. Since f is finite (3) is equivalent to saying that f is flat (cf. [AtM69] Proposition 3.10). Hence f fulfils the assumptions of Proposition B.2, and we conclude that $\mathcal{O}_Y = f_* \mathcal{O}_{Y'}$ and the proposition follows from [Har77] III.11.3.¹⁸

Alternatively, from [GrD67] IV.15.5.9 (ii) it follows that there is an open dense subset $U \subseteq Y$ such that X_y is connected for all $y \in U$. Since, moreover, by the same theorem the number of connected components of the fibres is a lower semi-continuous function on Y the special fibres cannot have more connected components than the general ones, that is, all fibres are connected. \square

B.5 Corollary (Principle of Connectedness)

Let $\{X_t\}_{t \in T}$ be a flat family of closed subschemes $X_t \subseteq \mathbb{P}_{k(t)}^n$, where T is a connected noetherian scheme. Suppose that X_{t_0} is connected and reduced for some point $t_0 \in T$. Then X_t is connected for all $t \in T$.

Proof: The X_t are the fibres of a flat projective morphism

$$\begin{array}{ccc} \mathcal{X} & \hookrightarrow & \mathbb{P}_T^n \\ & \searrow \pi & \downarrow \\ & & T, \end{array}$$

where \mathcal{X} is a closed subscheme of \mathbb{P}_T^n . The result thus follows from Proposition B.4.¹⁹ \square

B.6 Lemma

Under the hypotheses of Proposition B.1 let $C \in |\mathcal{B}|_a$ then C is connected.

¹⁸Compare the result with [GrD67] III.4.3.10, which deals with the case of X also being integral, but weakens the hypothesis on π to proper and universally open. Also the assumption on the fibre is reduced to $R(Y)$ being algebraically closed in $R(X)$ - in the case where the characteristic of $R(Y)$ is zero. Compare also [GrD67] IV.12.2.4 (vi) and III.15.5.9, which both deal with an arbitrary number of connected components.

¹⁹Compare the result with [Har77] Ex. III.11.4.

Proof: Consider the universal family

$$\begin{array}{ccc}
 |B|_a \times \Sigma & \longleftarrow & \bigcup_{C \in |B|_a} \{C\} \times C =: S \\
 & \searrow \text{pr}_{|B|_a} & \downarrow \pi \text{ flat} \\
 & & |B|_a
 \end{array} \tag{B.3}$$

over the connected projective scheme $|B|_a \subseteq \text{Hilb}_\Sigma$.

Since the projection π is a flat projective morphism, and since the fibre $\pi^{-1}(B) = \{B\} \times B$ is connected and reduced, the result follows from Proposition B.4. \square

B.7 Lemma

Under the hypotheses of Proposition B.1 let $C \in |B|_a$ with $B \subseteq C$, then $C = B$.

Proof: Suppose $B \subsetneq C$, then the Hilbert polynomials of B and C are different in contradiction to $B \sim_a C$. \square

B.8 Lemma

Under the hypotheses of Proposition B.1 let $C \in |B|_a$ with $C \neq B$, then $C \cap B = \emptyset$.

Proof: Since B is irreducible by Lemma B.7 B and C do not have a common component. Suppose $B \cap C = \{p_1, \dots, p_r\}$, then $B^2 = B.C \geq r > 0$ in contradiction to $B^2 = 0$. \square

B.9 Proposition (Zariski's Lemma)

Under the hypotheses of Proposition B.1 let $C = \sum_{i=1}^r n_i C_i \in |B|_a$, where the C_i are pairwise different irreducible curves and $n_i > 0$ for $i = 1, \dots, r$.

Then the intersection matrix $Q = (C_i.C_j)_{i,j=1,\dots,r}$ is negative semi-definite, and, moreover, C , considered as an element of the vectorspace $\bigoplus_{i=1}^r \mathbb{Q} \cdot C_i$, generates the annihilator of Q .

In particular, $D^2 \leq 0$ for all curves $D \subseteq C$, and, moreover, $D^2 = 0$ if and only if $D = C$.

Proof: By Lemma B.6 C is connected. We are going to apply [BPV84] I.2.10, and thus we have to verify three conditions.

- (a') $C.C_i = B.C_i = 0$ for all $i = 1, \dots, r$ by Lemma B.8. Thus C is an element of the annihilator of Q with $n_i > 0$ for all $i = 1, \dots, r$.
- (b) $C_i.C_j \geq 0$ for all $i \neq j$.
- (c) Since C is connected there is no non-trivial partition $I \cup J$ of $\{1, \dots, r\}$ such that $C_i.C_j = 0$ for all $i \in I$ and $j \in J$.

Thus [BPV84] I.2.10 implies that $-Q$ is positive semi-definite. \square

B.10 Lemma

Under the hypotheses of Proposition B.1 let $C, C' \in |B|_a$ be two distinct curves, then $C \cap C' = \emptyset$.

Proof: Suppose $C = A + D$ and $C' = A + D'$ such that D and D' have no common component.

We have

$$0 = B^2 = (A + D)^2 = (A + D')^2 = (A + D).(A + D'),$$

and thus

$$(A + D)^2 + (A + D')^2 = 2(A + D).(A + D'),$$

which implies that

$$D^2 + D'^2 = 2D.D',$$

where each summand on the left hand side is less than or equal to zero by Proposition B.9, and the right hand side is greater than or equal to zero, since the curves D and D' have no common component. We thus conclude

$$D^2 = D'^2 = D.D' = 0.$$

But then again Proposition B.9 implies that $D = C$ and $D' = C'$, that is, C and C' have no common component.

Suppose $C \cap C' = \{p_1, \dots, p_r\}$, then $B^2 = C.C' \geq r > 0$ would be a contradiction to $B^2 = 0$. Hence, $C \cap C' = \emptyset$. \square

B.11 Lemma

Under the hypotheses of Proposition B.1 consider once more the universal family (B.3) together with its projection onto Σ ,

$$\begin{array}{ccc}
 |B|_a \times \Sigma & \xrightarrow{\text{pr}_\Sigma} & \Sigma \\
 \swarrow & \searrow \pi' & \\
 S & & \\
 \downarrow \pi & & \\
 |B|_a & &
 \end{array}
 \quad (\text{B.4})$$

Then S is an irreducible projective surface, $|B|_a$ is an irreducible curve, and π' is surjective.

Proof:

Step 1: S is an irreducible projective surface and π' is surjective.

The universal property of $|B|_a$ implies that S is an effective Cartier divisor of $|B|_a \times \Sigma$, and thus in particular projective of dimension at least $2 \leq \dim |B|_a + \dim(\Sigma) - 1$. Since π' is projective, its image is closed in Σ and of dimension 2, hence it is the whole of Σ , since Σ is irreducible.

By Lemma B.10 the fibres of π' are all single points, and thus, by [Har92] Theorem 11.14, S is irreducible. Moreover, $\dim(S) = \dim(\Sigma) + \dim(\text{fibre}) = 2$.

Step 2: $\dim |B|_a = \dim(|B|_a \times \Sigma) - \dim(\Sigma) = \dim(S) + 1 - 2 = 1$.

Step 3: $|B|_a$ is irreducible.

Let V be any irreducible component of $|B|_a$ of dimension one, then we have a universal family over V and the analogue of Step 1 for V shows that the curves in V cover Σ . But then by Lemma B.10 there can be no further curve in $|B|_a$, since any further curve would necessarily have a non-empty intersection with one of the curves in V . \square

B.12 Lemma

Let's consider the following commutative diagram of projective morphisms

$$\begin{array}{ccc}
 S & \xrightarrow{\pi'} & \Sigma \\
 \pi \downarrow & \swarrow & \uparrow \varphi' \\
 |B|_a & \xleftarrow{\varphi} & S_{\text{red}}
 \end{array}
 \tag{B.5}$$

The map $\varphi' : S_{\text{red}} \rightarrow \Sigma$ is birational.²⁰

Proof: Since S_{red} and Σ are irreducible and reduced, and since φ' is surjective, we may apply [Har77] III.10.5, and thus there is an open dense subset $U \subseteq S_{\text{red}}$ such that $\varphi'_| : U \rightarrow \Sigma$ is smooth. Hence, in particular $\varphi'_|$ is flat and the fibres are single reduced points.²¹ Since $\varphi'_| : U \rightarrow \varphi'(U)$ is projective and quasi-finite, it is finite (cf. [Har77] Ex. III.11.2), and it follows from Proposition B.2 that $\varphi'_|$ is an isomorphism onto its image, i. e. φ' is birational. \square

B.13 Lemma

If $\psi : \Sigma \dashrightarrow S_{\text{red}}$ denotes the rational inverse of the map φ' in (B.5), then ψ is indeed a morphism, i. e. φ' is an isomorphism.

Proof: By Lemma B.10 the fibres of φ' over the possible points of indeterminacy of φ' are just points, and thus the result follows from [Bea83] Lemma II.9. \square

B.14 Lemma

The map $g : \Sigma \rightarrow |B|_a$ assigning to each point $p \in \Sigma$ the unique curve $C \in |B|_a$ with $p \in C$ is a morphism, and is thus a fibration whose fibres are the curves in $|B|_a$.

Proof: We just have $g = \varphi \circ \psi$. \square

Proof of Proposition B.1: Let $\nu : H \rightarrow |B|_a$ be the normalisation of the irreducible curve $|B|_a$. Then H is a smooth irreducible curve.

²⁰The proof uses that the characteristic of the ground field is zero, even though one might perhaps avoid this.

²¹By definition $\varphi'_|$ is étale, and hence the completed local rings of the fibres are isomorphic to the completed local rings of their base points and hence regular. But then the local rings themselves are regular and thus reduced.

Moreover, since Σ is irreducible and smooth, and since $g : \Sigma \rightarrow |B|_a$ is surjective, g factorises over H , i. e. we have the following commutative diagram

$$\begin{array}{ccc} \Sigma & \xrightarrow{g} & |B|_a \\ \exists f \downarrow & \nearrow v & \\ H & & \end{array}$$

Then f is the desired fibration. □

C. Some Facts used for the Proof of the Lemma of Geng Xu

In this section we are, in particular, writing down some identifications of certain sheaves respectively of their global sections. Doing this we try to be very formal. However, in a situation of the kind $X \xrightarrow{i} Y \xrightarrow{\pi} Z$ we do not distinguish between \mathcal{O}_X and $i_*\mathcal{O}_X$, or between π and the restriction of π to X .

C.1 Lemma

Let $\varphi(x, y, t) = \sum_{i=0}^{\infty} \varphi_i(x, y) \cdot t^i \in \mathbb{C}\{x, y, t\}$ with $\varphi(x, y, t) \in (x, y)^m$ for every fixed t in some small disc Δ around 0. Then $\varphi_i(x, y) \in (x, y)^m$ for every $i \in \mathbb{N}_0$.

Proof: We write the power series as $\varphi = \sum_{\alpha+\beta=0}^{\infty} \left(\sum_{i=0}^{\infty} c_{\alpha,\beta,i} \cdot t^i \right) \cdot x^\alpha y^\beta$. $\varphi(x, y, t) \in (x, y)^m$ for every $t \in \Delta$ implies

$$\sum_{i=0}^{\infty} c_{\alpha,\beta,i} \cdot t^i = 0 \quad \forall \alpha + \beta < m \text{ and } t \in \Delta.$$

The identity theorem for power series in \mathbb{C} then implies that

$$c_{\alpha,\beta,i} = 0 \quad \forall \alpha + \beta < m \text{ and } i \geq 0.$$

□

C.2 Lemma

Let X be a noetherian scheme, $i : C \hookrightarrow X$ a closed subscheme, \mathcal{F} a sheaf of modules on C , and \mathcal{G} a sheaf of modules on X . Then

$$(C.1) \quad i_*\mathcal{F} \cong i_*\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_C,$$

$$(C.2) \quad H^0(C, \mathcal{F}) = H^0(X, i_*\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_C),$$

$$(C.3) \quad \mathcal{G} \otimes_{\mathcal{O}_X} \mathcal{O}_C \cong i_*i^*(\mathcal{G} \otimes_{\mathcal{O}_X} \mathcal{O}_C), \text{ and}$$

$$(C.4) \quad H^0(C, i^*(\mathcal{G} \otimes_{\mathcal{O}_X} \mathcal{O}_C)) = H^0(X, \mathcal{G} \otimes_{\mathcal{O}_X} \mathcal{O}_C).$$

Proof:

(C.1) For $U \subseteq X$ open, we define

$$\begin{array}{ccc} \Gamma(U, i_*\mathcal{F}) & \rightarrow & \Gamma(U, i_*\mathcal{F}) \otimes_{\Gamma(U, \mathcal{O}_X)} \Gamma(U, \mathcal{O}_C) \subseteq \Gamma(U, i_*\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_C) \\ s & \mapsto & s \otimes 1. \end{array}$$

This morphism induces on the stalks the isomorphism

$$i_*\mathcal{F}_x = \left\{ \begin{array}{ll} \mathcal{F}_x, \text{ (if } x \in C) & = \mathcal{F}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{X,x}/I_{C,x} \\ 0, \text{ (else)} & = 0 \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{X,x}/I_{C,x} \end{array} \right\} \cong i_*\mathcal{F}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{C,x},$$

where $I_{C,x}$ is the ideal defining C in X locally at x .

(C.2) The identification (C.1) together with [Har77] III.2.10 gives:

$$H^0(C, \mathcal{F}) = H^0(X, i_*\mathcal{F}) = H^0(X, i_*\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_C).$$

(C.3) The adjoint property of i_* and i^* together with $i^*i_* \cong \text{id}$ gives rise to the following isomorphisms:

$$\begin{aligned} \text{End}(i_*i^*(\mathcal{G} \otimes \mathcal{O}_C)) &\cong \text{Hom}(i^*i_*i^*(\mathcal{G} \otimes \mathcal{O}_C), i^*(\mathcal{G} \otimes \mathcal{O}_C)) \\ &\cong \text{End}(i^*(\mathcal{G} \otimes \mathcal{O}_C)) \cong \text{Hom}(\mathcal{G} \otimes \mathcal{O}_C, i_*i^*(\mathcal{G} \otimes \mathcal{O}_C)). \end{aligned}$$

That means, that the identity morphism on $i_*i^*(\mathcal{G} \otimes \mathcal{O}_C)$ must correspond to an isomorphism from $\mathcal{G} \otimes \mathcal{O}_C$ to $i_*i^*(\mathcal{G} \otimes \mathcal{O}_C)$ via these identifications.

(C.4) follows from (C.3) and once more [Har77] III.2.10.

□

C.3 Corollary

In the situation of Lemma II.2.4 we have:

$$(C.5) \quad H^0(C, \pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)) = H^0(\Sigma, \pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)), \text{ and}$$

$$(C.6) \quad H^0(C, \pi_*\mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)) = H^0(\Sigma, \pi_*\mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)).$$

Proof: We denote by $j : \tilde{C} \hookrightarrow \tilde{\Sigma}$ and $i : C \hookrightarrow \Sigma$ respectively the given embeddings.

(C.5) By (C.2) in Lemma C.2 we have:

$$H^0(C, \pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)) = H^0(\Sigma, i_*(\pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_C} \mathcal{O}_C(C)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C).$$

By the projection formula this is just equal to:

$$\begin{aligned} H^0(\Sigma, (i_*\pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_\Sigma(C)) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C) &= H^0(\Sigma, \pi_*j_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)) \\ &=_{\text{def}} H^0(\Sigma, \pi_*\mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)). \end{aligned}$$

(C.6) Using (C.4) in Lemma C.2 we get:

$$\begin{aligned} H^0(C, \pi_*\mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)) &=_{\text{def}} H^0(C, i^*(\pi_*\mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C))) = \\ &= H^0(\Sigma, \pi_*\mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_C(C)). \end{aligned}$$

□

C.4 Lemma

Let \mathcal{F} be any coherent sheaf on Σ . Then the kernel of the natural map

$$\bigotimes_{i=1}^r \mathcal{J}_{X(m_i; z_i)/\Sigma} \otimes \mathcal{F} \xrightarrow{\delta} \mathcal{J}_{X(\underline{m}; \underline{z})/\Sigma} \cdot \mathcal{F}$$

has support contained in $\{z_1, \dots, z_r\}$.

Moreover, if \mathcal{F} is locally free, then δ is an isomorphism.

Proof: We have $\text{Ker}(\delta)_z = \text{Ker}(\delta_z)$ and for $z \notin \{z_1, \dots, z_r\}$ the map δ_z is given by

$$\begin{aligned} \bigotimes_{i=1}^r \mathcal{O}_{\Sigma, z} \otimes \mathcal{F}_z &\longrightarrow \mathcal{O}_{\Sigma, z} \cdot \mathcal{F}_z \\ f_1 \otimes \dots \otimes f_r \otimes g &\longmapsto f_1 \cdots f_r \cdot g, \end{aligned}$$

and is thus an isomorphism.

Suppose now that \mathcal{F} is locally free and that $z \in \{z_1, \dots, z_r\}$. Then δ_z is given by

$$\begin{aligned} \mathcal{O}_{\Sigma, z_i} \otimes \dots \otimes \mathfrak{m}_{\Sigma, z_i}^{m_i} \otimes \dots \otimes \mathcal{O}_{\Sigma, z_i} \otimes \mathcal{O}_{\Sigma, z_i} &\longrightarrow \mathfrak{m}_{\Sigma, z_i}^{m_i} \\ f_1 \otimes \dots \otimes f_r \otimes g &\longmapsto f_1 \cdots f_r \cdot g, \end{aligned}$$

which again is an isomorphism. \square

C.5 Lemma

With the notation of Lemma II.2.4 we show that $\text{supp}(\text{Ker}(\gamma)) \subseteq \{z_1, \dots, z_r\}$.

Proof: Since $\pi : \tilde{\Sigma} \setminus (\bigcup_{i=1}^r E_i) \longrightarrow \Sigma \setminus \{z_1, \dots, z_r\}$ is an isomorphism, we have for any sheaf \mathcal{F} of $\mathcal{O}_{\tilde{\Sigma}}$ -modules and $y \in \tilde{\Sigma} \setminus (\bigcup_{i=1}^r E_i)$:

$$(\pi_* \mathcal{F})_{\pi(y)} = \lim_{\pi(y) \in V} \mathcal{F}(\pi^{-1}(V)) = \lim_{y \in U} \mathcal{F}(U) = \mathcal{F}_y.$$

In particular,

$$(\pi_* \mathcal{O}_{\tilde{\Sigma}}(E) \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C))_{\pi(y)} \cong \mathcal{O}_{\tilde{\Sigma}, y} \otimes_{\mathcal{O}_{\Sigma, \pi(y)}} \mathcal{O}_{C, \pi(y)} \cong \mathcal{O}_{\Sigma, \pi(y)} \otimes_{\mathcal{O}_{\Sigma, \pi(y)}} \mathcal{O}_{C, \pi(y)},$$

and

$$(\pi_* \mathcal{O}_{\tilde{C}}(E) \otimes_{\mathcal{O}_{\Sigma}} \mathcal{O}_C(C))_{\pi(y)} \cong \mathcal{O}_{\tilde{C}, y} \otimes_{\mathcal{O}_{\Sigma, \pi(y)}} \mathcal{O}_{C, \pi(y)} \cong \mathcal{O}_{C, \pi(y)} \otimes_{\mathcal{O}_{\Sigma, \pi(y)}} \mathcal{O}_{C, \pi(y)}.$$

Moreover, the morphism $\gamma_{\pi(y)}$ becomes under these identifications just the morphism given by $a \otimes \bar{b} = 1 \otimes \overline{ab} \mapsto \bar{a} \otimes \bar{b} = 1 \otimes \overline{ab}$, which is injective. Thus, $0 = \text{Ker}(\gamma_{\pi(y)}) = \text{Ker}(\gamma)_{\pi(y)}$, and $\pi(y) \notin \text{supp}(\text{Ker}(\gamma))$. \square

C.6 Lemma

Let X be an irreducible noetherian scheme, \mathcal{F} a coherent sheaf on X , and $s \in H^0(X, \mathcal{F})$ such that $\dim(\text{supp}(s)) < \dim(X)$. Then $s \in H^0(X, \text{Tor}(\mathcal{F}))$.

Proof: The multiplication by s gives rise to the following exact sequence:

$$0 \longrightarrow \text{Ker}(\cdot s) \longrightarrow \mathcal{O}_X \xrightarrow{\cdot s} \mathcal{F}.$$

Since \mathcal{O}_X and \mathcal{F} are coherent, so is $\text{Ker}(\cdot s)$, and hence $\text{supp}(\text{Ker}(\cdot s))$ is closed in X . Now,

$$\begin{aligned} \text{supp}(\text{Ker}(\cdot s)) &= \{z \in X \mid \exists 0 \neq r_z \in \mathcal{O}_{X,z} : r_z \cdot s_z = 0\} \\ &= \{z \in X \mid s_z \in \text{Tor}(\mathcal{F}_z)\}. \end{aligned}$$

But then the complement $\{z \in X \mid s_z \notin \text{Tor}(\mathcal{F}_z)\}$ is open and is contained in $\text{supp}(s)$ (since $s_z = 0$ implies that $s_z \in \text{Tor}(\mathcal{F}_z)$), and is thus empty since X is irreducible and $\text{supp}(s)$ of lower dimension. \square

D. Some Facts on Divisors on Curves

D.1 Remark

Let $C = C_1 \cup \dots \cup C_k$ be a reduced curve on a smooth projective surface Σ over \mathbb{C} , where the C_i are irreducible, and let \mathcal{L} be a line bundle on C . Then we define the *degree* of \mathcal{L} with the aid of the normalisation $\nu : C' \rightarrow C$. We have $H^2(C, \mathbb{Z}) \cong \bigoplus_{i=1}^k H^2(C'_i, \mathbb{Z}) = \mathbb{Z}^k$, and thus the image of \mathcal{L} in $H^2(C, \mathbb{Z})$, which is the first Chern class of \mathcal{L} , can be viewed as a vector (l_1, \dots, l_k) of integers, and we may define the degree of \mathcal{L} by

$$\text{deg}(\mathcal{L}) := l_1 + \dots + l_k.$$

In particular, if C is irreducible, we get:

$$\text{deg}(\mathcal{L}) = \text{deg}(\nu^* \mathcal{L}) = c_1(\nu^* \mathcal{L}).$$

Since $H^0(C, \mathcal{L}) \neq 0$ implies that $H^0(C', \nu^* \mathcal{L}) \neq 0$, and since the existence of a non-vanishing global section of $\nu^* \mathcal{L}$ on the smooth curve C' implies that the corresponding divisor is effective, we get the following lemma. (cf. [BPV84] Section II.2)

D.2 Lemma

Let C be an irreducible reduced curve on a smooth projective surface Σ , and let \mathcal{L} be a line bundle on C . If $H^0(C, \mathcal{L}) \neq 0$, then $\text{deg}(\mathcal{L}) \geq 0$.

When studying divisors on geometrically ruled surfaces or on products of curves, we need the following estimation of the number of independent sections of a divisor on a curve.

D.3 Lemma

Let C be a smooth projective curve over \mathbb{C} of genus g and let $\mathfrak{c} \in \text{Div}(C)$ with $\text{deg}(\mathfrak{c}) \geq 0$. Then $h^0(C, \mathfrak{c}) \leq \text{deg}(\mathfrak{c}) + 1$.

Proof: Let us suppose that \mathfrak{c} is effective, since otherwise $h^0(C, \mathfrak{c}) \leq 1$ anyway. Then the inclusion

$$\mathcal{O}_C(K_C - \mathfrak{c}) \xrightarrow{+\mathfrak{c}} \mathcal{O}_C(K_C)$$

implies that $h^0(C, K_C - \mathfrak{c}) \leq h^0(C, K_C) = g$. Hence by the Riemann-Roch Formula we get

$$h^0(C, \mathfrak{c}) = \text{deg}(\mathfrak{c}) + 1 - g + h^0(C, K_C - \mathfrak{c}) \leq \text{deg}(\mathfrak{c}) + 1.$$

□

E. Some Facts on Divisors on Surfaces

In this section we gather some simple facts on divisors on smooth surfaces which are used throughout the paper.

E.1 Lemma

If $\text{NS}(\Sigma) = \mathbb{L} \cdot \mathbb{Z}$, then either \mathbb{L} or $-\mathbb{L}$ is ample.

Proof: Since Σ is projective, there exists some very ample divisor H , and by assumption $H \sim_{\mathbb{Q}} h \cdot \mathbb{L}$ for some $0 \neq h \in \mathbb{Z}$. W. l. o. g. $h > 0$. Thus

$$\mathbb{L}^2 = \frac{1}{h^2} \cdot H^2 > 0,$$

and for any irreducible curve $C \subset \Sigma$

$$\mathbb{L} \cdot C = \frac{1}{h} \cdot H \cdot C > 0.$$

Therefore, by the Nakai-Moishezon Criterion \mathbb{L} is ample. □

E.2 Lemma

Let Σ be any smooth projective surface and let $A, B \in \text{Div}(\Sigma)$ be such that A is effective and $-A^2 > A \cdot B \geq 0$. Then

- (a) $h^0(\Sigma, A + B) = h^0(\Sigma, B)$.
- (b) If $(B - K_{\Sigma}) \cdot A > 0$, then $h^1(\Sigma, A + B) = h^1(\Sigma, B)$.

Proof:

- (a) We note that the divisor $\mathcal{O}_A(A + B)$ has degree $A \cdot (A + B) < 0$ and is thus not effective, i. e. $h^0(A, \mathcal{O}_A(A + B)) = 0$.

Considering the exact sequence

$$0 \longrightarrow \mathcal{O}_{\Sigma}(B) \xrightarrow{+A} \mathcal{O}_{\Sigma}(A + B) \longrightarrow \mathcal{O}_A(A + B) \longrightarrow 0, \quad (\text{E.1})$$

the statement is implied by its long exact cohomology sequence

$$0 \longrightarrow H^0(\Sigma, B) \longrightarrow H^0(\Sigma, A + B) \longrightarrow H^0(A, \mathcal{O}_A(A + B)) = 0.$$

- (b) By assumption $0 > A \cdot (K_{\Sigma} - B) = \deg(\mathcal{O}_A(K_{\Sigma} - B)) = \deg(\mathcal{O}_A(K_A) \otimes \mathcal{O}_A(-A - B))$. Hence

$$0 = h^0(A, \mathcal{O}_A(K_A) \otimes \mathcal{O}_A(-A - B)) = h^1(A, \mathcal{O}_A(A + B)).$$

Hence once more the long exact cohomology sequence of (E.1) finishes the proof:

$$0 \longrightarrow H^1(\Sigma, B) \longrightarrow H^1(\Sigma, A + B) \longrightarrow H^1(A, \mathcal{O}_A(A + B)) = 0.$$

□

E.3 Lemma

Let \mathbb{L} be very ample over \mathbb{C} on the smooth projective surface Σ .

- (a) *Given two distinct points $z, z' \in \Sigma$, there is a smooth connected curve through z and z' in $|L|_1$. Indeed, a generic curve in $|L|_1$ through z and z' will be so.*
- (b) *Given distinct points $z_1, \dots, z_r \in \Sigma$, the set \mathcal{U} of points z_0 such that for any point z^* there is a smooth connected curve through z_0 and z^* containing at most one of the z_j for $j = 1, \dots, r$ is open and dense in Σ .*

Proof: Considering the embedding $\Sigma \subseteq \mathbb{P}_\mathbb{C}^n$ defined by L the curves in $|L|_1$ are in one-to-one correspondence with the hyperplane sections.

- (a) There is an $n - 2$ -dimensional family of hyperplane sections going through two fixed points of Σ , which in local coordinates w. l. o. g. is given by the family of equations $\mathcal{F} = \{a_1x_1 + \dots + a_{n-1}x_{n-1} = 0 \mid (a_1 : \dots : a_{n-1}) \in \mathbb{P}_\mathbb{C}^{n-2}\}$. Since the local analytic rings of Σ in every point are smooth, hence, in particular complete intersections, they are given as $\mathbb{C}\{x_1, \dots, x_n\}$ modulo some ideal generated by $n - 2$ power series f_1, \dots, f_{n-2} forming a regular sequence. Thus, having $n - 2$ free indeterminates in our family \mathcal{F} of equations, a generic equation g will lead to a regular sequence f_1, \dots, f_{n-2}, g , i. e. the hyperplane section defined by g is smooth in each of the two points, and thus everywhere. Moreover, by [Har77] III.7.9 it is connected.²²
- (b) Given $i, j \in \{1, \dots, r\}$, $i \neq j$, there is a unique line L_{ij} in $\mathbb{P}_\mathbb{C}^n$ through z_i and z_j . We set $\mathcal{U}_{ij} = \Sigma \setminus L_{ij}$, which is an open and dense subset of Σ . Let $z_0 \in \mathcal{U}_{ij}$ and $z^* \in \Sigma$ be given. W. l. o. g. $z^* \neq z_0$. Since $z_0 \notin L_{ij}$ a generic hyperplane in $\mathbb{P}_\mathbb{C}^n$ through z_0 and z^* does not contain L_{ij} , and thus a generic hyperplane section through z_0 and z^* does not contain both z_i and z_j . However, by part (a) a generic hyperplane section is a smooth connected curve through z_0 and z^* . Setting $\mathcal{U} = \bigcap_{i \neq j} \mathcal{U}_{ij}$, which is open and dense in Σ , we are done.

□

We reformulate the Hodge Index Theorem from [BPV84] in the following well known equivalent form which suits our purposes better.

E.4 Theorem (Hodge Index Theorem)

Let $H, D \in \text{Div}(\Sigma) \otimes \mathbb{Q}$ be \mathbb{Q} -divisors with $H^2 > 0$. Then $H^2 \cdot D^2 \leq (H \cdot D)^2$ with equality if and only if $D \sim_n \alpha H$ for some $\alpha \in \mathbb{Q}$.

Proof: We set $\alpha = -\frac{H \cdot D}{H^2} \in \mathbb{Q}$. Then $H \cdot (D + \alpha H) = 0$ and by the standard Hodge Index Theorem (cf. [BPV84] Corollary 2.15) we have

$$0 \geq (D + \alpha H)^2 = H^2 \cdot \alpha^2 + 2 \cdot H \cdot D \cdot \alpha + D^2.$$

²²See also [Har77] Proof of Corollary III.9.13.

We define a polynomial function

$$\varphi : \mathbb{R} \rightarrow \mathbb{R} : t \mapsto H^2 \cdot t^2 + 2 \cdot H \cdot D \cdot t + D^2.$$

Since the leading term H^2 is positive, there is some value $\beta \in \mathbb{R}$ such that $\varphi(\beta) > 0$. However, $\varphi(\alpha) \leq 0$, and thus by the Intermediate Value Theorem φ has at least one zero. This, in turn, implies that the discriminant of the polynomial φ is non-negative, i. e.

$$4 \cdot (D \cdot H)^2 - 4 \cdot D^2 \cdot H^2 \geq 0.$$

Moreover, equality holds precisely if φ has just one zero and is thus always non-negative, in particular we must have $0 = \varphi(\alpha) = H \cdot (D + \alpha H)$. But then by the standard Hodge Index Theorem we get $D + \alpha H \sim_n 0$, i. e. $D \sim_n -\alpha H$. \square

Finally we state the Theorem of Bertini in a version which is suitable for our purposes and which can be found in [Wae73] § 47, Satz 3 and Satz 4.

E.5 Theorem (Theorem of Bertini)

Let Σ be a smooth projective surface, $D \in \text{Div}(\Sigma)$ a divisor and $\mathcal{D} \subseteq |D|_1$ a linear system without fixed part. Suppose that every curve in \mathcal{D} is reducible.

- (a) Two generic curves in \mathcal{D} intersect only in the base points of \mathcal{D} .
- (b) There exists a one-dimensional algebraic family \mathcal{F} of curves in Σ such that the irreducible components of any curve in \mathcal{D} belong to \mathcal{F} .

F. The Hilbert Scheme Hilb_Σ^n

Hilb_Σ^n denotes the Hilbert scheme of zero-dimensional schemes of degree n .

F.1 Lemma

Let $D \in \text{Div}(\Sigma)$ be a divisor and $X \subset \Sigma$ be a zero-dimensional scheme of degree n with $h^1(\Sigma, \mathcal{J}_{X/\Sigma}(D)) = 0$. Then there is an open, dense, irreducible neighbourhood $\mathcal{U} = \mathcal{U}(X) \subseteq \text{Hilb}_\Sigma^n$ of X such that $h^1(\Sigma, \mathcal{J}_{Y/\Sigma}(D)) = 0$ for any $Y \in \mathcal{U}$.

Proof: We consider the universal family of Hilb_Σ^n :

$$\begin{array}{ccc} \mathcal{X} & \hookrightarrow & \text{Hilb}_\Sigma^n \times \Sigma \xrightarrow{\text{pr}_\Sigma} \Sigma, \\ \pi \downarrow & \swarrow \text{pr} & \\ \text{Hilb}_\Sigma^n & & \end{array} \quad (\text{F.1})$$

where pr and pr_Σ denote the canonical projections. In particular π and pr are flat, that is $\mathcal{O}_{\text{Hilb}_\Sigma^n \times \Sigma}$ and $\mathcal{O}_\mathcal{X}$ are flat $\mathcal{O}_{\text{Hilb}_\Sigma^n}$ -modules via pr and π respectively. From the exact sequence

$$0 \rightarrow \mathcal{J}_\mathcal{X} \rightarrow \mathcal{O}_{\text{Hilb}_\Sigma^n \times \Sigma} \rightarrow \mathcal{O}_\mathcal{X} \rightarrow 0$$

it thus follows that the ideal sheaf $\mathcal{J}_\mathcal{X}$ of \mathcal{X} is flat over $\mathcal{O}_{\text{Hilb}_\Sigma^n}$. Moreover, by Lemma F.3 also the sheaf $\mathcal{J}_\mathcal{X}(D) := \mathcal{J}_\mathcal{X} \otimes_{\mathcal{O}_{\text{Hilb}_\Sigma^n \times \Sigma}} \text{pr}_\Sigma^* \mathcal{O}_\Sigma(D)$ is flat over $\mathcal{O}_{\text{Hilb}_\Sigma^n}$.

The Semicontinuity Theorem (cf. [Har77] III.12.8) therefore implies that the function

$$\alpha : \text{Hilb}_\Sigma^n \rightarrow \mathbb{Z} : Y \mapsto \dim_{k(Y)} H^1(\text{pr}^{-1}(Y), \mathcal{J}_X(D)_Y)$$

is upper semi continuous.

Now, for $Y \in \text{Hilb}_\Sigma^n$ the restriction $\text{pr}_{\Sigma|} : \text{pr}^{-1}(Y) = \{Y\} \times \Sigma \rightarrow \Sigma$ is an isomorphism, and this isomorphism induces

$$Y \cong \{Y\} \times Y = \pi^{-1}(Y) \hookrightarrow \text{pr}^{-1}(Y) = \{Y\} \times \Sigma \cong \Sigma$$

and $\mathcal{J}_{X,Y} \cong \text{pr}_{\Sigma*} \mathcal{J}_{X,Y} = \mathcal{J}_{Y/\Sigma}$. With the aid of the projection formula we deduce

$$(\mathcal{J}_X(D))_Y \cong \text{pr}_{\Sigma*} (\mathcal{J}_X \otimes_{\mathcal{O}_{\text{Hilb}_\Sigma^n \times \Sigma}} \text{pr}_\Sigma^* \mathcal{O}_\Sigma(D)) \cong \text{pr}_{\Sigma*} \mathcal{J}_X \otimes_{\mathcal{O}_\Sigma} \mathcal{O}_\Sigma(D) \cong \mathcal{J}_{Y/\Sigma}(D).$$

Thus $\dim_{k(Y)} H^1(\text{pr}^{-1}(Y), \mathcal{J}_X(D)_Y) = h^1(\Sigma, \mathcal{J}_{Y/\Sigma}(D))$, and the upper semicontinuity of α implies that the set $U = \{Y \in \text{Hilb}_\Sigma^n \mid h^1(\Sigma, \mathcal{J}_{Y/\Sigma}(D)) = 0\}$ is a non-empty open subset of Hilb_Σ^n . Since the latter is smooth and connected, hence irreducible, U is dense and irreducible. \square

F.2 Lemma

Let $\varphi : B \rightarrow A$ and $\psi : C \rightarrow A$ be morphisms of rings. If the A -module M is flat over B and if N is a free C -module, then $M \otimes_C N$ is a flat B -module.

Proof: Let $0 \rightarrow K' \rightarrow K \rightarrow K'' \rightarrow 0$ be an exact sequence of B -modules. Since M is flat over B , we have

$$0 \rightarrow K' \otimes_B M \rightarrow K \otimes_B M \rightarrow K'' \otimes_B M \rightarrow 0$$

is an exact sequence of B -modules, and thus via φ also of A -modules, but then via ψ of C -modules. Since N is free, hence flat, over C we moreover have

$$0 \rightarrow K' \otimes_B M \otimes_C N \rightarrow K \otimes_B M \otimes_C N \rightarrow K'' \otimes_B M \otimes_C N \rightarrow 0$$

is an exact sequence of C -modules, and thus it is exact as a sequence of B -modules. \square

F.3 Lemma

Let $f : X \rightarrow Y$ and $g : X \rightarrow Z$ be two morphisms of schemes. If the \mathcal{O}_X -module \mathcal{F} is flat over \mathcal{O}_Y and if \mathcal{G} is a locally free \mathcal{O}_Z -module, then $\mathcal{F} \otimes_{\mathcal{O}_X} g^* \mathcal{G}$ is a flat \mathcal{O}_Y -module.

Proof: Let $x \in X$ be given. We set $y = f(x)$, $z = g(x)$, $A = \mathcal{O}_{X,x}$, $B = \mathcal{O}_{Y,y}$, $C = \mathcal{O}_{Z,z}$, $\varphi = f^\# : B \rightarrow A$, $\psi = g^\# : C \rightarrow A$, $M = \mathcal{F}_x$ and $N = \mathcal{G}_z$. By assumption M is flat over B and N is free over C . Moreover,

$$(\mathcal{F} \otimes_{\mathcal{O}_X} g^* \mathcal{G})_x = (\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_X \otimes_{g^{-1}\mathcal{O}_Z} g^{-1}\mathcal{G})_x = \mathcal{F}_x \otimes_{\mathcal{O}_{Z,z}} \mathcal{G}_z = M \otimes_C N.$$

Thus the statement follows from Lemma F.2 \square

G. Examples of Surfaces

G.a. Geometrically Ruled Surfaces

Let $\Sigma = \mathbb{P}(\mathcal{E}) \xrightarrow{\pi} C$ be a geometrically ruled surface with normalised bundle \mathcal{E} (in the sense of [Har77] V.2.8.1). The Néron–Severi group of Σ is

$$\mathrm{NS}(\Sigma) = C_0\mathbb{Z} \oplus F\mathbb{Z},$$

with intersection matrix

$$\begin{pmatrix} -e & 1 \\ 1 & 0 \end{pmatrix},$$

where $F \cong \mathbb{P}_C^1$ is a fibre of π , C_0 a section of π with $\mathcal{O}_\Sigma(C_0) \cong \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$, $g = g(C)$ the genus of C , $\epsilon = \Lambda^2 \mathcal{E}$ and $e = -\deg(\epsilon) \geq -g$.²³ For the canonical divisor we have

$$K_\Sigma \sim_a -2C_0 + (2g - 2 - e) \cdot F,$$

where $g = g(C)$ is the genus of the base curve C .

In [Har77] V.2.18, V.2.20 and V.2.21 there one finds numerical criteria for an algebraic divisor class $aC_0 + bF$ to be ample respectively to contain irreducible curves. The following lemma on the nefness of $aC_0 + bF$ is an easy deduction of these statements.

G.1 Lemma

Let $C \sim_a aC_0 + bF$ be given.

- (a) If $e \geq 0$, then C is nef if and only if $a \geq 0$ and $b \geq ae$.
- (b) If $e \leq 0$, then C is nef if and only if $a \geq 0$ and $b \geq \frac{ae}{2}$.
- (c) If $e \leq 0$, then any curve $C \subset \Sigma$ is nef.

Proof: For (a) and (b) we note that a divisor is nef if and only if it belongs to the closure of the ample cone in $\mathrm{NS}(\Sigma) \otimes \mathbb{R}$ (cf. [IsS96] p. 162 and p. 175). The results thus follow from [Har77] V.2.20 b. and V.2.21 b.

It suffices to show (c) for irreducible curves, since the sum of nef divisors is nef. If, however, C is irreducible, then by [Har77] V.2.20 a. and V.2.21 a. the divisor is nef in view of (a) and (b). □

In Chapter II we have to examine special irreducible curves on Σ . Thus the following Lemma is of interest.

G.2 Lemma

Let $B \in |aC_0 + bF|_a$ be an irreducible curve with $B^2 = 0$ and $\dim |B|_a \geq 1$. Then we are in one of the following cases

²³By [Nag70] Theorem 1 there is some section $D \sim_a C_0 + bF$ with $g \geq D^2 = 2b - e$. Since D is irreducible, by [Har77] V.2.20/21 $b \geq 0$, and thus $-g \leq e$.

(G.1) $a = 0, b = 1, \text{ and } B \sim_a F,$

(G.2) $e = 0, a \geq 1, b = 0, \text{ and } B \sim_a aC_0, \text{ or}$

(G.3) $e < 0, a \geq 2, b = \frac{ae}{2} < 0, \text{ and } B \sim_a aC_0 + \frac{ae}{2} \cdot F.$

Moreover, if $a = 1$, then $\Sigma \cong C_0 \times \mathbb{P}_c^1$.

Proof: Since B is irreducible, we have

$$0 \leq B.F = a \quad \text{and} \quad 0 \leq B.C_0 = b - ae. \tag{G.4}$$

If $a = 0$, then $|B|_a = |bF|_a$, but since the general element of $|B|_a$ is irreducible, b has to be one, and we are in case (G.1).

We, therefore, may assume that $a \geq 1$. Since $B^2 = 0$ we have

$$0 = B^2 = 2a \left(b - \frac{ae}{2} \right), \quad \text{hence} \quad b = \frac{ae}{2}. \tag{G.5}$$

Combining this with (G.4) we get $e \leq 0$.

Moreover, if $e = 0$, then of course $b = 0$, while, if $e < 0$, then $a \geq 2$ by [Har77] V.2.21, since otherwise b would have to be non-negative. This brings us down to the cases (G.2) and (G.3).

It remains to show, that $B.F = a = 1$ implies $\Sigma \cong C_0 \times \mathbb{P}_c^1$. But by assumption the elements of $|B|_a$ are disjoint sections of the fibration π . Thus, by Lemma G.3, $\Sigma \cong C \times \mathbb{P}_c^1$. □

G.3 Lemma

If $\pi : \Sigma \rightarrow C$ has three disjoint sections, then Σ is isomorphic to $C \times \mathbb{P}_c^1$ as a ruled surface, i. e. there is an isomorphism $\alpha : \Sigma \rightarrow C \times \mathbb{P}_c^1$ such that the following diagram is commutative:

$$\begin{array}{ccc} \Sigma & \xrightarrow{\alpha} & C \times \mathbb{P}_c^1 \\ & \searrow \pi & \swarrow \text{pr} \\ & C & \end{array}$$

Proof: π is a locally trivial \mathbb{P}_c^1 -bundle, thus C is covered by a finite number of open affine subsets $U_i \subset C$ with trivialisations

$$\begin{array}{ccc} \pi^{-1}(U_i) & \xrightarrow[\varphi_i]{\cong} & U_i \times \mathbb{P}_c^1 \\ & \searrow \pi & \swarrow \text{pr} \\ & U_i & \end{array}$$

which are linear on the fibres.

The three disjoint sections on Σ , say S_0, S_1 , and S_∞ , give rise to three sections S_0^i, S_1^i , and S_∞^i on $U_i \times \mathbb{P}_c^1$. For each point $z \in U_i$ there is a unique linear projectivity on the fibre $\{z\} \times \mathbb{P}_c^1$ mapping the three points $p_{0,z} = S_0^i \cap (\{z\} \times \mathbb{P}_c^1)$, $p_{1,z} = S_1^i \cap (\{z\} \times \mathbb{P}_c^1)$, and $p_{\infty,z} = S_\infty^i \cap (\{z\} \times \mathbb{P}_c^1)$ to the standard basis $0 \equiv (z, (1 :$

$0))$, $1 \equiv (z, (1 : 1))$, and $\infty \equiv (z, (0 : 1))$ of $\mathbb{P}_c^1 \cong \{z\} \times \mathbb{P}_c^1$. If $p_{0,z} = (z, (x_0 : y_0))$, $p_{1,z} = (z, (x_1 : y_1))$, and $p_{\infty,z} = (z, (x_\infty : y_\infty))$, the projectivity is given by the matrix

$$A = \begin{pmatrix} \frac{(x_0 y_1 - y_0 x_1) y_\infty}{y_0 y_1 x_\infty^2 - y_0 x_1 x_\infty y_\infty - x_0 y_1 x_\infty y_\infty + x_0 x_1 y_\infty} & \frac{(x_0 y_1 - y_0 x_1) x_\infty}{y_0 y_1 x_\infty^2 - y_0 x_1 x_\infty y_\infty - x_0 y_1 x_\infty y_\infty + x_0 x_1 y_\infty} \\ \frac{y_0}{x_0 y_\infty - y_0 x_\infty} & \frac{x_0}{x_0 y_\infty - y_0 x_\infty} \end{pmatrix},$$

whose entries are rational functions in the coordinates of $p_{0,z}$, $p_{1,z}$, and $p_{\infty,z}$. Inserting for the coordinates local equations of the sections, A finally gives rise to an isomorphism of \mathbb{P}_c^1 -bundles, i. e. a morphism which is a linear isomorphism on the fibres,

$$\alpha_i : U_i \times \mathbb{P}_c^1 \rightarrow U_i \times \mathbb{P}_c^1$$

mapping the sections S_0^i , S_1^i , and S_∞^i to the trivial sections.

The transition maps

$$U_{ij} \times \mathbb{P}_c^1 \xrightarrow{\alpha_{ij}^{-1}} U_{ij} \times \mathbb{P}_c^1 \xrightarrow{\varphi_{ij}^{-1}} \pi^{-1}(U_{ij}) \xrightarrow{\varphi_{ij}} U_{ij} \times \mathbb{P}_c^1 \xrightarrow{\alpha_{ij}} U_{ij} \times \mathbb{P}_c^1,$$

with $U_{ij} = U_i \cap U_j$, are linear on the fibres and fix the three trivial sections. Thus they must be the identity maps, which implies that the $\alpha_i \circ \varphi_i$, $i = 1, \dots, r$, glue together to an isomorphism of ruled surfaces:

$$\begin{array}{ccc} \Sigma & \xrightarrow{\alpha} & C \times \mathbb{P}_c^1 \\ & \searrow \pi & \swarrow \text{pr} \\ & C & \end{array}$$

See also [IsS96] p. 229. □

Knowing the algebraic equivalence classes of irreducible curves in Σ which satisfy the assumptions in Condition (II.1.5) we can give a better formulation of the vanishing theorem in the case of geometrically ruled surfaces.

In order to do the same for the existence theorems, we have to study very ample divisors on Σ . These, however, depend very much on the structure of the base curve C ,²⁴ and the general results which we give may be not the best possible. Only in the case $C = \mathbb{P}_c^1$ we can give a complete investigation.

G.4 Remark

The geometrically ruled surfaces with base curve \mathbb{P}_c^1 are, up to isomorphism, just the Hirzebruch surfaces $F_e = \mathbb{P}(\mathcal{O}_{\mathbb{P}_c^1} \oplus \mathcal{O}_{\mathbb{P}_c^1}(-e))$, $e \geq 0$. Note that $\text{Pic}(F_e) = \text{NS}(F_e)$, that is, algebraic equivalence and linear equivalence coincide. Moreover, by [Har77] V.2.18 a divisor class $L = \alpha C_0 + \beta F$ is very ample

²⁴Cf. [Har77] V.2.22.2, Ex. V.2.11 and Ex. V.2.12.

over \mathbb{C} if and only if $\alpha > 0$ and $\beta > \alpha e$. The conditions throughout the existence theorems turn out to be optimal if we work with $L = C_0 + (e + 1) \cdot F$, while for other choices of L they become more restrictive.²⁵

In the case $C \not\cong \mathbb{P}_\mathbb{C}^1$, we may choose an integer $l \geq \max\{e + 1, 2\}$ such that the algebraic equivalence class $|C_0 + lF|_a$ contains a very ample divisor L , e. g. $l = e + 3$ will do, if C is an elliptic curve.²⁶ In particular, $l \geq 2$ as soon as $\Sigma \not\cong \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$.

With the above choice of L we have $g(L) = 1 + \frac{L^2 + L \cdot K_\Sigma}{2} = 1 + \frac{(-e+2l) + (e-2l+2g-2)}{2} = g$, and hence the generic curve in $|L|_1$ is a smooth curve whose genus equals the genus of the base curve.

Finally, in several situations we need a thorough knowledge on the cohomology groups of a divisor on a surface. We therefore prove the following lemmata.

G.5 Lemma

Let $D \sim_1 aC_0 + bF$ with $b \in \text{Div}(C)$ of degree $b = \deg(b)$, let $a' = \max\{k \mid a \geq k \geq 0, b \geq ke\}$, $a'' = \min\{k \mid a \geq k \geq 0, b \geq ke\}$, $a^* = \min\{k \mid a \geq k \geq 0, b \leq ke + 2g - 2\}$ and $a^{**} = \max\{k \mid a \geq k \geq 0, b \leq ke + 2g - 2\}$.²⁷

(a) Let $a < 0$, then $h^0(\Sigma, D) = 0$.

(b) Let $a \geq 0$. If $e \geq 0$, then

$$\begin{aligned} h^0(\Sigma, D) &= h^0(\Sigma, a'C_0 + bF) \leq \sum_{k=0}^{a'} h^0(C, \mathcal{O}_C(k \cdot e + b)) \\ &\leq a'b + a' + b + 1 - e \cdot \frac{a'(a'+1)}{2}, \end{aligned}$$

while if $e \leq 0$, then

$$\begin{aligned} h^0(\Sigma, D) &\leq \sum_{k=a''}^a h^0(C, \mathcal{O}_C(k \cdot e + b)) \\ &\leq \left(ab + a + b + 1 - e \cdot \frac{a(a+1)}{2} \right) - \left(a''b + a'' - e \cdot \frac{a''(a''-1)}{2} \right). \end{aligned}$$

²⁵ Let $L' = \alpha C_0 + \beta F$, then $D - L' - K_{F_e} = (a + 1 - \alpha) \cdot C_0 + (b + 1 + e - \beta) \cdot F$, and thus the optimality of the conditions follows from

(III.1.8b) $(D - L' - K_{F_e})^2 = (a + 1 - \alpha) \cdot (2 \cdot (b + 1 + e - \beta) - (a + 1 - \alpha) \cdot e) \leq a((2b - ae) + (\alpha e + e + 2 - 2\beta)) \leq a \cdot (2b - ae) = (D - L - K_{F_e})^2,$

(III.1.9b) $(D - L' - K_{F_e}) \cdot F = a + 1 - \alpha \leq a = (D - L - K_{F_e}) \cdot F$, and for $e = 0$, $(D - L' - K_{F_e}) \cdot C_0 = b + 1 - \beta \leq b = (D - L - K_{F_e}) \cdot C_0$, and

(III.1.11b) $b + 1 + e - \beta \geq e(a + 1 - \alpha)$ implies $b \geq b + e\alpha + 1 - \beta \geq ae$.

²⁶ l will be the degree of a suitable very ample divisor ϑ on C . Now ϑ defines an embedding of C into some $\mathbb{P}_\mathbb{C}^N$ such that the degree of the image C' is just $\deg(\vartheta)$. Therefore, $\deg(\vartheta) \geq 2$, unless C' is linear (cf. [Har77] Ex. I.7.6), which implies $C \cong \mathbb{P}_\mathbb{C}^1$.

²⁷If a minimum does not exist, we set its value to $a + 1$, and likewise we set a non-existing maximum to -1 .

(c) Let $a \geq 0$. If $e \geq 0$, then

$$h^1(\Sigma, D) \leq \sum_{k=a^*}^a h^1(C, \mathcal{O}_C(k \cdot e + b)),$$

while if $e \leq 0$, then

$$h^1(\Sigma, D) \leq \sum_{k=0}^{a^{**}} h^1(C, \mathcal{O}_C(k \cdot e + b)).$$

(d) If \mathcal{E} is decomposable and $a \geq 0$, then for any $i \in \mathbb{Z}$ and $a \geq 0$

$$h^i(\Sigma, D) = \sum_{k=0}^{a'} h^i(C, \mathcal{O}_C(k \cdot e + b)).$$

(e) If $e = 0$ and D nef, then $a' = a$ and $a'' = 0$.

(f) Let $g(C) = 1$ and $a \geq 0$. If $e \geq 0$, then

$$h^0(\Sigma, D) \leq a'b + b + 1 - e \cdot \frac{a'(a'+1)}{2},$$

and if in addition $b > a'e$, then even

$$h^0(\Sigma, D) \leq a'b + b - e \cdot \frac{a'(a'+1)}{2}.$$

If²⁸ $e = -1$, then

$$h^0(\Sigma, D) \leq \begin{cases} ab + b + \frac{a(a+1)}{2}, & \text{if } b > 0, \\ ab + \frac{a(a+1)}{2} + \frac{b(b+1)}{2}, & \text{if } 0 \geq b \geq -\frac{a}{2} \text{ and } b \not\sim_1 b \cdot e, \\ ab + 1 + \frac{a(a+1)}{2} + \frac{b(b+1)}{2}, & \text{if } 0 \geq b \geq -\frac{a}{2} \text{ and } b \sim_1 b \cdot e, \\ 0, & \text{if } -\frac{a}{2} > b. \end{cases}$$

(g) If $e < 0$ and $a \geq 0$, then setting $b' = b - \frac{ae}{2}$

$$h^0(\Sigma, D) \leq ab' + a + b' + 1 - \frac{1}{2e} \cdot (b + 1 + \frac{e}{2})^2.$$

Moreover, if D is effective, then

$$h^0(\Sigma, D) \leq ab' + a + b' + 1 - \frac{1}{2e} \cdot (b' + \frac{3ea}{4})^2$$

and

$$h^0(\Sigma, D) \leq ab' + a + b' + 1 - \frac{9e}{32} \cdot a^2.$$

Proof:

(a) If $a < 0$, then $0 > a = D.F$, which is impossible for any curve.

(b) We consider the exact sequence

$$0 \rightarrow \mathcal{O}_\Sigma((a-1)C_0 + bF) \rightarrow \mathcal{O}_\Sigma(aC_0 + bF) \rightarrow \mathcal{O}_{C_0}(aC_0 + bF) \rightarrow 0. \quad (\text{G.6})$$

Taking into account that $\mathcal{O}_{C_0}(aC_0 + bF) \cong \mathcal{O}_C(ae + b)$ the long exact cohomology sequence of (G.6) gives

$$h^0(\Sigma, aC_0 + bF) \leq h^0(\Sigma, (a-1)C_0 + bF) + h^0(C, \mathcal{O}_C(ae + b)).$$

²⁸If $g = 1$ and $e < 0$, then $e = -1$.

By induction on a , starting with $a = 0$, we thus get

$$h^0(\Sigma, D) \leq \sum_{k=0}^a h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})).$$

For the induction basis $a = 0$ we note that by [Har77] Ex. III.8.3 and Ex. III.8.4 and by the projection formula we have for all $i > 0$

$$R^i \pi_*(\mathfrak{b}F) = R^i \pi_*(\mathcal{O}_\Sigma \otimes \pi^* \mathfrak{b}) = R^i \pi_* \mathcal{O}_\Sigma \otimes \mathfrak{b} = 0,$$

and thus by [Har77] Ex. III.8.1,

$$h^i(\Sigma, \mathfrak{b}F) = h^i(C, \mathfrak{b}) \quad \text{for all } i \geq 0. \quad (\text{G.7})$$

If $e \geq 0$ and $k > a'$ respectively if $e \leq 0$ and $k < a''$, then $\deg(k\epsilon + \mathfrak{b}) = -ke + \mathfrak{b} < 0$ and

$$h^0(C_0, \mathcal{O}_{C_0}(kC_0 + \mathfrak{b}F)) = h^0(C, \mathcal{O}_C(k\epsilon + \mathfrak{b})) = 0.$$

Thus, if $e \geq 0$ we have

$$h^0(\Sigma, D) \leq \sum_{k=0}^{a'} h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})),$$

and if $e \leq 0$

$$h^0(\Sigma, D) \leq \sum_{k=a''}^a h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})).$$

By Lemma D.3 for $e \geq 0$ and $k \leq a'$ respectively $e \leq 0$ and $k \geq a''$ we have $h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})) \leq \mathfrak{b} - ke + 1$. Thus if $e \geq 0$

$$\sum_{k=0}^{a'} h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})) \leq a' \mathfrak{b} + a' + \mathfrak{b} + 1 - e \cdot \frac{a'(a'+1)}{2},$$

and

$$\begin{aligned} & \sum_{k=a''}^a h^0(C, \mathcal{O}_C(k \cdot \epsilon + \mathfrak{b})) \\ & \leq \left(a\mathfrak{b} + a + \mathfrak{b} + 1 - e \cdot \frac{a(a+1)}{2} \right) - \left(a''\mathfrak{b} + a'' - e \cdot \frac{a''(a''-1)}{2} \right). \end{aligned}$$

It remains to show that for $e \geq 0$ and $a > a'$,

$$h^0(\Sigma, aC_0 + \mathfrak{b}F) = h^0(\Sigma, a'C_0 + \mathfrak{b}F).$$

Since $a > a'$, $\deg(a\epsilon + \mathfrak{b}) = -a\epsilon + \mathfrak{b} < 0$, the long exact cohomology sequence of (G.6) implies

$$h^0(\Sigma, aC_0 + \mathfrak{b}F) = h^0(\Sigma, (a-1)C_0 + \mathfrak{b}F).$$

The result thus follows by descending induction on a for $a > a'$.

(c) The long exact cohomology sequence of (G.6) gives

$$H^1(\Sigma, (a-1)C_0 + bF) \rightarrow H^1(\Sigma, aC_0 + bF) \rightarrow H^1(C_0, \mathcal{O}_{C_0}(aC_0 + bF)).$$

We thus get

$$h^1(\Sigma, aC_0 + bF) \leq h^1(\Sigma, (a-1)C_0 + bF) + h^1(C, \mathcal{O}_C(a\epsilon + b)),$$

and by induction on a , starting with $a = 0$ – see (G.7) –, we find

$$h^1(\Sigma, D) \leq \sum_{k=0}^a h^1(C, \mathcal{O}_C(k \cdot \epsilon + b)).$$

However, if $e \geq 0$ and $k < a^*$ respectively if $e \leq 0$ and $k > a^{**}$, then $\deg(k\epsilon + b) = b - ke > 2g - 2$ and thus $h^1(C, k\epsilon + b) = 0$.

- (d) See [FuP00] Lemma 2.10. Replacing the section C_0 by a section in $|C_0 - \epsilon F|_a$, which only exists in the decomposable case, they do basically the same considerations as in (a) – however, they get surjectivity at the important positions in the long exact cohomology sequence.
- (e) If $e = 0$ and D nef, then by Lemma G.1 we know that $a, b \geq 0$ and thus $a' = a$ and $a'' = 0$.
- (f) Let us consider the case $e \geq 0$ first. By part (a) we know $h^0(\Sigma, D) \leq \sum_{k=0}^{a'} h^0(C, \mathcal{O}_C(k \cdot \epsilon + b))$. Moreover, we have

$$\deg(k \cdot \epsilon + b) = b - ke \begin{cases} > 0, & \text{if } k = 0, \dots, a' - 1, \\ & \text{and if } k = a' \text{ and } b > a'e, \\ \geq 0, & \text{if } k = a'. \end{cases}$$

By Lemma G.14 we thus get

$$h^0(C, \mathcal{O}_C(k \cdot \epsilon + b)) \leq \begin{cases} b - ke, & \text{if } k = 0, \dots, a' - 1, \\ & \text{and if } k = a' \text{ and } b > a'e, \\ b - ke + 1, & \text{if } k = a'. \end{cases}$$

Thus the claim for $e \geq 0$ follows. Let now $e = -1$. The claim then is implied by the following considerations.

If $b > 0$, then obviously $b > -k = ek$ for all $k = 0, \dots, a$ and thus $a'' = 0$. Moreover, as above we see $\deg(k \cdot \epsilon + b) = b + k > 0$ for all $k = 0, \dots, a$ in this case, and thus $h^0(C, \mathcal{O}_C(k \cdot \epsilon + b)) = b + k$ for all k .

If $0 \geq b \geq -a$, then $a'' = -b$ and $\deg(k \cdot \epsilon + b) = b + k > 0$ for $k = -b + 1, \dots, a$. Therefore we get $h^0(C, \mathcal{O}_C(k \cdot \epsilon + b)) = b + k$ for all $k = -b + 1, \dots, a$. If, moreover, $b - b \cdot e \not\sim 0$, then $h^0(C, \mathcal{O}_C(-b \cdot \epsilon + b)) = 0$, while if $b - b \cdot e \sim 0$, then $h^0(C, \mathcal{O}_C(-b \cdot \epsilon + b)) = 1$.

Finally, if $-\frac{a}{2} > b$ then $|D|_1$ is empty by [Har77] V.2.21.

- (g) Since e is negative, any of the expressions $-\frac{1}{2e} \cdot (b+1+\frac{e}{2})^2$, $-\frac{1}{2e} \cdot (b'+\frac{3ea}{4})^2$ and $-\frac{9e}{32} \cdot a^2$ is non-negative. Moreover, for $b \geq 0$ we have $a'' = 0$ and thus $h^0(\Sigma, D) = ab' + a + b' + 1$. The inequalities are thus fulfilled for $b \geq 0$.

It remains to verify them for $b < 0$, that is, for $b < 0$ we have to show that $-(a''b + a'' - e \cdot \frac{a'' \cdot (a''-1)}{2})$ is bounded from above by any of these three expressions.

Let $b < 0$ be given, and consider the function

$$\varphi : \mathbb{R} \rightarrow \mathbb{R} : t \mapsto \frac{e}{2} \cdot t^2 - (b+1+\frac{e}{2}) \cdot t.$$

The derivative $\frac{\partial \varphi}{\partial t}(t) = e \cdot t - (b+1+\frac{e}{2})$ has its unique zero at $t = \frac{1}{e} \cdot (b+1+\frac{e}{2})$ and φ thus takes its maximum there. That is,

$$\begin{aligned} -\left(a''b + a'' - e \cdot \frac{a'' \cdot (a''-1)}{2}\right) &= \varphi(a'') \\ &\leq \varphi\left(\frac{1}{e} \cdot (b+1+\frac{e}{2})\right) = -\frac{1}{2e} \cdot (b+1+\frac{e}{2})^2. \end{aligned}$$

This proves the first inequality.

If D is effective and $b < 0$, then by [Har77] V.2.21 $a \geq 2$ and $0 > b \geq \frac{ae}{2}$. But then $b' + \frac{3ea}{4} = b + \frac{ea}{4} < b+1+\frac{e}{2} < 0$ and therefore

$$-\frac{1}{2e} \cdot (b+1+\frac{e}{2})^2 < -\frac{1}{2e} \cdot (b'+\frac{3ea}{4})^2,$$

which gives the second inequality. And finally, since $b' = b - \frac{ae}{2} \geq 0$ in this situation, we get $\frac{3ea}{4} \leq b' + \frac{3ea}{4} < 0$ and thus

$$-\frac{1}{2e} \cdot (b'+\frac{3ea}{4})^2 \leq -\frac{9e}{32} \cdot a^2.$$

□

G.6 Lemma

Let $\Sigma = \mathbb{F}_e$ and let $D = aC_0 + bF$, $a' = \max\{k \mid a \geq k, b \geq ke\}$ and $a^* = \max\{k \mid -a-2 \geq k, b+2+e \leq ke\}$.

$$(a) \quad h^1(\Sigma, D) = \begin{cases} 0, & \text{if } a = -1 \text{ or } (a \geq 0 \text{ and } b \geq ae - 1) \\ & \text{or } (a \leq -2 \text{ and } b \leq -(a+1) \cdot e - 1), \\ e \cdot \frac{a \cdot (a+1)}{2} - ab - a - b - 1 \neq 0, & \text{if } a \geq 0 \text{ and } b < 0 \\ & \text{and } (a, b) \neq (0, -1), \\ e \cdot \frac{l \cdot (l+1)}{2} - l \cdot (k+1) \neq 0, & \text{if } a > 0 \text{ and} \\ & 0 \leq b = (a-l) \cdot e + k \leq ae - 2 \\ & \text{with } 1 < l \leq a, 0 \leq k < e, \\ h^1(\Sigma, K_\Sigma - D), & \text{if } a \leq -2. \end{cases}$$

$$(b) \quad h^0(\Sigma, D) = \begin{cases} 0, & \text{if } a < 0 \text{ or } b < 0, \\ ab + a + b + 1 - e \cdot \frac{a \cdot (a+1)}{2}, & \text{if } 0 \leq a \text{ and } ae \leq b, \\ a'b + a' + b + 1 - e \cdot \frac{a'(a'+1)}{2}, & \text{if } 0 \leq a \text{ and } 0 \leq b \leq ae. \end{cases}$$

$$(c) \quad h^2(\Sigma, D) = \begin{cases} 0 & \text{if } a > -2 \text{ or } b > -2 - e, \\ ab + a + b + 1 - e \cdot \frac{a \cdot (a+1)}{2} & \text{if } a \leq -2, b + 2 + e \leq ae, \\ a^*b + a^* + b + 1 - e \cdot \frac{a^*(a^*+1)}{2} & \text{if } a \leq -2 \text{ and} \\ & ae < b + 2 + e \leq 0. \end{cases}$$

Proof:

(a) By the Theorem of Riemann-Roch we have

$$\chi(D) = \frac{D^2 - D \cdot K_\Sigma}{2} + \chi(\mathcal{O}_\Sigma) = ab + a + b + 1 - e \cdot \frac{a(a+1)}{2}.$$

The result is thus an application of part (b) and (c).

(b) This follows from Lemma G.5 (d) taking into account that for a divisor $\mathcal{O}_{\mathbb{P}_c^1}(d)$ on \mathbb{P}_c^1 of non-negative degree we have $h^0(\mathbb{P}_c^1, \mathcal{O}_{\mathbb{P}_c^1}(d)) = d + 1$.

(c) By Serre duality we have

$$h^2(\Sigma, D) = h^0(\Sigma, K_\Sigma - D) = h^0(\Sigma, (-a - 2) \cdot C_0 + (-b - 2 - e) \cdot F).$$

The result thus follows by simple calculations from part (b).

□

G.b. Products of Curves

Let C_1 and C_2 be two smooth projective curves of genera $g_1 \geq 0$ and $g_2 \geq 0$ respectively. The surface $\Sigma = C_1 \times C_2$ is naturally equipped with two fibrations $\text{pr}_i : \Sigma \rightarrow C_i$, $i = 1, 2$, and by abuse of notation we denote two generic fibres $\text{pr}_2^{-1}(p_2) = C_1 \times \{p_2\}$ resp. $\text{pr}_1^{-1}(p_1) = \{p_1\} \times C_2$ again by C_1 resp. C_2 .

G.7 Lemma

Let $D \in \text{Div}(\Sigma)$ be a divisor such that $D \sim_1 \text{pr}_2^* a + \text{pr}_1^* b$ with $a \in \text{Div}(C_2)$ of degree a and $b \in \text{Div}(C_1)$ of degree b .

- (a) $D \sim_a aC_1 + bC_2$.
- (b) If a is very ample on C_2 and b is very ample on C_1 , then $\text{pr}_2^* a \otimes \text{pr}_1^* b$ is very ample on Σ .
- (c) If $a \geq 2g_2 + 1$ and $b \geq 2g_1 + 1$, then $\text{pr}_2^* a \otimes \text{pr}_1^* b$ is very ample on Σ .
- (d) D is ample if and only if $a, b > 0$.
- (e) D is nef if and only if $a, b \geq 0$.
- (f) D is effective if and only if $a, b \geq 0$ and $(a, b) \neq (0, 0)$.

$$(g) \quad K_{\Sigma} \cong \text{pr}_2^* K_{C_2} + \text{pr}_1^* K_{C_1} \sim_a (2g_2 - 2)C_1 + (2g_1 - 2)C_2.$$

Proof:

- (a) Obviously $D \sim_1 \text{pr}_2^* a + \text{pr}_1^* b \sim_a a'C_1 + b'C_2$ for some $a', b' \in \mathbb{Z}$. But then $a' = D.C_2 = \deg(\mathcal{O}_{C_2}(D)) = \deg(a)$ and $b' = D.C_1 = \deg(\mathcal{O}_{C_1}(D)) = \deg(b)$.
- (b) a and b induce embeddings of C_2 and C_1 into some projective spaces $\mathbb{P}_{\mathbb{C}}^n$ and $\mathbb{P}_{\mathbb{C}}^m$ respectively. Thus $\text{pr}_2^* a \otimes \text{pr}_1^* b$ induces the embedding into the product of these spaces and is the pullback of the very ample line bundle $\mathcal{O}_{\mathbb{P}_{\mathbb{C}}^n}(1) \otimes \mathcal{O}_{\mathbb{P}_{\mathbb{C}}^m}(1)$ which induces the Segre embedding of $\mathbb{P}_{\mathbb{C}}^n \times \mathbb{P}_{\mathbb{C}}^m$.
- (c) By [Har77] IV.3.2 then a and b are very ample, and we are done with (b).
- (d) If D is ample, then $a = D.C_2 > 0$ and $b = D.C_1 > 0$. On the other hand, if $a, b > 0$ we may choose some $n > 0$ such that $na \geq 2g_2 + 1$ and $nb \geq 2g_1 + 1$, and thus nD is very ample, which implies that D itself is ample.
- (e) If D is nef, then $a = D.C_2 \geq 0$ and $b = D.C_1 \geq 0$.
Now let $a, b \geq 0$. Since any irreducible curve which is not a fibre of pr_i is intersected by some fibre of pr_i , the fibres of pr_i are nef, $i = 1, 2$. Hence, D is nef if $a = 0$ or $b = 0$. If, however, both are strictly positive, then D is even ample, hence in particular nef.
- (f) If D is effective, then $0 < D.(nC_1 + mC_2) = ma + nb$ for all $n, m > 0$. Hence, neither a nor b can be negative. The converse is obvious.
- (g) See [Har77] Exercise II.8.3 and Example IV.1.3.3.

□

In many situations we are interested in the cohomology groups of a divisor of the type $\text{pr}_1^* a + \text{pr}_2^* b$. The Künneth-Formula implies the following useful results.

G.8 Lemma

Let $a \in \text{Div}(C_2)$ and $b \in \text{Div}(C_1)$ and let $i \in \mathbb{Z}$, then

$$h^i(\Sigma, \text{pr}_2^* a + \text{pr}_1^* b) = \sum_{k=0}^i h^k(C_2, a) \cdot h^{i-k}(C_1, b).$$

In particular, $h^0(\Sigma, \text{pr}_2^* a + \text{pr}_1^* b) = h^0(C_2, a) \cdot h^0(C_1, b)$.

Proof: The Künneth-Formula (cf. [Dan96] I.1.7) gives

$$H^i(\Sigma, \text{pr}_2^* a + \text{pr}_1^* b) = \sum_{k=0}^i H^k(C_2, a) \otimes H^{i-k}(C_1, b),$$

and thus the result follows. □

G.9 Corollary

Let $D \in \text{Div}(\Sigma)$ be a divisor such that $D \sim_1 aC_1 + bC_2$ with $a \in \text{Div}(C_2)$ of degree $a \geq 0$ and $b \in \text{Div}(C_1)$ of degree $b \geq 0$. Then:

$$h^0(\Sigma, \text{pr}_2^* a + \text{pr}_1^* b) \leq (a + 1) \cdot (b + 1).$$

Proof: By Lemma G.8 it suffices to show that for a divisor $c \in \text{Div}(C)$ on a smooth projective curve C we have $h^0(C, c) \leq \deg(c) + 1$. This, however, follows from Lemma D.3. \square

G.10 Corollary

Suppose that $g_2 = 0$ and $g_1 = 1$, and let $D \in \text{Div}(\Sigma)$ be a divisor such that $D \sim_1 \text{pr}_2^* a + \text{pr}_1^* b$ with $a \in \text{Div}(C_2)$ of degree $a \geq 0$ and $b \in \text{Div}(C_1)$ of degree $b \geq 1$. Then:

$$h^0(\Sigma, \text{pr}_2^* a + \text{pr}_1^* b) = (a + 1) \cdot b.$$

Proof: This follows by Lemma G.8, taking into account that $h^0(C_2, a) = a + 1$ (cf. Lemma G.6) and $h^0(C_1, b) = b$ (cf. Lemma G.14). \square

G.11 Remark

We are going to show that for a generic choice of the curves C_1 and C_2 the Néron–Severi group $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z}$ of Σ is two-dimensional²⁹ with intersection matrix

$$(C_i \cdot C_j)_{i,j} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Thus, the only irreducible curves $B \subset \Sigma$ with selfintersection $B^2 = 0$ are the fibres C_1 and C_2 , and for any irreducible curve $B \sim_a aC_1 + bC_2$ the coefficients a and b must be non-negative.

Supposed that one of the curves is rational, the surface is geometrically ruled and the Picard number of Σ is two. Whereas, if both C_1 and C_2 are of strictly positive genus, this need no longer be the case as we will see in Remark G.18. Thus the following proposition is the best we may expect.

G.12 Proposition

For a generic choice of smooth projective curves C_1 and C_2 the Néron–Severi group of $\Sigma = C_1 \times C_2$ is $\text{NS}(\Sigma) \cong C_1\mathbb{Z} \oplus C_2\mathbb{Z}$.

More precisely, fixing g_1 and g_2 there is a very general subset $U \subseteq M_{g_1} \times M_{g_2}$ such that for any $(C_1, C_2) \in U$ the Picard number of $C_1 \times C_2$ is two, where M_{g_i} denotes the moduli space of smooth projective curves of genus g_i , $i = 1, 2$.

Proof: As already mentioned, if either g_1 or g_2 is zero, then we may take $U = M_{g_1} \times M_{g_2}$.

²⁹In the case that C_1 and C_2 are elliptic curves, generic means precisely, that they are not isogenous - see Section G.c.

Suppose that $g_1 = g_2 = 1$. Given an elliptic curve C_1 there is a countable union V of proper subvarieties of M_1 such that for any $C_2 \in M_1 \setminus V$ the Picard number of $C_1 \times C_2$ is two - namely, if τ_1 and τ_2 denote the periods as in Section G.c, then we have to require that there exists no invertible integer matrix $\begin{pmatrix} z_1 & z_2 \\ z_3 & z_4 \end{pmatrix}$ such that $\tau_2 = \frac{z_4 - z_3 \tau_1}{z_2 - z_1 \tau_1}$. (Compare also [GrH94] p. 286.)

We, therefore, may assume that $g_1 \geq 2$ and $g_2 \geq 1$. The claim then follows from Lemma G.13, which is due to Denis Gaitsgory. \square

G.13 Lemma (Denis Gaitsgory)

Let C_2 be any smooth projective curve of genus $g_2 \geq 1$. Then for any $g_1 \geq 2$ there is a very general subset \mathcal{U} of the moduli space M_{g_1} of smooth projective curves of genus g_1 such that the Picard number of $C_1 \times C_2$ is two for any $C_1 \in \mathcal{U}$.

Proof: We note that a curve $B \subset \Sigma = C_1 \times C_2$ with $C_1 \not\curvearrowright_a B \not\curvearrowright_a C_2$ induces a non-trivial morphism $\mu_B : C_1 \rightarrow \text{Pic}(C_2) : p \mapsto \text{pr}_{2*}(\text{pr}_1^*(p))$. It thus makes sense to study the moduli problem of (non-trivial) maps from curves of genus g_1 into $\text{Pic}(C_2)$.

More precisely, let $k \in \mathbb{N}$ and let $0 \neq \beta \in H_2(\text{Pic}_k(C_2), \mathbb{Z}) = \mathbb{Z}^{2g_2}$ be given, where $\text{Pic}_k(C_2)$ is the Picard variety of divisors of degree k on C_2 . Following the notation of [FuP97] we denote by $M_{g_1,0}(\text{Pic}_k(C_2), \beta)$ the moduli space of pairs (C_1, μ) , where C_1 is a smooth projective curve of genus g_1 and $\mu : C_1 \rightarrow \text{Pic}_k(C_2)$ a morphism with $\mu_*([C_1]) = \beta$. We then have the canonical morphism

$$F_{k,\beta} : M_{g_1,0}(\text{Pic}_k(C_2), \beta) \rightarrow M_{g_1} : (C_1, \mu) \mapsto C_1,$$

just forgetting the map μ , and the proposition reduces to the following claim.

Claim: For no choice of $k \in \mathbb{N}$ and $0 \neq \beta \in H_2(\text{Pic}_k(C_2), \mathbb{Z})$ the morphism $F_{k,\beta}$ is dominant.

Let $\mu : C_1 \rightarrow \text{Pic}_k(C_2)$ be any morphism with $\mu_*([C_1]) = \beta$. Then μ is not a contraction and the image of C_1 is a projective curve in the abelian variety $\text{Pic}_k(C_2)$. Moreover, we have the following exact sequence of sheaves

$$0 \longrightarrow \mathcal{T}_{C_1} \xrightarrow{d\mu} \mu^* \mathcal{T}_{\text{Pic}_k(C_2)} = \mathcal{O}_{C_1}^{g_2} \longrightarrow \mathcal{N}_\mu := \text{coker}(d\mu) \longrightarrow 0. \quad (\text{G.8})$$

Since $d\mu$ is a non-zero inclusion, its dual

$$d\mu^\vee : (\mu^* \mathcal{T}_{\text{Pic}_k(C_2)})^\vee = \mathcal{O}_{C_1}^{g_2} \rightarrow \Omega_{C_1} = \omega_{C_1}$$

is not zero on global sections, that is

$$\begin{array}{ccc} H^0(d\mu^\vee) : H^0(C_1, \mathcal{O}_{C_1}^{g_2}) & \longrightarrow & H^0(C_1, \omega_{C_1}) \\ \parallel & & \parallel \\ \text{Hom}_{\mathcal{O}_{C_1}}(\mathcal{O}_{C_1}^{g_2}, \mathcal{O}_{C_1}) & & \text{Hom}_{\mathcal{O}_{C_1}}(\mathcal{T}_{C_1}, \mathcal{O}_{C_1}) \end{array}$$

is not the zero map.

Since $g_1 \geq 2$ we have $h^0(C_1, \omega_{C_1}) = 2g_1 - 2 > 0$, and thus ω_{C_1} has global sections. Therefore, the induced map $H^0(C_1, \omega_{C_1} \otimes \mathcal{O}_{C_1}^{g_2}) \rightarrow H^0(C_1, \omega_{C_1} \otimes \omega_{C_1})$ is not the zero map, which by Serre duality gives that the map

$$H^1(d\mu) : H^1(C_1, \mathcal{T}_{C_1}) \rightarrow H^1(C_1, \mu^* \mathcal{T}_{\text{Pic}_k(C_2)})$$

from the long exact cohomology sequence of (G.8) is not zero. Hence the coboundary map

$$\delta : H^0(C_1, \mathcal{N}_\mu) \rightarrow H^1(C_1, \mathcal{T}_{C_1})$$

cannot be surjective. According to [Har98] p. 96 we have

$$\delta = dF_{k,\beta} : \mathcal{T}_{M_{g_1,0}(\text{Pic}_k(C_2),\beta)} = H^0(C_1, \mathcal{N}_\mu) \longrightarrow \mathcal{T}_{M_{g_1}} = H^1(C_1, \mathcal{T}_{C_1}).$$

But if the differential of $F_{k,\beta}$ is not surjective, then $F_{k,\beta}$ itself cannot be dominant. \square

G.c. Products of Elliptic Curves

Let $C_1 = \mathbb{C}/\Lambda_1$ and $C_2 = \mathbb{C}/\Lambda_2$ be two elliptic curves, where $\Lambda_i = \mathbb{Z} \oplus \tau_i \mathbb{Z} \subset \mathbb{C}$ is a lattice and τ_i is in the upper half plane of \mathbb{C} . We denote the natural group structure on each of the C_i by $+$ and the neutral element by 0 .

Our interest lies in the study of the surface $\Sigma = C_1 \times C_2$. This surface is naturally equipped with two fibrations $\text{pr}_i : \Sigma \rightarrow C_i$, $i = 1, 2$, and by abuse of notation we denote the fibres $\text{pr}_2^{-1}(0) = C_1 \times \{0\}$ resp. $\text{pr}_1^{-1}(0) = \{0\} \times C_2$ again by C_1 resp. C_2 . The group structures on C_1 and C_2 extend to Σ so that Σ itself is an abelian variety. Moreover, for $p = (p_1, p_2) \in \Sigma$ the mapping $\tau_p : \Sigma \rightarrow \Sigma : (a, b) \mapsto (a + p_1, b + p_2)$ is an automorphism of abelian varieties. Due to these translation morphisms we know that for any curve $B \subset \Sigma$ the algebraic family of curves $|B|_a$ covers the whole of Σ , and in particular $\dim |B|_a \geq 1$. This also implies $B^2 \geq 0$.

Since Σ is an abelian surface, $\text{NS}(\Sigma) = \text{Num}(\Sigma)$, $K_\Sigma = 0$, and the Picard number $\rho = \rho(\Sigma) \leq 4$ (cf. [LaB92] 4.11.2 and Ex. 2.5). But the Néron–Severi group of Σ contains the two independent elements C_1 and C_2 , so that $\rho \geq 2$. The general case³⁰ is indeed $\rho = 2$, however ρ might also be larger (see Example G.17), in which case the additional generators may be chosen to be graphs of surjective morphisms from C_1 to C_2 (cf. [IsS96] 3.2 Example 3). This shows:

$$\rho(\Sigma) = 2 \text{ if and only if } C_1 \text{ and } C_2 \text{ are not isogenous.}$$

The following lemma provides a better knowledge on the dimension of the global sections of a divisor on an elliptic curve, and leads to better results in the irreducibility and smoothness theorems, when products of elliptic curves are studied.

³⁰The abelian surfaces with $\rho \geq 2$ possessing a principle polarisation are parametrised by a countable number of surfaces in a three-dimensional space, and the Picard number of such an abelian surface is two unless it is contained the intersection of two or three of these surfaces (cf. [IsS96] 11.2). See also [GrH94] p. 286, and Proposition G.12.

G.14 Lemma

Let C be a smooth elliptic curve over \mathbb{C} and let a be a divisor on C of degree $a = \deg(a)$. Then:

- (a) $h^0(C, a) = \chi(a) = \deg(a)$ and $h^1(C, a) = 0$, if $a > 0$.
- (b) $h^0(C, a) = h^1(C, a) = 0$, if $a = 0$ and $a \not\sim \mathcal{O}_C$.
- (c) $h^0(C, a) = h^1(C, a) = 1$, if $a = 0$ and $a \sim \mathcal{O}_C$.

Proof:

- (a) Since $\deg(-a) = -a < 0$ and since K_C is trivial, $h^1(C, a) = h^0(C, -a) = 0$. Thus the Theorem of Riemann-Roch gives

$$h^0(C, a) = \chi(a) = \deg(a) + g(C) - 1 = \deg(a).$$

- (b) and (c) follow from [Har77] IV.1.2. □

G.15 Lemma

Let $D \subset \Sigma$ be an effective divisor such that $D \sim_1 \text{pr}_1^* a + \text{pr}_2^* b$, where $a \in \text{Div}(C_1)$ with $a = \deg(a) \geq 0$ and $b \in \text{Div}(C_2)$ with $b = \deg(b) \geq 0$.

- (a) If $a, b > 0$, then $h^0(\Sigma, D) = ab$ and $h^1(\Sigma, D) = 0$.
- (b) $h^0(\Sigma, D) = h^1(\Sigma, D) \leq \begin{cases} a, & \text{if } a > 0, b = 0 \\ b, & \text{if } a = 0, b > 0. \end{cases}$
- (c) $h^2(\Sigma, D) = 0$.

Proof: We first of all note, that $a, b \geq 0$ and $(a, b) \neq (0, 0)$ by Lemma G.7. Since D is effective, $0 = h^0(\Sigma, -D) = h^2(\Sigma, K_\Sigma + D) = h^2(\Sigma, D)$.

By Lemma G.8 we know that

$$h^0(\Sigma, D) = h^0(C_2, a) \cdot h^0(C_1, b)$$

and

$$h^1(\Sigma, D) = h^1(C_2, a) \cdot h^0(C_1, b) + h^0(C_2, a) \cdot h^1(C_1, b).$$

The result thus follows from Lemma G.14. □

G.16 Lemma

Let $B \subset \Sigma$ be an irreducible curve, $B \not\sim_a C_k$, $k = 1, 2$, and $\{i, j\} = \{1, 2\}$.

- (a) If $B^2 = 0$, then B is smooth, $g(B) = 1$, and $\text{pr}_{ij} : B \rightarrow C_i$ is an unramified covering of degree $B \cdot C_j$.
- (b) If $B^2 = 0$, then $\#(B \cap \tau_p(C_i)) = B \cdot C_j$ for any $p \in \Sigma$, and if $q, q' \in B$, then $\tau_{q-q'}(B) = B$.
- (c) If $B^2 = 0$, then the base curve H in the fibration $\pi : \Sigma \rightarrow H$ with fibre B , which exists according to Proposition B.1, is an elliptic curve.
- (d) If $B \cdot C_i = 1$, then $B^2 = 0$ and $C_j \cong B$.

- (e) If $B.C_i = 1 = B.C_j$, then $C_1 \cong C_2$.
- (f) If B is the graph of a morphism $\alpha : C_i \rightarrow C_j$, then $B.C_j = 1$ and $B^2 = 0$.

Proof:

- (a) The adjunction formula gives

$$p_a(B) = 1 + \frac{B^2 + K_\Sigma \cdot B}{2} = 1.$$

Since $|C_2|_a$ covers the whole of Σ and $B \not\sim_a C_2$, the two irreducible curves B and C_2 must intersect properly, that is, B is not a fibre of pr_1 . But then the mapping $\text{pr}_{1|} : B \rightarrow C_1$ is a finite surjective morphism of degree $B.C_2$. If B was a singular curve its normalisation would have to have arithmetical genus 0 and the composition of the normalisation with $\text{pr}_{1|}$ would give rise to a surjective morphism from \mathbb{P}_C^1 to an elliptic curve, contradicting Hurwitz's formula. Hence, B is smooth and $g(B) = p_a(B) = 1$. We thus may apply the formula of Hurwitz to $\text{pr}_{1|}$ and the degree of the ramification divisor R turns out to be

$$\deg(R) = 2 \cdot \left(g(B) - 1 + (g(C_1) - 1) \cdot \deg(\text{pr}_{1|}) \right) = 0.$$

The remaining case is treated analogously.

- (b) W. l. o. g. $i = 2$. For $p = (p_1, p_2) \in \Sigma$ we have $\tau_p(C_2) = \text{pr}_1^{-1}(p_1)$ is a fibre of pr_1 , and since $\text{pr}_{1|}$ is unramified, $\#(B \cap \tau_p(C_2)) = \deg(\text{pr}_{1|}) = B.C_2$. Suppose $q, q' \in B$ with $\tau_{q-q'}(B) \neq B$. Then $q = \tau_{q-q'}(q') \in B \cap \tau_{q-q'}(B)$, and hence $B^2 = B \cdot \tau_{q-q'}(B) > 0$, which contradicts the assumption $B^2 = 0$.
- (c) Since $\chi(\Sigma) = 0$, [FrM94] Lemma I.3.18 and Proposition I.3.22 imply that $g(H) = p_g(\Sigma) = h^0(\Sigma, K_\Sigma) = 1$.
- (d) W. l. o. g. $B.C_2 = 1$. Let $0 \neq p \in C_2$. We claim that $B \cap \tau_p(B) = \emptyset$, and hence $B^2 = B \cdot \tau_p(B) = 0$. Suppose $(a, b) \in B \cap \tau_p(B)$, then there is an $(a', b') \in B$ such that $(a, b) = \tau_p(a', b') = (a', b' + p)$, i. e. $a = a'$ and $b = b' + p$. Hence, $(0, b), (0, b') \in \tau_{-a}(B) \cap C_2$. But, $\tau_{-a}(B).C_2 = B.C_2 = 1$, and thus $b' = b = b' + p$ in contradiction to the choice of p . $C_1 \cong B$ via $\text{pr}_{1|}$ follows from (a).
- (e) By (d) we have $C_1 \cong B \cong C_2$.
- (f) $\text{pr}_{i|} : B \rightarrow C_i$ is an isomorphism (cf. [GrD67] I.5.1.4), and has thus degree one. But $\deg(\text{pr}_{i|}) = B.C_j$. Thus we are done with (d).

□

G.17 Example

- (a) Let $C_1 = C_2 = C = \mathbb{C}/\Lambda$ with $\Lambda = \mathbb{Z} \oplus \tau\mathbb{Z}$, and $\Sigma = C_1 \times C_2 = C \times C$. The Picard number $\rho(\Sigma)$ is then either three or four, depending on whether the group $\text{End}_0(C)$ of endomorphisms of C fixing 0 is just \mathbb{Z} or

larger. Using [Har77] Theorem IV.4.19 and Exercise IV.4.11 we find the following classification.

Case 1: $\exists d \in \mathbb{N}$ such that $\tau \in \mathbb{Q}[\sqrt{-d}]$, i. e. $\mathbb{Z} \subsetneq \text{End}_0(\mathbb{C})$.

Then $\rho(\Sigma) = 4$ and $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z} \oplus C_3\mathbb{Z} \oplus C_4\mathbb{Z}$ where C_3 is the diagonal in Σ , i. e. the graph of the identity map from \mathbb{C} to \mathbb{C} , and C_4 is the graph of the morphism $\alpha : \mathbb{C} \rightarrow \mathbb{C} : p \mapsto (b\tau) \cdot p$ of degree $|b\tau|^2$, where $0 \neq b \in \mathbb{N}$ minimal with $b \cdot (\tau + \bar{\tau}) \in \mathbb{Z}$ and $b\tau\bar{\tau} \in \mathbb{Z}$. Setting $a := C_3 \cdot C_4 \geq 1$, the intersection matrix is

$$(C_j \cdot C_k)_{j,k=1,\dots,4} = \begin{pmatrix} 0 & 1 & 1 & |b\tau|^2 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & a \\ |b\tau|^2 & 1 & a & 0 \end{pmatrix}.$$

If e. g. $\tau = i$, then $C_4 = \{(c, ic) \mid c \in \mathbb{C}\}$ and

$$(C_j \cdot C_k)_{j,k=1,\dots,4} = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}.$$

Case 2: $\nexists d \in \mathbb{N}$ such that $\tau \in \mathbb{Q}[\sqrt{-d}]$, i. e. $\mathbb{Z} = \text{End}_0(\mathbb{C})$.

Then $\rho(\Sigma) = 3$ and $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z} \oplus C_3\mathbb{Z}$ where again C_3 is the diagonal in Σ . The intersection matrix in this case is

$$(C_j \cdot C_k)_{j,k=1,2,3} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

- (b) Let $C_1 = \mathbb{C}/\Lambda_1$ and $C_2 = \mathbb{C}/\Lambda_2$ with $\Lambda_1 = \mathbb{Z} \oplus \tau_1\mathbb{Z}$, $\tau_1 = i$, and $\Lambda_2 = \mathbb{Z} \oplus \tau_2\mathbb{Z}$, $\tau_2 = \frac{1}{2}i$. Then $C_1 \not\cong C_2$.³¹

We consider the surjective morphisms $\alpha_j : C_1 \rightarrow C_2$, $j = 3, 4$, induced by multiplication with the complex numbers $\alpha_3 = 1$ and $\alpha_4 = i$ respectively. Denoting by C_j the graph of α_j , we claim, $C_1 \cdot C_3 = \deg(\alpha_3) = 2$ and $C_1 \cdot C_4 = \deg(\alpha_4) = 2$. α_j being an unramified covering, we can calculate its degree by counting the preimages of 0. If $p = [a + ib] \in \mathbb{C}/\Lambda_1 = C_1$

³¹Suppose $C_1 \cong C_2$, then there are integers $a, b, c, d \in \mathbb{Z}$ with $ad - bc = \pm 1$ such that $\frac{1}{2}i = \frac{ai+b}{ci+d}$ (cf. [Har77] Theorem IV.15B). But this leads to $ai + b = \frac{1}{2}di - \frac{1}{2}c$, i. e. $2a = d$ and $-2b = c$. Inserting these relations in the determinant equation we get $\pm 1 = 2a^2 + 2b^2 = 2(a^2 + b^2)$ which would say that two divides one.

with $0 \leq a, b < 1$, then

$$\begin{aligned}\alpha_3(p) = 0 &\Leftrightarrow a + ib = \alpha_3 \cdot (a + ib) \in \Lambda_2 \\ &\Leftrightarrow \exists r, s \in \mathbb{Z} : a = r \text{ and } b = \frac{1}{2} \cdot s \\ &\Leftrightarrow a = 0 \text{ and } b \in \{0, \frac{1}{2}\}.\end{aligned}$$

and

$$\begin{aligned}\alpha_4(p) = 0 &\Leftrightarrow ia - b = \alpha_4 \cdot (a + ib) \in \Lambda_2 \\ &\Leftrightarrow \exists r, s \in \mathbb{Z} : -b = r \text{ and } a = \frac{1}{2} \cdot s \\ &\Leftrightarrow b = 0 \text{ and } a \in \{0, \frac{1}{2}\}.\end{aligned}$$

Moreover, the graphs C_3 and C_4 intersect only in the point $(0, 0)$ and the intersection is obviously transversal, so $C_3 \cdot C_4 = 1$.

Thus $\Sigma = C_1 \times C_2$ is an example for a product of non-isomorphic elliptic curves with $\rho(\Sigma) = 4$, $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z} \oplus C_3\mathbb{Z} \oplus C_4\mathbb{Z}$, and intersection matrix

$$(C_j \cdot C_k)_{j,k=1,\dots,4} = \begin{pmatrix} 0 & 1 & 2 & 2 \\ 1 & 0 & 1 & 1 \\ 2 & 1 & 0 & 1 \\ 2 & 1 & 1 & 0 \end{pmatrix}.$$

- (c) See [HuR98] p. 4 for examples $\Sigma = C_1 \times C_2$ with $\rho(\Sigma) = 3$ and intersection matrix

$$\begin{pmatrix} 0 & 1 & a \\ 1 & 0 & 1 \\ a & 1 & 0 \end{pmatrix}, \quad a \neq 1.$$

G.18 Remark

- (a) If C_1 and C_2 are isogenous, then there are irreducible curves $B \subset \Sigma$ with $B \cdot C_i$ arbitrarily large.

For this just note, that we have a curve $\Gamma \subset \Sigma$ which is the graph of an isogeny $\alpha : C_1 \rightarrow C_2$. Denoting by $n_{C_2} : C_2 \rightarrow C_2$ the morphism induced by the multiplication with $n \in \mathbb{N}$, we have a morphism $n_{C_2} \circ \alpha$ whose degree is just $n^2 \cdot \deg(\alpha)$. But the degree is the intersection number of the graph with C_1 . The dual morphism of $n_{C_2} \circ \alpha$ has the the same degree, which then is the intersection multiplicity of its graph with C_2 . (Cf. [Har77] Ex. IV.4.7.)

- (b) If C_1 and C_2 are isogenous, then Σ might very well contain smooth irreducible elliptic curves B which are neither isomorphic to C_1 nor to C_2 , and hence cannot be the graph of an isogeny between C_1 and C_2 . But being an elliptic curve we have $B^2 = 0$ by the adjunction formula. If now $\text{NS}(\Sigma) = \bigoplus_{i=1}^{\rho(\Sigma)} C_i\mathbb{Z}$, where the additional generators are graphs, then $B \sim_a \sum_{i=1}^{\rho(\Sigma)} n_i C_i$ with some $n_i < 0$. (Cf. [LaB92] Ex. 10.6.)

Throughout the remaining part of the subsection we will restrict our attention to the general case, that is that C_1 and C_2 are *not* isogenous. This makes the formulae look much nicer, since then $\text{NS}(\Sigma) = C_1\mathbb{Z} \oplus C_2\mathbb{Z}$.

G.19 Lemma

Let C_1 and C_2 be non-isogenous elliptic curves, $D \in \text{Div}(\Sigma)$ with $D \sim_a aC_1 + bC_2$.

- (a) D is nef if and only if $a, b \geq 0$.
- (b) D is ample if and only if $a, b > 0$.
- (c) D is very ample if and only if $a, b \geq 3$.
- (d) $D^2 = 0$ if and only if $a = 0$ or $b = 0$.
- (e) If D is an irreducible curve, then we are in one of the following cases:
 - (1) $a = 0$ and $b = 1$,
 - (2) $a = 1$ and $b = 0$,
 - (3) $a, b > 0$,

and, conversely, if we are in one of these cases, then there is an irreducible curve algebraically equivalent to D .

- (f) If D is an irreducible curve and $D^2 = 0$, then either $D \sim_a C_1$ or $D \sim_a C_2$.

Proof: (a) and (b) follow from Lemma G.7.

- (c) In Lemma G.7 we proved that $a, b \geq 3$ if D is very ample.

Conversely, if $a < 3$, then $D \cap C_2$ is a divisor of degree $D.C_2 = a < 3$ on the elliptic curve C_2 and hence not very ample (cf. [Har77] Example IV.3.3.3). But then D is not very ample. Analogously if $b < 3$.

- (d) $0 = D^2 = 2ab$ if and only if $a = 0$ or $b = 0$.

- (e) Let us first consider the case that D is irreducible.

If $a = 0$ or $b = 0$, then D is algebraically equivalent to a multiple of a fibre of one of the projections pr_i , $i = 1, 2$. In this situation $D^2 = 0$ and thus the irreducible curve D does not intersect any of the fibres properly. Hence it must be a union of several fibres, and being irreducible it must be a fibre. That is we are in one of the first two cases.

Suppose now that $a, b \neq 0$. Thus D intersects C_i properly, and $0 < D.C_1 = b$ and $0 < D.C_2 = a$.

It now remains to show that the mentioned algebraic systems contain irreducible curves, which is clear for the first two of them. Let therefore a and b be positive. Then obviously the linear system $|aC_1 + bC_2|_l$ contains no fixed component, and being ample by (v) its general element is irreducible according to [LaB92] Theorem 4.3.5.

- (f) Follows from (d) and (e).

□

G.d. Surfaces in $\mathbb{P}_\mathbb{C}^3$ and Complete Intersections

A smooth projective surface Σ in $\mathbb{P}_\mathbb{C}^3$ is given by a single equation $f = 0$ with $f \in \mathbb{C}[w, x, y, z]$ homogeneous, and by definition the degree of Σ , say n , is just the degree of f – for $n = 1$, $\Sigma \cong \mathbb{P}_\mathbb{C}^2$; for $n = 2$, $\Sigma \cong \mathbb{P}_\mathbb{C}^1 \times \mathbb{P}_\mathbb{C}^1$; and for $n = 3$, Σ is isomorphic to $\mathbb{P}_\mathbb{C}^2$ blown up in six points in general position. Thus the Picard number $\rho(\Sigma)$, i. e. the rank of the Néron–Severi group, in these cases is 1, 2, and 7 respectively.³² Note that these are also precisely the cases where Σ is rational.

In general the Picard number $\rho(\Sigma)$ of a surface in $\mathbb{P}_\mathbb{C}^3$ may be arbitrarily large,³³ but the Néron–Severi group always contains a very special member, namely the class $H \in \text{NS}(\Sigma)$ of a hyperplane section with $H^2 = n$. And the class of the canonical divisor is then just $(n - 4) \cdot H$. Moreover, if the degree of Σ is at least four, that is, if Σ is not rational, then it is likely that $\text{NS}(\Sigma) = H\mathbb{Z}$. More precisely, if $n \geq 4$, Noether’s Theorem says that $\{\Sigma \mid \rho(\Sigma) = 1, \deg(\Sigma) = n\}$ is a very general subset of the projective space of smooth projective surfaces in $\mathbb{P}_\mathbb{C}^3$ of fixed degree n , i. e. its complement is an at most countable union of lower dimensional subvarieties (cf. [Har75] Corollary 3.5 or [IsS96] p. 146).

In Chapter V we need to know that hypersurface sections on surfaces in $\mathbb{P}_\mathbb{C}^3$, or more generally on complete intersections, are always non-special. If $\Sigma = H_1 \cap \dots \cap H_{N-2} \subseteq \mathbb{P}_\mathbb{C}^N$ is a complete intersection of hypersurfaces $H_i \subset \mathbb{P}_\mathbb{C}^N$ of degree d_i , $i = 1, \dots, N - 2$, with $d_1 \geq \dots \geq d_{N-2} > 1$, we say Σ is a *complete intersection of type* (d_1, \dots, d_{N-2}) .

G.20 Proposition

Let $\Sigma = H_1 \cap \dots \cap H_{N-2} \subseteq \mathbb{P}_\mathbb{C}^N$ be a smooth complete intersection of type (d_1, \dots, d_{N-2}) , and let $H \subset \Sigma$ be a hyperplane section.

Then $K_\Sigma \sim_l (-N - 1 + \sum_{i=1}^{N-2} d_i) \cdot H$, $H^2 = d_1 \cdots d_{N-2}$ and³⁴ for $d \in \mathbb{Z}$

$$(a) \quad h^1(\Sigma, dH) = 0,$$

$$(b) \quad h^0(\Sigma, dH) = \binom{N+d}{N} + \sum_{k=1}^{N-2} \sum_{1 \leq j_1 < \dots < j_k \leq N-2} (-1)^k \cdot \binom{N+d - \sum_{i=1}^k d_{j_i}}{N}, \text{ and}$$

$$(c) \quad \chi(\mathcal{O}_\Sigma) = 1 + \binom{-1 + \sum_{i=1}^{N-2} d_i}{N} + \sum_{k=1}^{N-2} \sum_{1 \leq j_1 < \dots < j_k \leq N-2} (-1)^k \cdot \binom{-1 + \sum_{i=1}^{N-2} d_i - \sum_{l=1}^k d_{j_l}}{N}.$$

Proof: Denoting by H^i a hyperplane section in $\Sigma_i = H_1 \cap \dots \cap H_i$, $i = 0, \dots, N - 2$, we claim that we have indeed $K_{\Sigma_i} \sim_l (-N - 1 + \sum_{k=1}^i d_k) \cdot H^i$ and for $d \in \mathbb{Z}$

$$(a') \quad h^k(\Sigma_i, dH^i) = 0 \text{ for all } 0 < k < N - i.$$

³²See e. g. [Har77] Example II.8.20.3 and Remark V.4.7.1.

³³E. g. the n -th Fermat surface, given by $w^n + x^n + y^n + z^n = 0$ has Picard number $\rho \geq 3 \cdot (n - 1) \cdot (n - 2) + 1$, with equality if $\gcd(n, 6) = 1$ (cf. [Shi82] Theorem 7, see also [AoS83] pp. 1f. and [IsS96] p. 146).

³⁴If $m < n$, we set the binomial coefficient $\binom{m}{n} = 0$.

$$\begin{aligned}
\text{(b')} \quad h^0(\Sigma_i, dH^i) &= h^0(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d)) + \sum_{k=1}^i \sum_{1 \leq j_1 < \dots < j_k \leq i} (-1)^k \cdot h^0\left(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d - \sum_{l=1}^k d_{j_l})\right) \\
&= \binom{N+d}{N} + \sum_{k=1}^i \sum_{1 \leq j_1 < \dots < j_k \leq i} (-1)^k \cdot \binom{N+d - \sum_{l=1}^k d_{j_l}}{N}.
\end{aligned}$$

We do the proof by induction on i , where for $i = 0$ we are in the case $\Sigma_0 = \mathbb{P}_\mathbb{C}^N$ with $K_{\mathbb{P}_\mathbb{C}^N} \sim_{\mathbb{1}} (-N - 1)H^0$ and (a') follows from [Har77] III.5.1, while (b') is obvious. We may thus assume that $i \geq 1$.

The inclusion $\Sigma_i \hookrightarrow \Sigma_{i-1}$ gives rise to the exact sequence

$$0 \longrightarrow \mathcal{O}_{\Sigma_{i-1}}(-d_i H^{i-1}) \xrightarrow{\cdot H^i} \mathcal{O}_{\Sigma_{i-1}} \longrightarrow \mathcal{O}_{\Sigma_i} \longrightarrow 0, \quad (\text{G.9})$$

and by [Har77] II.8.20 $\omega_{\Sigma_i} \cong \omega_{\Sigma_{i-1}} \otimes \mathcal{O}_{\Sigma_i}(d_i H^i)$. But then by induction

$$K_{\Sigma_i} = \left(-N - 1 + \sum_{k=1}^i d_k \right) \cdot H^i.$$

The long exact cohomology sequence of (G.9) twisted by dH^{i-1} then gives

$$\begin{array}{ccc}
0 \rightarrow H^0(\Sigma_{i-1}, (d - d_i)H^{i-1}) \rightarrow H^0(\Sigma_{i-1}, dH^{i-1}) \longrightarrow H^0(\Sigma_i, dH^i) & (\text{G.10}) \\
& & \downarrow \\
& & H^1(\Sigma_{i-1}, (d - d_i)H^{i-1})
\end{array}$$

and for any $k \geq 1$

$$H^k(\Sigma_{i-1}, dH^{i-1}) \longrightarrow H^k(\Sigma_i, dH^i) \longrightarrow H^{k+1}(\Sigma_{i-1}, (d - d_i)H^{i-1}). \quad (\text{G.11})$$

(a') If $0 < k < N - i$, then $h^k(\Sigma_{i-1}, dH^{i-1}) = 0 = h^{k+1}(\Sigma_{i-1}, (d - d_i) \cdot H^{i-1})$ for any d by induction. Thus by G.11 also $h^k(\Sigma_i, dH^i) = 0$.

(b') Since $h^1(\Sigma_i, dH^i) = 0$, by (G.10) and induction we get

$$\begin{aligned}
h^0(\Sigma_i, dH^i) &= h^0(\Sigma_{i-1}, dH^{i-1}) + h^0(\Sigma_{i-1}, (d - d_i) \cdot H^{i-1}) \\
&= h^0(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d)) + \sum_{k=1}^{i-1} \sum_{1 \leq j_1 < \dots < j_k \leq i-1} (-1)^k \cdot h^0\left(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d - \sum_{l=1}^k d_{j_l})\right) \\
&\quad - h^0(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d - d_i)) - \sum_{k=1}^{i-1} \sum_{1 \leq j_1 < \dots < j_k \leq i-1} (-1)^k \cdot h^0\left(\mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d - d_i - \sum_{l=1}^k d_{j_l})\right). \\
&= h^0(\mathbb{P}_\mathbb{C}^N, \mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d)) + \sum_{k=1}^i \sum_{1 \leq j_1 < \dots < j_k \leq i} h^0\left(\mathbb{P}_\mathbb{C}^N, \mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(d - \sum_{l=1}^k d_{j_l})\right).
\end{aligned}$$

Knowing that for any $\delta \in \mathbb{Z}$ the dimension $h^0(\mathbb{P}_\mathbb{C}^N, \mathcal{O}_{\mathbb{P}_\mathbb{C}^N}(\delta)) = \binom{N+\delta}{N}$ we have proved (b').

(c) This follows immediately by (a), (b) and Serre duality, since

$$\chi(\mathcal{O}_\Sigma) = 1 + h^0(\Sigma, K_\Sigma).$$

Finally, we note that H^2 is just the degree of $\Sigma \subseteq \mathbb{P}_\mathbb{C}^N$, which is just $d_1 \cdots d_{N-2}$ by the Theorem of Bézout (cf. [Har77] I.7.7.). \square

We note that complete intersections of type (4), (3,2) and (2,2,2) are the only smooth complete intersection surfaces which are K3-surfaces, since for no other choice of the d_i and N we can achieve that $\sum_{i=1}^{N-2} d_i - N - 1 = 0$.

Since a smooth hypersurface in $\mathbb{P}_\mathbb{C}^3$ is a complete intersection, we get the following corollary.

G.21 Corollary

Let $\Sigma \subset \mathbb{P}_\mathbb{C}^3$ be a smooth hypersurface of degree n in $\mathbb{P}_\mathbb{C}^3$, and let $H \subset \mathbb{P}_\mathbb{C}^3$ be any hyperplane section.

Then $K_\Sigma \sim_{\mathbb{L}} (n-4) \cdot H$ and³⁵ for $d \in \mathbb{Z}$

- (a) $h^1(\Sigma, dH) = 0$,
- (b) $h^0(\Sigma, dH) = \binom{d+3}{3} - \binom{d-n+3}{3}$, and
- (c) $\chi(\mathcal{O}_\Sigma) = 1 + \binom{n-1}{3}$.

G.e. K3-Surfaces

We note that if Σ is a K3-surface then the Néron–Severi group $\text{NS}(\Sigma)$ and the Picard group $\text{Pic}(\Sigma)$ of Σ coincide, i. e. $|D|_a = |D|_l$ for every divisor D on Σ . The Picard number $\rho(\Sigma)$ of a K3-surface may take values between 1 and 20, and there are K3-surfaces Σ with $\rho(\Sigma) = \rho$ for each $\rho \in \{1, \dots, 20\}$. The generic case is $\rho(\Sigma) = 1$ – more precisely, the K3-surfaces with $\rho(\Sigma) = \rho$ are parametrised by points of an at most countable union of irreducible varieties of dimension $20 - \rho$ (cf. [IsS96] 12.5 Corollary 4). Moreover, if $\text{NS}(\Sigma) = \mathbb{L} \cdot \mathbb{Z}$, then $\pi_\Sigma = \frac{1}{2} \cdot \mathbb{L}^2$ may be any even positive number (cf. [IsS96] 12.5 Corollary 2). In general, we denote by $\pi_\Sigma = \min \{D^2 \mid D \in \text{NS}(\Sigma), D^2 > 0\}$. E. g. if Σ is a double cover of $\mathbb{P}_\mathbb{C}^2$ branched along a sextic, then $\pi_\Sigma = 1$; if Σ is quartic in $\mathbb{P}_\mathbb{C}^3$, then $\pi_\Sigma = 2$; if Σ is a complete intersection of type (3,2) in $\mathbb{P}_\mathbb{C}^4$, then $\pi_\Sigma = 3$; if Σ is a complete intersection of type (2,2,2) in $\mathbb{P}_\mathbb{C}^5$, then $\pi_\Sigma = 4$ (cf. [IsS96] p. 219). Moreover, the surfaces of these types are each parametrised by a 19-dimensional variety (cf. [Mér85] 5.4), and thus by [IsS96] 12.5 Corollary 2 a generic member must have Picard number one.

An irreducible curve B has self-intersection $B^2 = 0$ if and only if the arithmetical genus of B is one. In that case $|B|_l$ is a pencil of elliptic curves without base points endowing Σ with the structure of an elliptic fibration over $\mathbb{P}_\mathbb{C}^1$ (cf. [Mér85] or Proposition B.1). However, a generic K3-surface does not possess an elliptic fibration.

³⁵If $m < n$, we set the binomial coefficient $\binom{m}{n} = 0$.

Bibliography

- [AGZV85] Victor Arnold, Sabir Gusein-Zade, and Alexander Varchenko, *Singularities of differentiable maps*, vol. I, Birkhäuser, 1985.
- [AlH00] James Alexander and Alexandre Hirschowitz, *An asymptotic vanishing theorem for generic unions of multiple points*, *Inv. Math.* **140** (2000), no. 2, 303–325.
- [AoS83] Noboru Aoki and Tetsuji Shioda, *Generators of the Néron-Severi group of a Fermat surface*, *Arithmetic and Geometry* (Michael Artin et al., ed.), vol. I, *Progress in Mathematics*, no. 35, Birkhäuser, 1983, pp. 1–12.
- [ArC81] Enrico Arbarello and Maurizio Cornalba, *Footnotes to a paper of Beniamino Segre*, *Math. Ann.* **256** (1981), 341–362.
- [ArC83] Enrico Arbarello and Maurizio Cornalba, *A few remarks about the variety of irreducible plane curves of given degree and genus*, *Ann. Sci. École Norm. Sup.* **16** (1983), no. 3, 467–488.
- [AtM69] Michael F. Atiyah and Ian G. MacDonal, *Introduction to commutative algebra*, Addison-Wesley, 1969.
- [Bar93a] Didier Barkats, *Irréductibilité des variétés des courbes planes à noeuds et à cusps*, Preprint no. 364, Univ. de Nice-Sophia-Antipolis, 1993.
- [Bar93b] Didier Barkats, *Non vacuité des variétés des courbes planes à noeuds et à cusps*, Preprint no. 363, Univ. de Nice-Sophia-Antipolis, 1993.
- [Bar98] Didier Barkats, *Étude des variétés des courbes planes à noeuds et à cusps*, *Algebraic Geometry, Papers for the EUROPROJ Conferences held in Catania, Italy, September 1993, and Barcelona, Spain, September 1994* (Peter E. Newstead, ed.), *Lecture Notes Pure Appl. Math.* no. 200, 1998, pp. 25–35.
- [Bea83] Arnaud Beauville, *Complex algebraic surfaces*, *LMS LNS*, no. 68, CUP, 1983.
- [BeS95] Mauro C. Beltrametti and Andrew J. Sommese, *The adjunction theory of complex projective varieties*, Walter de Gruyter, 1995.
- [Bog79] Fedor A. Bogomolov, *Holomorphic tensors and vector bundles on projective varieties*, *Math. USSR Izvestija* **13** (1979), no. 3, 499–555.
- [BPV84] Wolf Barth, Christian Peters, and Antonius Van de Ven, *Compact complex surfaces*, Springer, 1984.
- [BrG81] Bill Bruce and Peter J. Gibblin, *A stratification of the space of plane quartic curves*, *Proc. London Math. Soc.* **442** (1981), 270–298.
- [Bru81] Bill Bruce, *Singularities on a projective hypersurface*, *Bull. LMS* **13** (1981), 47–50.
- [Bru99] Andrea Bruno, *Limit linear series and families of equisingular plane curves*, Preprint, 1999.
- [ChC99] Luca Chiantini and Ciro Ciliberto, *On the Severi variety of surfaces in $\mathbb{P}_{\mathbb{C}}^3$* , *J. of Alg. Geom.* **8** (1999), 67–83.
- [ChR93] Mei-Chou Chang and Ziv Ran, *Divisors on some generic hypersurfaces*, *J. of Diff. Geom.* **38** (1993), 671–678.
- [ChS97] Luca Chiantini and Edoardo Sernesi, *Nodal curves on surfaces of general type*, *Math. Ann.* **307** (1997), 41–56.
- [Cil01] Ciro Ciliberto, *Linear systems with multiple assigned base points*, To appear in: *Proceedings of the Conference Zero Dimensional Schemes and Applications*, Feb. 2000, Naples, 2001.
- [CiM98] Ciro Ciliberto and Rick Miranda, *Degenerations of planar linear systems*, *J. Reine Angew. Math.* **501** (1998), 191–220.

- [Dan96] Vladimir I. Danilov, *Cohomology of algebraic varieties*, Algebraic Geometry, Vol. II (Igor R. Shafarevich, ed.), Encyclopedia of Mathematical Sciences, no. 35, Springer, 1996, pp. 1–126.
- [Dav86] Edward Davis, *Zero-dimensional subschemes of $\mathbb{P}_\mathbb{C}^2$: New applications of Castelnuovo's function*, Ann. Univ. Ferrara **32** (1986), 93–107.
- [Deg87] Alexander Degtyarev, *Isotopy classification of complex plane projective curves of degree five*, Preprint P-3-87, Steklov Math. Inst., Leningrad branch, 1987.
- [Deg95] Alexander Degtyarev, *Quintics in \mathbb{C}^2 with non-abelian fundamental group*, Preprint, Max–Planck–Institut für Mathematik, 1995.
- [DeP00] Theo De Jong and Gerhard Pfister, *Local analytic geometry*, Vieweg, 2000.
- [DiH88] Steven Diaz and Joe Harris, *Ideals associated to deformations of singular plane curves*, Trans. AMS **309** (1988), 433–468.
- [EiL93] Lawrence Ein and Robert Lazarsfeld, *Sheshadri constants on smooth surfaces*, Journées de Géométrie Algébrique d'Orsay: Juillet 1992, Astérisque, no. 218, SMF, 1993, pp. 177–186.
- [Eva99] Laurent Evain, *La fonction de Hilbert de la réunion de 4^h gros points génériques de $\mathbb{P}_\mathbb{C}^2$* , J. Alg. Geom. **8** (1999), 787–796.
- [Fla01] Flaminio Flamini, *Moduli of nodal curves on smooth surfaces of general type*, math.AG/0103205, 2001.
- [FIM01] Flaminio Flamini and C. Madonna, *Geometric linear normality for nodal curves on some projective surfaces*, to appear in: Boll. Un. Mat. Ital. 2001.
- [Fra41] Alfredo Franchetta, *Sulle curve eccezionali di prima specie appartenenti a una superficie algebrica*, Boll. Un. Mat. Ital. **2** (1940-41), no. 3, 28–29.
- [FrM94] Robert Friedman and John W. Morgan, *Smooth four-manifolds and complex surfaces*, vol. 3, Ergebnisse der Mathematik und ihrer Grenzgebiete, no. 27, Springer, 1994.
- [FuP97] William Fulton and Rahul Pandharipande, *Notes on stable maps and quantum cohomology*, Algebraic Geometry, Santa Cruz 1995 (János Kollár et al., ed.), Proceedings of Symposia in Pure Mathematics, no. 62,2, AMS, 1997, pp. 45–96.
- [FuP00] Luis Fuentes and Manuel Pedreira, *The projective theory of ruled surfaces*, math.AG/0006204, June 2000.
- [GD84] Claudine Giacinti-Diebolt, *Variétés des courbes projectives planes de degré et lieu singulier donnés*, Math. Ann. **266** (1984), no. 3, 321–350.
- [Gim87] Alessandro Gimigliano, *On linear systems of plane curves*, Ph.D. thesis, Queen's University, Kingston, Ontario, 1987.
- [GLS97] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *New asymptotics in the geometry of equisingular families of curves*, Int. Math. Res. Not. **13** (1997), 595–611.
- [GLS98a] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *Geometry of families of nodal curves on the blown-up projective plane*, Trans. AMS **350** (1998), 251–274.
- [GLS98b] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *On the irreducibility of families of curves*, Unpublished Manuscript, 1998.
- [GLS98c] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *Plane curves of minimal degree with prescribed singularities*, Inv. math. **133** (1998), 539–580.
- [GLS00] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *Castelnuovo function, zero-dimensional schemes, and singular plane curves*, J. of Alg. Geom. **9** (2000), no. 4, 663–710.
- [GLS02] Gert-Martin Greuel, Christoph Lossen, and Eugenii Shustin, *Singular algebraic curves*, Springer, 2002.

- [GrD67] Alexandre Grothendieck and Jean Dieudonné, *Éléments de Géométrie Algébrique*, vol. I-IV, Publication Mathématiques, no. 4, 8, 11, 17, 20, 24, 28, and 32, IHES, 1960-67.
- [GrH94] Phillip Griffith and Joseph Harris, *Principles of algebraic geometry*, John Wiley, 1994.
- [GrK89] Gert-Martin Greuel and Ulrich Karras, *Families of varieties with prescribed singularities*, Comp. math. **69** (1989), 83–110.
- [GrL96] Gert-Martin Greuel and Christoph Lossen, *Equianalytic and equisingular families of curves on surfaces*, Man. math. **91** (1996), 323–342.
- [GrM85] Monique A. Gradolato and Emilia Mezzetti, *Curves with nodes, cusps and ordinary triple points*, Ann.Univ. Ferrara, sez. 7, **31** (1985), 23–47.
- [GrM88] Monique A. Gradolato and Emilia Mezzetti, *Families of curves with ordinary singular points on regular surfaces*, Ann. mat. pura et appl. **150** (1988), 281–298.
- [GrS99] Gert-Martin Greuel and Eugenii Shustin, *Geometry of equisingular families of curves*, Singularity Theory (Bill Bruce and David Mond, eds.), LMS LNS, no. 263, CUP, 1999, pp. 79–108.
- [GTU66] D. A. Gudkov, M. L. Tai, and G. A. Utkin, *Complete classification of irreducible curves of the 4th order*, Mat. Sbornik **69** (1966), no. 2, 222–256.
- [GuN83] Sabir M. Gusein-Zade and Nikolay N. Nekhoroshev, *Contiguity of A_k -singularities at points of the stratum $\mu = \text{const}$ of a singularity*, Funct. Anal. Appl. **17** (1983), 82–83.
- [GuS84] D. A. Gudkov and Eugenii Shustin, *On the intersection of the close algebraic curves*, Topology (Albrecht Dold and Benno Eckmann, eds.), Lecture Notes in Mathematics, no. 1060, Springer, 1984, pp. 278–289.
- [Har75] Robin Hartshorne, *Equivalence relations on algebraic cycles and subvarieties of small codimension*, Algebraic Geometry, Arcata 1974 (Robin Hartshorne, ed.), AMS Proceedings of Symposia in Pure Mathematics, no. 29, 1975, pp. 129–164.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer, 1977.
- [Har85a] Brian Harbourne, *Complete linear systems on rational surfaces*, Trans. AMS **289** (1985), 213–226.
- [Har85b] Joe Harris, *On the Severi problem*, Inv. Math. **84** (1985), 445–461.
- [Har86] Brian Harbourne, *The geometry of rational surfaces and Hilbert functions of points in the plane*, Can. Math. Soc. Conf. Proc. **6** (1986), 95–111.
- [Har92] Joe Harris, *Algebraic geometry, a first course*, Graduate Texts in Mathematics, no. 133, Springer, 1992.
- [Har98] Joe Harris, *Moduli of curves*, Graduate Texts in Mathematics, no. 187, Springer, 1998.
- [Har01] Brian Harbourne, *Survey on fat points in $\mathbb{P}_{\mathbb{C}}^2$* , math.AG/0101112, 2001.
- [Hir85] Alexandre Hirschowitz, *La méthode d'Horace pour l'interpolation à plusieurs variables*, Man. Math. **50** (1985), 337–388.
- [Hir86] Friedrich Hirzebruch, *Singularities of algebraic surfaces and characteristic numbers*, Contemp. Math. (1986), 141–155.
- [Hir89] Alexandre Hirschowitz, *Une conjecture pour la cohomologie des diviseurs sur les surfaces rationnelles génériques*, J. Reine Angew. Math. **397** (1989), 208–213.
- [Hir92] Atsuko Hirano, *Constructions of plane curves with cusps*, Saitama Math. J. **10** (1992), 21–24.
- [HuR98] Klaus Hulek and Kristian Ranestad, *Abelian surfaces with two plane cubic curve fibrations and Calabi-Yau threefolds*, Preprint Series 17, University of Oslo, 1998.

- [IsS96] Vassilii A. Iskovskih and Igor R. Shafarevich, *Algebraic surfaces*, Algebraic Geometry, Vol. II (Igor R. Shafarevich, ed.), Encyclopedia of Mathematical Sciences, no. 35, Springer, 1996, pp. 127–254.
- [ItS99] Ilia Itenberg and Eugenii Shustin, *Complexification of the Viro theorem and topology of real and complex combinatorial hypersurfaces*, MPI 99-111, 1999.
- [Ivi85] Kestutis Ivinskis, *Normale Flächen und die Miyaoka-Kobayashi-Ungleichung*, Master's thesis, Universität Bonn, 1985.
- [Kan89a] Pyung-Lyun Kang, *A note the variety of plane curves with nodes and cusps*, Proc. AMS **106** (1989), no. 2, 309–312.
- [Kan89b] Pyung-Lyun Kang, *On the variety of plane curves with δ nodes and k cusps*, Trans. AMS **316** (1989), no. 1, 165–192.
- [Kaw82] Yujiro Kawamata, *A generalisation of Kodaira-Ramanujam's vanishing theorem*, Math. Ann. **261** (1982), 43–46.
- [Koe86] Robert Jan Koelman, *Over de cusp*, Master's thesis, Univ. Leiden, 1986.
- [LaB92] Herbert Lange and Christina Birkenhake, *Complex abelian varieties*, Grundlehren der mathematischen Wissenschaften, no. 302, Springer, 1992.
- [Laz97] Robert Lazarsfeld, *Lectures on linear series*, Complex Algebraic Geometry (János Kollár, ed.), IAS/Park City Mathematics Series, no. 3, AMS, 1997, pp. 161–219.
- [Los98] Christoph Lossen, *The geometry of equisingular and equianalytic families of curves on a surface*, Phd thesis, FB Mathematik, Universität Kaiserslautern, Aug. 1998, <http://www.mathematik.uni-kl.de/~wwwagag/preprints/reports/Lossen/Lossen002/Lossen002.ps.gz>.
- [Los99] Christoph Lossen, *New asymptotics for the existence of plane curves with prescribed singularities*, Comm. in Algebra **27** (1999), 3263–3282.
- [Lue87a] Ignacio Luengo, *The μ -constant stratum is not smooth*, Inv. Math. **90** (1987), 139–152.
- [Lue87b] Ignacio Luengo, *On the existence of complete families of plane curves, which are obstructed*, J. LMS **36** (1987), 33–43.
- [Mér85] Jean-Yves Mérindol, *Propriétés élémentaires des surfaces $K3$* , Géométrie des surfaces $K3$: Modules et périodes; Séminaire Palaiseau, Astérisque, no. 126, SMF, 1985, pp. 45–58.
- [Mig00] Thierry Mignon, *Systèmes des courbes planes à singularités imposées: le cas des multiplicités inférieures ou égales à quatre*, J. Pure Appl. Alg. **151** (2000), no. 2, 173–195.
- [Mig01] Thierry Mignon, *An asymptotic existence theorem for plane curves with prescribed singularities*, J. of Alg. Geom. **10** (2001), no. 2, 281–297.
- [Mil68] John Milnor, *Singular points of complex hypersurfaces*, PUP, 1968.
- [Mum66] David Mumford, *Lectures on curves on an algebraic surface*, PUP, 1966.
- [Nag59] Masayoshi Nagata, *On the 14th problem of Hilbert*, Amer. J. Math. **33** (1959), 766–772.
- [Nag60] Masayoshi Nagata, *On rational surfaces, II*, Memoirs of the College of Science, University of Kyoto, Ser. A, **33** (1960), no. 2, 271–293.
- [Nag70] Masayoshi Nagata, *On self-intersection number of a section on a ruled surface*, Nagoya Mathematical Journal **37** (1970), 191–196.
- [Ran89] Ziv Ran, *Families of plane curves and their limits: Enriques' conjecture and beyond*, Annals of Math. **130** (1989), no. 1, 121–157.
- [Roé01] Joaquim Roé, *On the existence of plane curves with imposed multiple points*, J. Pure Appl. Alg. **156** (2001), 115–126.
- [Sak93] Fumio Sakai, *Singularities of plane curves*, Geometry of Complex Projective Varieties, Seminars and Conferences 9. Mediterranean Press, 1993, pp. 257–273.

- [Seg24] Beniamino Segre, *Dei sistemi lineari tangenti ad un qualunque sistema di forme*, Atti Acad. Naz. Lincei Rendiconti, Ser. 5, **33** (1924), 182–185.
- [Seg29] Beniamino Segre, *Esistenza e dimensione di sistemi continui di curve piane algebriche con dati caratteri*, Atti Acad. Naz. Lincei Rendiconti, Ser. 6, **10** (1929), 31–38.
- [Seg62] Beniamino Segre, *Alcune questioni su insiemi finiti di punti in geometria algebrica*, Atti del Convegno Internazionale di Geometria Algebrica, Torino, 1961, 1962, pp. 15–33.
- [Sei99] James Seibert, *The dimension of quasi-homogeneous linear systems with multiplicity four*, math.AG/9905076, 1999.
- [Sev21] Francesco Severi, *Vorlesungen über Algebraische Geometrie*, Bibliotheca Mathematica Teubneriana, no. 32, Teubner, 1921.
- [Shi82] Tetsuji Shioda, *On the Picard number of a Fermat surface*, Journal Fac. Sci. Univ. Tokyo, Sec. IA, **28** (1982), 725–734.
- [ShT96] Eugenii Shustin and Ilya Tyomkin, *Linear systems of plane curves with generic multiple points*, Preprint Tel-Aviv University, 1996.
- [ShT99] Eugenii Shustin and Ilya Tyomkin, *Versal deformation of algebraic hypersurfaces with isolated singularities*, Math. Ann. **313** (1999), 297–314.
- [Shu87a] Eugenii Shustin, *New M-curve of the 8th degree*, Math. Notes of Acad. Sci. USSR **42** (1987), 606–610.
- [Shu87b] Eugenii Shustin, *Versal deformation in the space of plane curves of fixed degree*, Funct. An. Appl. **21** (1987), 82–84.
- [Shu91a] Eugenii Shustin, *Geometry of discriminant and topology of algebraic curves*, Proc. Intern. Congress Math., Kyoto 1990 (Tokyo, Berlin, New York), vol. 1, Springer, 1991.
- [Shu91b] Eugenii Shustin, *On manifolds of singular algebraic curves*, Selecta Math. Sov. **10** (1991), 27–37.
- [Shu93] Eugenii Shustin, *Real plane algebraic curves with prescribed singularities*, Topology **32** (1993), 854–856.
- [Shu94] Eugenii Shustin, *Smoothness and irreducibility of varieties of algebraic curves with nodes and cusps*, Bull. SMF **122** (1994), 235–253.
- [Shu96a] Eugenii Shustin, *Geometry of equisingular families of plane algebraic curves*, J. Alg. Geom. **5** (1996), 209–234.
- [Shu96b] Eugenii Shustin, *Smoothness and irreducibility of varieties of algebraic curves with ordinary singularities*, Israel Math. Conf. Proc., no. 9, AMS, 1996, pp. 393–416.
- [Shu97] Eugenii Shustin, *Smoothness of equisingular families of plane algebraic curves*, Math. Res. Not. **2** (1997), 67–82.
- [Shu98] Eugenii Shustin, *Gluing of singular and critical points*, Topology **37** (1998), no. 1, 195–217.
- [Shu99] Eugenii Shustin, *Lower deformations of isolated hypersurface singularities*, Algebra i Analiz **10** (1999), no. 5, 221–249.
- [Shu01] Eugenii Shustin, *Analytic order of singular and critical points*, Preprint University of Tel Aviv, 2001.
- [Tan80] Allen Tannenbaum, *Families of algebraic curves with nodes*, Comp. Math. **41** (1980), 107–126.
- [Tan82] Allen Tannenbaum, *Families of curves with nodes on K3-surfaces*, Math. Ann. **260** (1982), 239–253.
- [Tan84] Allen Tannenbaum, *On the classical characteristic linear series of plane curves with nodes and cuspidal points: Two examples of Beniamino Segre*, Comp. Math. **51** (1984), 169–183.

- [Ura83] Tohsuke Urabe, *On quartic surfaces and sextic curves with certain singularities*, Proc. Japan. Acad. Ser. A, vol. 59, 1983, pp. 434–437.
- [Ura84] Tohsuke Urabe, *On quartic surfaces and sextic curves with singularities of type $\tilde{E}_8, T_{2,3,7}, \tilde{E}_{12}$* , Publ. Inst. Math. Sci. Kyoto Univ. **80** (1984), 1185–1245.
- [Ura86] Tohsuke Urabe, *Singularities in a certain class of quartic surfaces and sextic curves and Dynkin graphs*, Proc. Vancouver Conf. Algebraic Geometry, July 2–12, 1984, 1986, pp. 4877–497.
- [Var83] Alexander N. Varchenko, *On semicontinuity of the spectrum and an upper estimate for the number of singular points of a projective hypersurface*, Sov. Math. Dokl. **27** (1983), 735–739.
- [Vas90] Victor A. Vassiliev, *Stable cohomology of complements to the discriminants of deformations of singularities of smooth functions*, J. Sov. Math. **52** (1990), 3217–3230.
- [Vie82] Eckart Viehweg, *Vanishing theorems*, Journ. Reine Ang. Math. **335** (1982), 1–8.
- [Wae73] Bartel Leendert van der Waerden, *Einführung in die algebraische Geometrie*, 2. ed., Springer, 1973.
- [Wah74a] Jonathan M. Wahl, *Deformations of plane curves with nodes and cusps*, Amer. J. Math. **96** (1974), 529–577.
- [Wah74b] Jonathan M. Wahl, *Equisingular deformations of plane algebroid curves*, Trans. AMS **193** (1974), 143–170.
- [Wal95] Charles T. C. Wall, *Geometry of quartic curves*, Math. Proc. Camb. Phil. Soc. **117** (1995), 415–423.
- [Wal96] Charles T. C. Wall, *Highly singular quintic curves*, Math. Proc. Camb. Phil. Soc. **119** (1996), 257–277.
- [Xu94] Geng Xu, *Curves in \mathbb{P}^2 and symplectic packings*, Math. Ann. **299** (1994), 609–613.
- [Xu95] Geng Xu, *Ample line bundles on smooth surfaces*, J. reine angew. Math. **469** (1995), 199–209.
- [Zar35] Otto Zariski, *Algebraic surfaces*, vol. III, Ergebnisse der Mathematik und ihrer Grenzgebiete, no. 5, Springer, 1935.

Index

- A_μ , 9
 $B(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$, 10
 D_μ , 9
 E_i , 1
 $H^\vee(X, D)$, 1
 $H^\vee(\mathcal{F})$, 1
 $I(C, T^*)$, 24
 $I^a(C, z)$, 7
 $I^a(f)$, 7
 $I^a(C, z)$, 6
 $I^{ea}(C, z)$, 5
 $I^{ea}(f)$, 5
 $I_{\text{fix}}^{ea}(C, z)$, 6
 $I_{\text{fix}}^{ea}(f)$, 6
 $I^{es}(C, z)$, 6
 $I_{\text{fix}}^{es}(C, z)$, 6
 $I^s(C, z)$, 6
 K_Σ , 1
 M_g , 152
 $T^*(C, z)$, 6, 9, 24
 $V_{B, \underline{n}}$, 16
 $V_{B, n}$, 121
 $V_{D, n}$, 123
 $V_{|D|}(\mathcal{S}_1, \dots, \mathcal{S}_r)$, 9
 $V_{|D|}(\mathfrak{m}_1, \dots, \mathfrak{m}_r)$, 10
 $V_{|D|}^{\text{fix}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$, 9
 $V_{|D|}^{\text{irr, fix}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$, 10
 $V_{|D|}^{\text{irr, reg}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$, 10
 $V_{|D|}^{\text{reg}}(\mathcal{S}_1, \dots, \mathcal{S}_r)$, 9
 $X(C)$, 6
 $X(C, T^*)$, 24
 $X(\underline{m}; \underline{z})$, 5
 $X : L$, 2
 $X \cap L$, 2
 $X^*(C)$, 6
 $X_{\text{fix}}^*(C)$, 6
 $X^a(C)$, 6
 $X^{ea}(C)$, 5
 $X_{\text{fix}}^{ea}(C)$, 6
 $X^{es}(C)$, 6
 $X_{\text{fix}}^{es}(C)$, 6
 $X^s(C)$, 6
 Y_y , 128
 $\#X$, 1
 $\text{Bl}_z(\Sigma)$, 1
 $C[x, y]_a$, 44
 $C[x, y]_{\leq a}^*$, 44
 $C[x, y]_{\leq a}$, 7, 44
 $C[x, y]_{\leq a}^{ea}$, 44
 $C[x, y]_{\leq a}^{es}$, 44
 $\mathcal{C}\ell(C, T^*(C, z))$, 6
 $\mathcal{C}\ell(C, T^*)$, 24
 $\text{Div}(\Sigma)$, 1
 $\text{Hilb}_\Sigma^{\text{red}}$, 1
 $\text{Hilb}(\mathcal{S})$, 8
 Hilb_Σ^n , 140
 Hilb_Σ , 1, 121
 Hilb_Σ^n , 121
 Hilb_Σ^n , 11
 $\text{Hilb}_z(\mathcal{S})$, 8
 $\text{NS}(\Sigma)$, 1
 $\mathbb{P}(\mathcal{E})$, 142
 $\text{Pic}(\Sigma)$, 1
 $\Psi_D(k_1\mathcal{S}_1, \dots, k_r\mathcal{S}_r)$, 11
 Σ , 1
 $\deg(X)$, 1
 $\deg(X^a(\mathcal{S}))$, 9
 $\deg(X^a(\mathcal{S}))$, 8, 9
 $\deg(X^s(\mathcal{S}))$, 8, 9
 $\deg(X^{ea}(\mathcal{S}))$, 8
 $\deg(X_{\text{fix}}^{ea}(\mathcal{S}))$, 8
 $\deg(X^{es}(\mathcal{S}))$, 8
 $\deg(X_{\text{fix}}^{es}(\mathcal{S}))$, 8
 $\deg \mathcal{I}(f)$, 7
 $\delta(C, z)$, 3
 $\delta(\mathcal{S})$, 3
 $\dim \text{Hilb}(\mathcal{S})$, 8
 $\dim \text{Hilb}_z(\mathcal{S})$, 8
 $\text{jet}_m(g)$, 7
 $\kappa(C, z)$, 3
 $\kappa(\mathcal{S})$, 3
 $\mathcal{I}^a(f)$, 7
 $\mathcal{J}_X : \mathcal{J}_L$, 2
 $\mathcal{J}_X = \mathcal{J}_{X/Y}$, 1

- $\mathcal{I}_{X(C, T^*)/\Sigma}$, [24](#)
 $\mathcal{I}_{X(\underline{m}; \underline{z})}$, [5](#)
 $\mathcal{I}_{X^a(C)/\Sigma}$, [6](#), [7](#)
 $\mathcal{I}_{X^{ea}(C)/\Sigma}$, [5](#), [7](#)
 $\mathcal{I}_{X_{fix}^{ea}(C)/\Sigma}$, [6](#)
 $\mathcal{I}_{X^{es}(C)/\Sigma}$, [6](#)
 $\mathcal{I}_{X^{es}(C)}$, [7](#)
 $\mathcal{I}_{X_{fix}^{es}(C)/\Sigma}$, [6](#)
 $\mathcal{I}_{X^s(C)/\Sigma}$, [6](#)
 $\mathcal{I}_{X^s(C)}$, [7](#)
 $\mathcal{O}_X(D)$, [1](#)
 $\mathcal{O}_{Y, z}$, [1](#)
 \mathcal{S} , [2](#)
 $m_{Y, z}$, [1](#)
 ϵ , [142](#)
 $|D|$, [1](#)
 $|D|_a$, [1](#)
 $|D|_l$, [1](#)
 $|D|_n$, [1](#)
 $\text{mod}(C, z)$, [3](#)
 $\text{mod}(\mathcal{S})$, [3](#)
 $\mu(C, z)$, [3](#)
 $\mu(\mathcal{S})$, [3](#)
 $\text{mult}(C, z)$, [3](#)
 $\text{mult}(X, z)$, [1](#)
 $\text{mult}(\mathcal{S})$, [3](#)
 $\nu^*(\mathcal{S})$, [3](#)
 $\nu^a(\mathcal{S})$, [3](#)
 $\nu^s(\mathcal{S})$, [3](#)
 $\text{pr}_i : \Sigma \rightarrow C_i$, [150](#)
 $\psi(k_1 \mathcal{S}_1, \dots, k_r \mathcal{S}_r)$, [11](#)
 \sim , [25](#)
 \sim_a , [1](#)
 \sim_c , [2](#)
 \sim_l , [1](#)
 \sim_n , [1](#)
 \sim_t , [2](#)
 $\text{supp}(X)$, [1](#)
 $\tau(C, z)$, [3](#)
 $\tau(\mathcal{S})$, [3](#)
 $\tau^*(\mathcal{S})$, [3](#), [44](#), [69](#)
 $\tau^{es}(\mathcal{S})$, [3](#)
 \tilde{C} , [1](#)
 $\tilde{I}^a(g)$, [7](#)
 $\tilde{\Sigma}$, [1](#)
 $\tilde{\mathcal{I}}(f)$, [7](#)
 e , [142](#)
 $e^*(\mathcal{S})$, [45](#)
 $e^a(\mathcal{S})$, [45](#)
 $e^s(\mathcal{S})$, [45](#)
 f_k , [44](#)
 $g(C)$, [1](#)
 $p_a(C)$, [1](#)
 $r(C, z)$, [3](#)
 $r(\mathcal{S})$, [3](#)

abelian variety, [154](#)
asymptotical behaviour, [62](#), [81](#)
asymptotically proper, [II](#), [39](#)

Bogomolov, [61](#), [63](#), [64](#), [82](#), [83](#), [85](#), [89](#), [91](#)

complete embedded resolution tree, [9](#)
Conjecture
 of Gimigliano, [38](#)
 of Harbourne–Hirschowitz, [38](#)
 of Nagata, [37](#), [38](#)
 of Severi, [I](#)
conversation of Milnor numbers, [51](#)
Cremona transformation, [125](#)
curve, [1](#)
 irreducible, [1](#)
 reduced, [1](#)

deformation determinacy
 analytical, [3](#)
 topological, [3](#)
degree
 of $\mathcal{I}(f)$, [7](#)
 of $X^a(\mathcal{S})$, [8](#), [9](#)
 of $X^s(\mathcal{S})$, [8](#), [9](#)
 of $X^{ea}(\mathcal{S})$, [8](#)
 of $X_{fix}^{ea}(\mathcal{S})$, [8](#)
 of $X^{es}(\mathcal{S})$, [8](#)
 of $X_{fix}^{es}(\mathcal{S})$, [8](#)
 of a divisor on a curve, [137](#)
 of a zero-dimensional scheme, [1](#)
delta invariant, [3](#)
determinacy bound, [49](#)
divisor
 algebraically equivalent, [1](#)
 canonical, [1](#)
 exceptional, [1](#)
 linearly equivalent, [1](#)
 numerically equivalent, [1](#)
 very ample, [138](#)

elliptic curve, [154](#)
embedded resolution tree, [6](#), [24](#)
equimultiple family, [25](#)
equisingular families of curves, [10](#)

- equisingularity ideal, 6
- Example of Kodaira, 125
- examples
 - \mathbb{P}_c^2 , 20, 27, 33, 38, 54, 62, 82, 110
 - complete intersection K3-surfaces, 162
 - complete intersections, 27, 34, 61, 63, 116, 160
 - geometrically ruled surfaces, 20, 27, 33, 55, 111, 142
 - hypersurfaces in \mathbb{P}_c^n , 38
 - K3-surfaces, 22, 27, 35, 59, 61, 118, 162
 - of elliptic curves with Picard number 3 and 4, 156
 - products of curves, 21, 27, 34, 56, 114, 150
 - products of elliptic curves, 21, 58, 116, 154
 - rational surfaces, 62, 82, 111, 125
 - rational surfaces in \mathbb{P}_c^3 , 160
 - surface with infinitely many exceptional curves, 125
 - surfaces in \mathbb{P}_c^3 , 22, 27, 34, 58, 61, 81, 117, 160
- genus
 - arithmetical, 1
 - geometrical, 1
- genus formula, 39
- geometrically ruled surface, 142
 - normalised bundle, 142
- Hilbert scheme
 - of curves on a surface, 1, 121
 - of curves on a surface with fixed Hilbert polynomial, 121
 - of singularity schemes, 8
 - of zero-dimensional schemes of degree n on a surface, 11, 140
- Hirzebruch surface, 27, 28, 33, 72, 73, 111, 144
- Hodge Index Theorem, 139
- in very general position, 2, 121
- invariants
 - analytical, 3
 - topological, 3
- isogenous, 154
- Lemma
 - of Geng Xu, 16
 - of Zariski, 131
- Milnor number, 3
- modality, 3
- moduli space of curves, 152
- multiplicity, 3
 - of X at z , 1
- Néron–Severi group, 1
- period, 154
- Picard group, 1
- Principle of Connectedness, 129, 130
- reduction resistant condition
 - on \mathcal{K} , 25
 - suitable, 26
- regular with respect to L , 27
- residue scheme, 2
- Severi variety, 37, 61, 78, 81
- singularity
 - analytically equivalent, 2
 - contact equivalent, 2
 - finitely determined, 49
 - plane curve, 2
 - topologically equivalent, 2
- singularity ideal, 6
 - analytical, 6
- singularity scheme, 6, 24
 - analytical, 6
 - equianalytical, 5
 - equisingularity, 6
 - generalised, 24
 - equivalent, 25
 - isomorphic, 24
 - ordinary fat point scheme, 5, 40, 43
- singularity type
 - analytical, 2
 - good representative, 44
 - topological, 2, 52, 53
- suitable collection of ideals, 7
- support, 1
- T-smooth, 10, 44, 61
- Theorem
 - Existence, 40, 43, 46, 52–59
 - Irreducibility, 84–88, 111, 112, 114, 116–118
 - Kodaira Vanishing Theorem, 27
 - of Noether, 117, 160
 - T-Smoothness, 63, 70, 72–79
 - Vanishing, 13, 15, 20–23, 33–35

Tjurina ideal, [5](#)

Tjurina number, [3](#)

trace scheme, [2](#)

very general, [2](#), [121](#), 123, 124, 140, 152

very general position, 16

Viro method, 38, 46

Wissenschaftlicher Werdegang

1988	Abitur (St. Josef Gymnasium Biesdorf / Eifel)
1989–1992 und 1993–1996	Studium der Fächer Mathematik, Geschichte und Biologie an der Johannes Gutenberg–Universität Mainz
1990–1996	Stipendiat der Studienstiftung des deutschen Volkes
1991	Zwischenprüfung in Mathematik und Geschichte (Johannes Gutenberg–Universität Mainz)
1992	Zwischenprüfung in Biologie (Johannes Gutenberg–Universität Mainz)
1992–1993	Studium des Faches Mathematik an der University of Warwick / Großbritannien
1993	Master of Science in Pure Mathematics (University of Warwick / Großbritannien)
1996	Staatsexamen in Mathematik, Geschichte und Biologie (Johannes Gutenberg–Universität Mainz)
seit 1996	wissenschaftlicher Angestellter der Universität Kaiserslautern