



Annual net primary productivity of a cyanobacteria-dominated biological soil crust in the Gulf Savannah, Queensland, Australia

Burkhard Büdel¹, Wendy J. Williams², and Hans Reichenberger¹

¹Plant Ecology and Systematics, University of Kaiserslautern, 67663 Kaiserslautern, Germany

²Arid Soil Ecosystems, Agriculture and Food Sciences, University of Queensland, Gatton, 4343, Australia

Correspondence: Burkhard Büdel (buedel@bio.uni-kl.de)

Received: 31 August 2017 – Discussion started: 15 September 2017

Revised: 7 December 2017 – Accepted: 8 December 2017 – Published: 26 January 2018

Abstract. Biological soil crusts (biocrusts) are a common element of the Queensland (Australia) dry savannah ecosystem and are composed of cyanobacteria, algae, lichens, bryophytes, fungi and heterotrophic bacteria. Here we report how the CO₂ gas exchange of the cyanobacteria-dominated biocrust type from Boodjamulla National Park in the north Queensland Gulf Savannah responds to the pronounced climatic seasonality and on their quality as a carbon sink using a semi-automatic cuvette system. The dominant cyanobacteria are the filamentous species *Symplocastrum purpurascens* together with *Scytonema* sp. Metabolic activity was recorded between 1 July 2010 and 30 June 2011, during which CO₂ exchange was only evident from November 2010 until mid-April 2011, representative of 23.6 % of the 1-year recording period. In November at the onset of the wet season, the first month (November) and the last month (April) of activity had pronounced respiratory loss of CO₂. The metabolic active period accounted for 25 % of the wet season and of that period 48.6 % was net photosynthesis (NP) and 51.4 % dark respiration (DR). During the time of NP, net photosynthetic uptake of CO₂ during daylight hours was reduced by 32.6 % due to water supersaturation. In total, the biocrust fixed 229.09 mmol CO₂ m⁻² yr⁻¹, corresponding to an annual carbon gain of 2.75 g m⁻² yr⁻¹. Due to malfunction of the automatic cuvette system, data from September and October 2010 together with some days in November and December 2010 could not be analysed for NP and DR. Based on climatic and gas exchange data from November 2010, an estimated loss of 88 mmol CO₂ m⁻² was found for the 2 months, resulting in corrected annual rates of 143.1 mmol CO₂ m⁻² yr⁻¹, equivalent to a carbon gain of 1.7 g m⁻² yr⁻¹. The bulk of the net photosynthetic activity

occurred above a relative humidity of 42 %, indicating a suitable climatic combination of temperature, water availability and light intensity well above 200 μmol photons m⁻² s⁻¹ photosynthetic active radiation. The Boodjamulla biocrust exhibited high seasonal variability in CO₂ gas exchange pattern, clearly divided into metabolically inactive winter months and active summer months. The metabolic active period commences with a period (of up to 3 months) of carbon loss, likely due to reestablishment of the crust structure and restoration of NP prior to about a 4-month period of net carbon gain. In the Gulf Savannah biocrust system, seasonality over the year investigated showed that only a minority of the year is actually suitable for biocrust growth and thus has a small window for potential contribution to soil organic matter.

1 Introduction

Biological soil crusts (named “biocrusts” throughout the text) are a consortium of heterotrophic bacteria, cyanobacteria, algae, fungi, lichens and bryophytes in different proportions with photoautotrophic organisms dominating their biomass. They cover dryland soil surfaces and can compose up to 70 % of a dryland ecosystem’s living cover (Belnap, 1995; Belnap et al., 2016), but also occur in other climatic regions where competition with vascular plants is low (Büdel, 2001; Büdel et al., 2014). Due to the poikilohydric character of biocrust organisms, biocrusts exhibit high resilience under extreme conditions and have a remarkable adaptation to various combinations of climatic factors (e.g. Karsten et al., 2016; Sancho et al., 2016, and citations

therein), thus making them excellent candidates for pioneering hostile environments on our planet. There is good evidence that cyanobacteria-dominated biocrusts have inhabited Earth's soil surfaces for at least 2600 million years (Watanabe et al., 2000; for an overview see also Beraldi-Campesi and Retallack, 2016). Lalonde and Konhauser (2015) point to the importance of oxygenic photosynthesis of early biocrusts providing sufficient equivalents for oxidative-weathering reactions in benthic and soil environments. This certainly also points to the role of biocrusts in soil formation and soil fertility, for example, in leaching carbon and nitrogen to initial soils. Consequently, there is growing interest in biocrust carbon gain (Lange and Belnap, 2016) and their CO₂ exchange rates are considered relevant on local and global scales (e.g. Castillo-Monroy et al. 2011; Wilske et al., 2009; Elbert et al., 2012; Porada et al., 2013, 2014). Process-based models as used by Porada et al. (2013, 2014) still rely on a few available data sets covering a small number of biocrust types, organisms, geographical regions and climatic conditions (see also summary in Sancho et al., 2016).

With the focus on biocrust CO₂ gas exchange over longer periods of time, a number of studies have been published either on the basis of long-term measurements or modelled from single or grouped measurements. From these results, two biocrust groups can be distinguished: one where biocrusts experienced net C uptake and the other where biocrusts experienced C loss. Examples from the net C uptake group include a biocrust from the Mojave Desert which gained 11.7 g C m⁻² yr⁻¹ (Brostoff et al., 2005), a biocrust in the northern Negev Desert, Israel, with a net C uptake of 0.7–5.1 g m⁻² yr⁻¹ (Wilske et al., 2008, 2009) and a biocrust from a desert region of north-west China showing a net C uptake of 3.5 to 6.1 g C m⁻² yr⁻¹ (Feng et al., 2014). Among the C-losing biocrusts are those of south-east Utah determined to be typical net C sources (Bowling et al., 2011), a biocrust from the Colorado Plateau, USA, also losing 62 ± 8 g C m⁻² yr⁻¹ (Darrouzet-Nardi et al., 2015) and finally biocrusts from the Gurbantunggut Desert, north-western China, showed a C release of 48.8 ± 5.4 to 50.9 ± 3.8 g m⁻² yr⁻¹ (Su et al., 2013). One can be fairly confident that persistent biocrusts must have a positive C balance. If they did not, they would certainly disappear from the reference habitats. However, despite the plausibility that biocrusts must have a positive net C balance, it is difficult to observe net CO₂ uptake. There are various reasons, however; two major ones are (1) positive CO₂ uptake might only occur during a small part of the year and (2) it is difficult to separate the C balance of the biocrust from C fluxes of other organisms like microbes and roots of higher vegetation or minerals like carbonate that occur below them. Nevertheless, biocrusts are only one constituent of mature soils, and it seems plausible that measurements that include soil layers other than the biocrust itself might result in CO₂ release because of a high percentage of heterotrophic organisms (Bowling et al., 2011; Darrouzet-Nardi et al., 2015; Su et al., 2013), while those that

are solely restricted to the biocrust layer might explain why they show net C uptake over the year (Brostoff et al., 2005; Wilske et al., 2008, 2009; Feng et al., 2014). We believe that seasonality (biocrust wet-up and dry-down) also plays an important role for several reasons. For example, cyanobacterial colonies exposed to wet–dry cycles apparently do not fully recover, in that quite a number of cells die during the dry period (e.g. Grilli-Caiola et al., 1993; Grilli-Caiola and Billi, 2007). Another important observation considered in the design of the present study was that the determination of CO₂ gas exchange of single species might not represent the biocrust. There is a strong influence on the outcome of the measurements when species are removed from the context of the biocrust, rather than studying the whole biocrust system (Colesie et al., 2016; Elbert et al., 2012), as this does not necessarily represent the ecological response of an intact biocrust (Weber et al., 2012).

Previously, it was observed that we could not resurrect the Australian Gulf Savannah biocrust photosynthetic activity in the middle of the dry season, even after soaking them in water for more than 24 h (Williams et al., 2014). This motivated us to perform a long-term study on an entire cyanobacteria-dominated biocrust common in northern Queensland, considered a mid-successional type in a highly seasonal environment. All these considerations led us to the following questions. (1) How do the cyanobacteria-dominated biocrusts of Boodjamulla respond to the pronounced seasonality of water availability? (2) Are these biocrusts sources or sinks for carbon at an annual timescale?

Here we focused on a common biocrust type occurring in the Gulf Plains bioregion characterized by woodlands and extensive perennial grasslands. Our investigation site was situated in Boodjamulla National Park in the Gulf Plains dry savannah region of north-eastern Australia, established in 1985, after which cattle grazing ceased. Cyanobacteria-dominated biocrusts are important drivers of ecosystem function throughout Queensland's dry savannah and especially in Boodjamulla National Park (Williams et al., 2014). There is very little rainfall during the winter dry season and the vast majority of rain commences during the summer wet season accompanied by high ambient air temperatures (> 40 °C) and high soil surface temperatures (60–74 °C). Heavy rains in the wet season often result in vast flooded plains and ephemeral wetlands (Williams et al., 2014).

2 Material and methods

2.1 Investigation site

Boodjamulla National Park (18.39° S, 138.62° E) is situated in the Gulf Savannah of north-eastern Australia covering an area of 2820 km². Mean annual rainfall is 641 mm falling mostly between December and February, although it can be highly variable with up to 1121 mm falling in the

wet years (www.bom.gov.au). Boodjamulla is mainly situated on sandstone, limestone, calcium carbonate and tufa formations sustaining *Eucalyptus* and *Melaleuca* woodlands, perennial grass floodplains, Spinifex grasslands and riparian vegetation (Fig. 1a). The biocrusts of this area are dominated by the cyanobacteria *Symplocastrum purpurascens* (Gomont ex Gomont) Anagnostidis (Fig. 2a, b, d–f), *Scytonema* sp. (Fig. 2a, b), *Symploca* sp., *Nostoc commune* Vaucher ex Bornet et Flahault and other *Nostoc* species. Other organisms occurring regularly in the Boodjamulla biocrust are the hairy liverwort *Riccia crinita* Taylor, the lichens *Peltula patellata* (Bagl.) Swinscow & Krog, *Heppia lutosa* (Ach.) Nyl., *Placidium squamulosum* (Ach.) Breuss and other small non-fertile lichen species. For a more detailed description of the locality and the biocrust see Williams and Büdel (2012) and Williams et al. (2014). We selected a site next to the Boodjamulla National Park ranger's station (Fig. 1a) with luxuriant biocrust growth (Fig. 1b), and in order to guarantee maximum control over the monitoring setup (Fig. 1c–e), kindly assisted by the park rangers. Here, the biocrust we used for the analysis was primarily formed by the two cyanobacteria *S. purpurascens* and *Scytonema* sp., with smaller amounts of other species including *Nostoc* sp., but not including bryophytes or lichens.

2.2 Sampling and sample treatment

Samples for the determination of light, temperature and water content, related to CO₂ gas exchange (environmental manipulation), and samples for the 1-year monitoring period were collected in the direct vicinity of the instrumental setup site (Fig. 1a–c). Great care was taken to ensure the homogeneity of all samples. For proper collection we removed only those top soil parts stabilized by the biocrust, which was a layer between 5 and 8 mm thick using a 8 cm wide spatula. Soil particles from underneath and not fixed to the crust by any filamentous structures were removed carefully using a soft brush and tweezers.

2.3 Environmental manipulation

For the analysis of the effect of the different environmental factors of light, temperature and water content (termed environmental manipulation throughout the text) on net photosynthesis, samples were air dried (at ~40 °C) in a 10 cm Petri dish, sealed and transported to the laboratory, where they were stored frozen (–20 °C) until measurement. This treatment has been tested in our laboratory many times with lichens of various different geographical origin, including from the tropics, and it has resulted in high survival rates (roughly 95 %) compared to storage in herbarium cabinets or boxes in the laboratory. Earlier gas exchange measurements on biocrusts, cyanobacteria, bryophytes and lichens before freezing and after thawing and re-moistening resulted in identical rates (unpub-

lished laboratory tests). Prior to measurement, samples were thawed at 23 °C for 12 h in an airtight box at low light intensities ($\ll 50 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) in order to avoid water condensation. Subsequently samples were passively rehydrated and were kept at 23 °C and natural day–night cycles ($\sim 150 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) for 2 days. Light, temperature and water content and related net photosynthesis (NP) measurements were performed using three independent replicates each. CO₂ gas exchange measurements were conducted under controlled laboratory conditions using minicuvette systems (CMS 400 and GFS 3000, Heinz Walz, Effeltrich, Germany). The response of NP and dark respiration (DR) was determined independently for light, temperature and water content (WC). Samples were weighed between measurements, and WC was calculated later in millimetres precipitation equivalent after final determination of the samples dry weight (exposed 5 days in a desiccator over silica gel at the end of the measurements). To obtain the NP response to light, fully hydrated samples ($n = 3$) were exposed to stepwise-increasing photosynthetic active radiation (PAR) from 0 to 2500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, near optimal temperature (32 °C) and ambient CO₂ concentration. The light cycle (about 30 min duration) was repeated until the samples were completely dry (after 3–4 h). Light saturation was defined as the PAR at 90 % of maximum NP. The temperature-related NP and DR were determined at increasing temperature steps, 22, 27, 32, 37, 42 and 47 °C, while light was constantly at 1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and WC was constantly at optimum ($n = 3$). The influence of WC on NP and DR was determined at constant, nearly saturating light (1500 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and at six different temperatures (22, 27, 32, 37, 42 and 47 °C), again using three replicates. Samples were completely soaked with water and exposed in the cuvette. Then, NP and DR were measured in short time intervals (roughly 10 min) until the samples were almost dry and did not show any NP nor DR reactions. After each time interval, the fresh weight of the sample was determined using a balance, and the corresponding WC to each data point was calculated using the dry weight of the sample (see above).

In all environmental manipulation, the CO₂ exchange rates of the samples were related to chlorophyll *a* content. For chlorophyll determination, the samples were ground into small pieces and then extracted twice with dimethyl sulfoxide (DMSO) at 60 °C for 90 min. The chlorophyll *a + b* content was determined and calculated according to Ronen and Galun (1984).

2.4 Field monitoring of CO₂ gas exchange

As there was only one semi-automatic cuvette system available, we could not replicate the measurements. To partly overcome this problem, we used several samples over the year. Samples were placed in a basket of thermoplastic resin with drain holes in the bottom to avoid standing water during




Figure 1. Boodjamulla National Park, measuring site. **(a)** Housing area for the park rangers and administration with location of biological soil crusts and the measuring site (white circle). **(b)** Dark patches of a biological soil crust between grass tussocks (red scale 2.5 cm). **(c)** Klapp cuvette system installed in two water-filled basins to prevent small animals from entering the device. **(d)** Measuring head of the clap cuvette system; the lid is open, exposing the wire mesh basket with the sample (IR means infrared thermocouple, L means light sensor for PAR, T means tubing for gas exchange, VP means vibration plate ensuring regular movement of the air when the cuvette is closed, H means light translucent head closing every 30 min for 3 min measurement). **(e)** Room with the data recording devices and control module.

rain events. The basket had a fixed size and all samples had exactly the same exposed surface of 16.5 cm^2 (Fig. 1d). All samples used were tested for a comparative large NP and DR rate under the given environmental conditions for two measurements (1 h) in the cuvette system and only those were used that had more-or-less identical NP and DR rates. We used 14 samples during field monitoring (Table 1) and exposed them in random mode. The “random” mode was determined by our ease of access (climatic conditions, days when authors not available) to the investigation site over the measurement period.

Water content of samples of the environmental manipulation and those of the field monitoring is always expressed as millimetre water column. As it was impossible to remove the sample after each measurement from the monitoring cuvette system to determine the fresh weight corresponding with the measured value by weighing it with a balance (measurements every half-hour; it was not possible for the authors to remain at the site for the whole period), the only method of obtaining matching values between field monitoring and controlled

experiments was to express it as rainfall in millimetres water column.

Field monitoring of the biocrust CO_2 gas exchange was recorded using a semi-automatic cuvette system (ACS) as described in detail by Lange (2002). Full technical details of the ACS (Heinz Walz, Effeltrich, Germany) are given in Lange et al. (1997a). We therefore focus on some major topics of the procedure here. The device is composed of two major parts: first the cuvette system itself, which is exposed in the natural environment of the biocrust (Fig. 1c), and secondly the controlling and data acquisition unit together with two infrared gas analysers (IRGA) for CO_2 ambient and CO_2 samples (BINOS, Rosemount, Hanau, Germany) and a pumping unit regulated by mass flow controllers (Fig. 1e). To secure data records, a data printer and a graphic plotter were added as well.

The soil crust samples were exposed on the lower part of the cuvette (Fig. 1d, arrow). When the upper lid was open (H in Fig. 1d), the sample was fully exposed to the natural environment. Measurements were taken every 30 min, during which the cuvette was closed for 3 min. We recorded the

Table 1. Samples used for monitoring (only those listed used during the active period).

1.	Sample A10	1 Jul–12 Dec 2010 (165 days)
2.	Sample C5	13–22 Dec 2010, 4–8 Jan, 12–14 Jan 2011 (18 days)
3.	Sample S1	23–29 Dec 2010 (7 days)
4.	Sample SC	30–31 Dec 2010 (2 days)
5.	Sample S4	1–3 Jan, 9–11 Jan 2011 (6 days)
6.	Sample S7	15–17 Jan 2011 (3 days)
7.	Sample 2B	18–25 Jan 2011 (8 days)
8.	Sample BS1	26 Jan–1 Feb 2011 (6 days)
9.	Sample BS2	2–13 Feb 2011 (12 days)
10.	Sample BS4	14 Feb–13 Mar 2011 (28 days)
11.	Sample BS7	14–24 Mar 2011 (11 days)
12.	Sample BS3	25 Mar–4 Apr 2011 (11 days)
13.	Sample C14	5–17 Apr 2011 (13 days)
14.	Sample C11B	18 Apr–30 Jun 2011 (73 days)

CO₂ exchange of the sample and absolute ambient CO₂ partial pressure as well as mass flow, air temperature, the sample surface temperature, air humidity and ambient photosynthetic radiation at the sample level. NP and DR were related to the area covered by the biocrust.

3 Results

The dominating cyanobacteria of the biocrust used for the long-term monitoring was the felt-like/tuft-like filamentous cyanobacterium *Symplocastrum purpurascens* forming a dark brownish stratum with erect tapering bunches of filaments (Fig. 2a–d) and the felt-like greyish *Scytonema* sp. inside and on top of the substratum (Fig. 2a–b). *S. purpurascens* is characterized by distinctly lamellate, reddish to purple-red sheath (colourless in the shade; Fig. 2e, f). After the first rain, new trichomes developed at the tips of the cyanobacterial layer.

3.1 Environmental manipulation

When exposed to stepwise-increasing PAR intensities, the biocrust did not reach full saturation of NP at optimal water content ($31.7 \pm 2.6 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at a WC of $0.7 \pm 0.1 \text{ mm}$ and 32°C ; $n = 3$) even at $2500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. At below-optimal WC (i.e. where at least 90 % of the maximum gas exchange rates are reached), a decline of NP ($21.3 \pm 5.7 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at a WC of $0.5 \pm 0.1 \text{ mm}$) was observed. This was also the case for well above optimal WC: $20.7 \pm 6.2 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at a WC of $1.0 \pm 0.1 \text{ mm}$, $7.6 \pm 3.7 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at a WC of $1.3 \pm 1.6 \text{ mm}$ and $2.3 \pm 0.2 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at a WC of $1.9 \pm 0.1 \text{ mm}$ (Fig. 3a).

Increasing air temperature from 22 to 47°C resulted in an increase of NP from 19.8 ± 1.4 to $32.4 \pm 4.5 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ ($n = 3$)

without saturation. The increase of DR was less pronounced and ranged from $-3.1 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at 22°C to $-6.3 \pm 1.4 \text{ nmol CO}_2 \text{ mg}^{-1} \text{ chlorophyll } a \text{ s}^{-1}$ at 47°C air temperature ($n = 3$; Fig. 3b).

Regarding CO₂ fixation, the optimal WC of the biocrust was $0.7 \pm 0.1 \text{ mm}$. At all temperatures the biocrust exhibited a clear optimum WC (range of 0.6 to 0.8 mm), where they reached their maximum NP. Water content below and above this optimum led to a strong decline or even a complete halt to NP (Fig. 3c). At WC of about 0.2 mm , the biocrust starts NP and DR, and with increasing WC, NP had a steep increase to maximum. A further increase of the WC created supersaturation. NP then strongly decreased to less than $1/10$ of maximal NP at optimal WC, and may have dropped down to zero (or even become negative at higher temperatures) and remained at this level (Fig. 3c).

3.2 Field monitoring of CO₂ gas exchange

Monitoring of diurnal CO₂ gas exchange of Boodjamulla biocrusts commenced on 1 July 2010 and lasted until 30 June 2011. Measurements were taken every 30 min day and night. There was no measurable gas exchange from July until the end of September 2010 and from mid-April to June 2011. With the onset of the first seasonal rains in November, the biocrust showed mainly CO₂ loss during the day, despite PAR levels of 2000 – $2500 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (Figs. 4, 5). CO₂ loss during the day was up to $1 \mu\text{mol m}^{-2} \text{ s}^{-1}$. Air temperature reached values of up to 46°C and relative air humidity increased up to 100 % during the night, dropping down to levels of 20 % during the day (Fig. S1 in the Supplement). The first positive NP was observed on 16 November (Fig. S1). From December 2010 to March 2011 rain events below 1.5 mm resulted in negative NP, whereas higher precipitation initiated positive NP of up to $8 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Figs. S2–S5). In April the rainy season ceased and small precipitation events resulted in a CO₂ loss during the day of up to $2 \mu\text{mol m}^{-2} \text{ s}^{-1}$ (Fig. S6).




Figure 2. (a) Close-up of the dry Boodjamulla biocrust at the measuring site, grey areas are dominated by the cyanobacterium *Scytonema* sp., dark-brownish areas dominated by *Symplocastrum purpurascens*. (b) Same as in (a), but after rehydration. (c) Gross fraction of the *S. purpurascens*-dominated biocrust and its stratification (EPS means extracellular polysaccharide sheath; LT-SEM). (d) In situ top view of the *S. purpurascens* biocrust. (e) Filament with red sheath from the top of the biocrust and (f) from beneath with a more-or-less colourless sheath.

The CO₂ content of the ambient air fluctuated between day and night from 370 to 470 ppm during the rainy season. Fluctuation was less pronounced in the dry season. During dark cloudy days with or without rain, fluctuation was diminished (Figs. S1–S6). In September and October 2010, the monitoring plot received the first rains. Nevertheless, although we could record metabolic activity, we were not able to calculate NP and DR due to malfunction of the ACS during these initial 2 months as well as on the following days: 1–2, 10–14, 18–20 November; 1–2, 12, 30–31 December; 22–23 March

2011. These data were excluded from further analyses (see Supplement Figs. S1–S6). An estimation on the basis of climatic data from September and October 2010 together with gas exchange data from November 2010 resulted in an estimated CO₂ loss of 88 mmol m⁻².

The sensitive interaction of the biocrust and the environmental factors can be observed in the reaction of diurnal CO₂ gas exchange over the months. For example in the night from 31 December 2010 to 1 January 2011, the biocrust was inactive but did show some DR at the end of the night and posi-




Figure 3. Response of net photosynthesis and dark respiration to water content, different PAR and temperature of the *S. purpurascens*-dominated biocrust. (a) Response of net photosynthesis to increasing PAR at different water content at 32 °C, one out of three replicates shown; (b) response of net photosynthesis and dark respiration to increasing air temperature at 1500 μmol photons m⁻² s⁻¹ and optimal WC, mean values of $n = 3$; (c) response of net photosynthesis and dark respiration of three replicates to increasing biocrust water content at 1500 μmol photons m⁻² s⁻¹ and an air temperature of 47 °C.

tive NP was measured from the morning until the afternoon (Fig. 5). As we did not record any rain, the biocrust must have been activated by dew fall or probably from some moisture in the soil as it was wet the day before. In the afternoon of 1 January, a heavy rainfall occurred, resulting in a strong water supersaturation of the biocrust. Net photosynthesis immediately became negative but recovered in the late afternoon, a pattern that could also be observed on 2 and 3 January. On 3 January PAR was so intense that the biocrust dried and metabolic activity ceased completely. The biocrust did not dry over the previous 2 days and there was DR during the whole night (Fig. 5). When comparing all positive NP values of the metabolic active period to the respective PAR and temperature values, the light saturation of the biocrust NP was reached at 2200 μmol m⁻² s⁻¹ (Fig. 6a), whereas the tem-

perature optimum was found at 37 °C (Fig. 6b). The comparison of NP with relative air humidity and PAR showed that almost all of NP was found at a relative air humidity above 42 % (Fig. 7).

While November 2010 and April 2011 had a negative CO₂ / carbon balance, December 2010 to March 2011 were positive (Table 2). Net primary productivity of the Boodjamulla biocrust was 229.1 mmol CO₂ m⁻² yr⁻¹, signifying a carbon fixation rate of 2.7 g m⁻² yr⁻¹ (Table 2; Fig. 4). Applying the September–October estimation, annual values were reduced to 144.1 mmol CO₂ m⁻² yr⁻¹, equivalent to 1.7 g C m⁻² yr⁻¹. Over the 8760 h of the 1-year measurement period, the biocrust was metabolically active for 2186 h, representing 25 % of the whole period. Of that 25 % total active period, 48.6 % was NP and 51.4 % DR (Fig. 8a). The biocrust




Figure 4. Daily CO₂ balance of the Boodjamulla biocrust. Black dots stand for dark respiration and open circles stand for gas exchange during daylight. Negative values during daylight indicate either supersaturation or water shortage.

suffered from reduced CO₂ uptake during NP periods due to water supersaturation for over 29.2 % of the photosynthetically active time (Fig. 8b).

4 Discussion

4.1 Seasonality and CO₂ balances

Apart from a clear seasonal activity pattern of the cyanobacteria-dominated biocrust from Boodjamulla National Park, Queensland, only a minority of the year was actually suitable for its growth during the 1-year CO₂ gas exchange field monitoring. An inactive winter period with no measurable CO₂ gas exchange lasted from July to mid-September 2010 and then from mid-April to the end of June 2011. Metabolic activity was found in the summer months only, starting from 23 September 2010, when the first rains commenced, continuing until 18 April 2011. Due to malfunction of the ACS, measurements from September and October and some days of November and December 2010 were not useable to calculate NP and DR. An estimation based on rainfall data from September and October, together with the reference gas exchange values from November, suggests a CO₂ loss of roughly 88 mmol m⁻². Net primary productivity was determined as 1.7 g C m⁻² yr⁻¹ (2.7 g C m⁻² yr⁻¹ without September–October correction). Our results showed that the Boodjamulla biocrust exhibited a positive net C uptake after the 1-year field monitoring period. This result is in line with the findings of several other studies but differs from all of them in the fact that our study focused on an environment with hot wet-season hydration, whereas all other studies were conducted in environments with cool season hydration. For example, a cyanobacteria-dominated biocrust in the Mojave Desert, USA, had a C gain of 11.5 g m⁻² yr⁻¹ (Brostoff et al., 2005), 6.7 times higher than the cyanobacteria-dominated Boodjamulla biocrust. Another biocrust dominated by cyanobacteria, algae, lichens and mosses from the Negev Desert, Israel, resulted in a C

gain of 0.7 to 5.1 g m⁻² yr⁻¹ (Wilske et al., 2008, 2009) and thus is pretty close to what we observed in our study, which also corresponds with the results from biocrusts composed of cyanobacteria, lichens and mosses of the Mu Us Desert in China with a C gain of 3.5 to 6.1 g m⁻² yr⁻¹ (Feng et al., 2014).

On the other hand, there are several studies that clearly demonstrate that biocrusts lose C to the atmosphere. When studying a cyanobacteria-dominated biocrust of the arid grassland in south-east Utah, USA, applying the eddy covariance method, Bowling et al. (2010) could not determine if this biocrust was a sink or a source as there were some grasses involved in the plot and hence their root respiratory CO₂ loss influenced the CO₂. When these authors applied a top soil chamber for gas exchange measurements, they found the same biocrust to be a typical C source (Bowling et al., 2011). However, this does not necessarily mean that overall they are a C source. A cyanolichen-dominated biocrust from the Gurbantunggut Desert, China, was reported as quite a large C source with a loss of -48.8 ± 5.4 to -50.9 ± 3.8 g C m⁻² yr⁻¹ (Su et al., 2012, 2013), and a very similar biocrust type of the arid grassland of the Colorado Plateau, USA, showed surprisingly similar values of -62 ± 8 g C m⁻² yr⁻¹ (Darrouzet-Nardi et al., 2015). How can this astonishing and, at first glance, contradictory fact be explained? Comparing methodology and how the measurements were taken sheds some light on this phenomenon. All investigations, including our own study that showed biocrusts having a net CO₂ uptake over the year, used gas exchange devices with a separate cuvette where the samples had to be removed from the biocrust (Brostoff et al., 2005; Feng et al., 2014) except the study of Wilske et al. (2008, 2009), in which a top soil chamber measured the biocrust in situ. All other studies with a negative C balance used top soil chambers where the biocrust was measured in situ (Bowling et al., 2011; Su et al., 2013; Darrouzet-Nardi et al., 2015). The main difference we could find in these studies was the thickness of the biocrust plus the sub-crust (soil) layer used. While the studies revealing biocrusts to be CO₂ emitters used collars penetrating 20 to 35 cm into the soil (Bowling et al., 2011; Su et al., 2013; Darrouzet-Nardi et al., 2015), the studies attributing biocrusts to be CO₂ gainers during the course of 1 year either used pieces of biocrust of 1 to 5 cm thickness (this study; Brostoff et al., 2005; Feng et al., 2014), or a collar penetrating only 5.5 cm into the soil (Wilske et al., 2008, 2009). The metabolic activity of heterotrophic organisms as well as respiration of roots from nearby plants of deeper soil levels apparently influence the CO₂ gas exchange measurements accordingly as indicated in the investigation of Bowling et al. (2011). However, soils are not a perpetual motion machine in terms of carbon balance; they can only respire as much carbon as introduced into the system. If carbon does not come from the autotrophic part of the soil system, it must be introduced from outside, either via litter transport, blown dust, animals or from run-on wa-




Figure 5. Detail of diurnal CO₂ gas exchange from January 2011, showing rain events resulting in water supersaturation of the biocrust. Blue bars indicate the approximate duration of rainfall. Green lines indicate gas exchange during daylight and black lines during the night.




Figure 6. (a) Net photosynthesis from all days related to light intensity (PAR). The biocrust shows a saturation at 2200 µmol photon m² s⁻¹ and a slight depression at 2400 µmol photons m² s⁻¹ and greater. (b) Net photosynthesis from all days related to air temperature. The optimum temperature is at 35 °C but the biocrust still performs very well at 42 °C.

ter from the surrounding environment. In a recent study using the eddy covariance method, Biederman et al. (2017) found a wide range of carbon sink/source functions. There was a mean annual net ecosystem productivity (NEP) varying from -350 to +330 g C m⁻² across sites with diverse vegetation types in the dryland ecosystems of south-western North America using evapotranspiration (ET) as a proxy for annual ecosystem water availability. Gross ecosystem productivity (GEP) and ecosystem respiration (R_{eco}) were negatively related to temperature, both interannually within and spatially across sites, and sites demonstrated a coherent response of GEP and NEP to anomalies in annual ET. Their investigation sites included one region with noteworthy biocrust cover not accompanied by dense vascular plant vegetation, the La Paz region of Baja California, with an annual C uptake (NEP) of roughly 90 g m⁻². Approximating annual C gain based on the maximal CO₂ uptake rates of four biocrust types composed of cyanobacteria, cyanolichens and chlorolichens from Baja California (Büdel et al., 2013), we approach an annual C gain for those biocrusts of 11 ± 4 g m⁻². The calculation was based on the following estimations: 90 active days per year with 34 of them having a sub-optimal CO₂ uptake rate of only 25 % of maximum due

to supersaturation. Daily rates were calculated by maximum NP for 5 h per day minus 10 h $R + DR$. This final estimation resulted in a 6.5 times higher C uptake than our pure cyanobacteria-dominated biocrust from Boodjamulla, but it is still 8 times less than found for the Baja California site in the study of Biederman et al. (2017). It could well be that later successional biocrusts with a wealth of different species groups, including bryophytes, lichens and green algae, other than cyanobacteria might reach higher productivity and thus higher annual carbon fixation rates. This should be the focus of further studies.

4.2 Carbon dioxide uptake rates and biocrust type

Maximum net CO₂ uptake rates of the Boodjamulla biocrust (8.3 µmol CO₂ m⁻² s⁻¹) clearly exceeded those of a comparable cyanobacteria-dominated biocrust from the Negev Desert, Israel, which reached maximal values of 1.1 µmol CO₂ m⁻² s⁻¹ (Lange et al., 1992), and from the Colorado Plateau, USA, with 2.0 µmol CO₂ m⁻² s⁻¹ (Darrouzet-Nardi et al., 2015). The higher NP rates of the Boodjamulla biocrusts are probably related to the felt-like structure on the soil surface, providing greater surface area for gas exchange (Fig. 2a–d), while the Negev

Table 2. Monthly net primary productivity of the Boodjamulla biological soil crust (values in brackets are an estimation only, not based on measurements; see text for explanation).

Month	NPP (NP-DR)	
	(mmol CO ₂ m ⁻² month ⁻¹)	(g C m ⁻² month ⁻¹)
July	0	0
August	0	0
September 2010	0 (−2.0)	0 (−0.02)
October 2010	0 (−83.0)	0 (−1.01)
November 2010	−210.26	−2.53
December 2010	110.20	1.32
January 2011	99.59	1.20
February 2011	80.99	0.97
March 2011	174.11	2.09
April 2011	−25.54	−0.31
May 2011	0	0
June 2011	0	0
Annual	229.09 mmol CO ₂ m ⁻² yr ⁻¹ (144.08 mmol CO ₂ m ⁻² yr ⁻¹)	2.74 g C m ⁻² yr ⁻¹ (1.71 g C m ⁻² yr ⁻¹)




Figure 7. Contour plot of net photosynthesis of the Boodjamulla biocrust based on linear interpolation between measured values. Shown is the active period from November 2010 to April 2011. Net photosynthesis is related to relative air humidity and photosynthetic active radiation (PAR). No dark respiration values shown. Colour key: yellow means no activity, orange to red means CO₂ loss during the day (supersaturation) and light green to violet means CO₂ uptake.

Desert biocrust was a thin layer of cyanobacterial filaments slightly beneath the surface. Annual carbon fixation rates of cyanobacteria-dominated arid region biocrusts are gen-

erally lower (this study; Brostoff et al., 2005; Feng et al., 2014; Wilske et al., 2009) compared with biocrusts including lichens and bryophytes (see summarizing Table 15.2 in Sancho et al., 2016; Elbert et al., 2012; Porada et al., 2013) or the carbon gain of isolated biocrust organisms, for example the green algal lichen *Lecanora muralis* (Schreber) Rabenh. from a rock crust with 21.5 g C m⁻² yr⁻¹ (Lange, 2002, 2003a, b). It follows that the developmental stage of a biocrust might be an important factor for photosynthetic performance, too. This view is supported by the results of a study (Housman et al., 2006) determining NP rates depending on successional stages rather than developmental stages. In this study the authors classified biocrusts as early successional (*Microcoleus*) or later successional stages (*Nostoc/Scytonema* or *Placidium/Collema*), and differences in NP, which was on average 1.2–2.8-fold higher in later successional crusts compared to the early successional stages (Housman et al., 2006). When considering the Boodjamulla biocrust as a mid-successional type, an increase in carbon gain might be expected in the future when lichens and bryophytes establish to form a later successional soil crust. Dealing here with a cyanobacteria-dominated biocrust of a mid-successional type might explain the low C uptake rates and also the apparent discrepancy in the values calculated for the global net primary productivity (NPP) by cryptogamic covers of Elbert et al. (2012).

4.3 Active times and water relation

The Boodjamulla biocrust had metabolic activity for only 25 % of the year, 12.3 % NP and 12.8 % DR (Fig. 8a). In 29.2 % of the photosynthetic active time, CO₂ fixation was lowered considerably by water supersaturation. For compar-




Figure 8. (a) Mean diel activity of the Boodjamulla biocrust: black means inactive; light grey means photosynthetically active; dark grey means dark respiration and hatched means metabolic activity, but due to technical failure of instrumentation, unclear if NP or DR. (b) Monthly extent of water supersaturated periods during the photosynthetic (NP) active time of the Boodjamulla biocrust. Black means periods of supersaturation and light grey means periods of conducive water supply.

ison, the lichen *L. muralis* from a temperate climate was active for 35.5 % of the year, 16.7 % NP and 18.9 % DR. During periods of photosynthesis, the lichen was heavily suppressed by water supersaturation at 38.5 % (Lange, 2003a, b). It is obvious that the strict seasonal rainfall pattern is a major contributing reason for the considerably lower metabolic activity of the savannah-type biocrust from Boodjamulla compared to the rock crust lichen *L. muralis* in a temperate climate with rainfall expanding over the whole year. As characteristic for poikilohydric organisms, both the Boodjamulla biocrust and the rock crust lichen suffer considerably from water supersaturation causing waterlogged gas diffusion channels and

thus drastically limiting CO₂ gas exchange (see Green et al., 2011, and references therein).

As metabolic activity is strictly bound to the presence of water it is important to know the role of water content on photosynthetic performance and respiratory CO₂ exchange. The Boodjamulla biocrust achieved maximum NP values at 0.5–0.8 mm WC and had a lower compensation point for NP at 0.1 mm WC. Comparable values were found for the Negev biocrust studied by Lange et al. (1992) and the rock crust lichen *L. muralis* (Lange, 2002). In the chlorolichens of a biocrust from Utah, comprised of *Diploschistes diacapsis* (Ach.) Lumbsch, *Psora cerebriformis* W. Weber and *Squamarina lentigera* (Weber) Poelt, photosynthetic metabolism was activated by extremely small amounts of moisture. The lower compensation point for NP is between 0.05 and 0.27 mm WC. Maximal NP occurred between 0.4 and 1.0 mm WC (Lange et al., 1997b). The values for the cyanobacterial soil crust lichen *Collema tenax* (Swartz) Ach., however, were considerably higher with the lower NP compensation point at 0.2 mm WC and maximal NP between 0.8 to 1.2 mm WC, but it performed NP under much higher temperatures than the above-mentioned green algal lichens (Lange et al., 1998).

Almost all gas exchange activity of the Boodjamulla biocrusts occurred at air relative humidity above 42 % (Fig. 7). This, however, must be taken with care as it does not mean that the biocrust is always active at this value and above. Like all cyanobacteria investigated so far, the cyanobacteria of the Boodjamulla biocrusts are also not activated by air humidity alone (e.g. Lange et al., 1992, 1993, 1994; own unpublished results). The value of 42 % relative humidity is merely a good indicator for the right combination of WC (rainfall dependent), temperature and light. A comparable observation has been made by Raggio et al. (2017), who found air relative humidity ($\ll 50\%$) and air temperature as the best predictors of metabolic activity duration for four different biocrust types across western Europe. In a number of cases we found activation of the Boodjamulla biocrust without any measurable precipitation (Fig. 5, Supplement Fig. S3, 4–5 January). This is likely explained by dew formation, a non-rainfall water source found playing an important role in biocrusts (Lange et al., 1994; Ouyang and Hu, 2017) and also observed at Boodjamulla during the wet season. There are a number of studies that found dew formation was important in biocrust systems, for example the study of Jacobs et al. (2000), where in a desert environment in Israel daily amounts of dew ranged between 0.1 and 0.3 mm night⁻¹. Dew formation determined for an inland dune biocrust community in Germany ranged from 0.04 and 0.18 kg m² over 2 days (Fischer et al., 2012). Even fog was identified as a major source of non-rainfall water driving biocrust productivity in the Atacama Desert of Chile, where approximately 8 to 24 % of the fog water flux was available to the biocrusts at the soil surface (Lehnert et al., 2018).

4.4 Reestablishment and resurrection after the dry season

What are the reasons for negative C balances of the biocrusts during the first active months after start of the rainy season? We suggest that in contrast to eukaryotic poikilohydric photoautotrophs such as liverworts, mosses or lichens, which resuscitate all thallus compartments after hydration, prokaryotic cyanobacteria show considerable dieback rates during longer dry periods or drought events (see Williams and Eldridge, 2011; Williams and Büdel, 2012). For example in the terrestrial, colony-forming unicellular genus *Chroococcidiopsis* the number of viable cells decreased with age of the colony and the length of exposure to drought (Grilli-Caiola et al., 1993; Grilli-Caiola and Billi, 2007). Desiccation-tolerant *Chroococcidiopsis* cells must either protect their components from desiccation-induced damage or repair it after rehydration. It was found that desiccation survivors limit genome fragmentation, preserve intact plasma membranes and have spatially reduced reactive oxygen species accumulation and dehydrogenase activity whereas damaged cells do not (Billi, 2009). In the abundant biocrust cyanobacterium *Microcoleus vaginatus* Gomont ex Gomont, immediate but transient induction of DNA repair and regulatory genes signalled the hydration event, and recovery of photosynthesis occurred within 1 h accompanied by up-regulation of anabolic pathways (Rajeev et al., 2013). In general, during the desiccated period homoiochlorophyllous cyanobacteria (maintaining their chlorophyll during desiccation) still suffer from photoinhibition induced by the typical high light intensities of their habitat. Nevertheless, resurrection of photosynthesis after desiccation occurs within hours or days, depending on the degree of damage (Lüttge, 2011), while reestablishment takes days or weeks, largely depending on the availability of water, and it also needs a positive C input. Our field monitoring uncovered numerous events of supersaturation during daylight combined with low NP rates after the onset of the active season. Supersaturation events later in the season are easily compensated for by high NP rates (Fig. 4). We interpret the early C loss phase after the drought, at least partly as a reestablishment period of the biocrusts structure, enabling the biocrust diminishing supersaturation events by means of erect cyanobacterial filament bundles, standing out of a covering water film and thus probably improving CO₂ gas diffusion (Fig. 2c, d).

4.5 Influence of temperature and global warming

In the environmental manipulation setup the Boodjamulla biocrusts did respond to air higher air temperatures (20–47 °C) with a continuous increase of both NP and DR, with NP increasing at slightly higher rates than DR (Fig. 3b). Even at 47 °C, CO₂ uptake did not show any reduction, nor did DR show a considerably stronger CO₂ release, and NP still exceeded DR by a factor of 5. However, in our field measure-

ments we did not observe biocrust activity above air temperatures of 43 °C (Fig. 6b); at this temperatures the biocrust was dry and inactive. During field monitoring the optimal temperature for positive NP was around 35 °C (Fig. 6b). Applying an experimental air temperature increase of 2–3 °C, Maestre et al. (2013) observed a drastic reduction in a lichen-dominated biocrust cover of ca. 44 % over 4 years in a dry-land ecosystem in Spain. Soil CO₂ efflux was increased and soil net CO₂ uptake was reduced with the additional warming. According to the field monitoring gas exchange rates of the Boodjamulla biocrusts, we would expect even shorter activity periods under the scenario of global warming and, relatedly, probably lower C uptake or even C loss resulting in a pronounced reduction of biocrust coverage. Other indirect effects of warming should be expected when it influences rainfall amount and regime. It can be speculated that fewer, but heavier, rain events would certainly effect the Boodjamulla biocrust by increasing supersaturation periods resulting in lower or even no carbon gain, also probably causing a pronounced reduction in coverage.

5 Conclusions

The Boodjamulla biocrust showed highly seasonal photosynthesis-related metabolic activity divided into four major periods: (1) a metabolically inactive winter time; (2) onset of the photosynthetic active period, starting with roughly 3 months of reestablishment, limited CO₂ uptake due to supersaturation and a hypothesized increased activity of heterotrophic organisms decomposing organic matter from old biocrusts; (3) a 4-month period of net C uptake; and (4) about 1 month with C loss until a complete cease of activity. During the four periods, NP and NPP rates vary strongly and thus seasonality plays an important role. For eco-physiological experiments, the period of the year biocrust material is sampled is absolutely crucial. The cyanobacteria-dominated Boodjamulla biocrust turned out to be a small but consistent sink of carbon as it grew, and it also potentially contributes to the soil organic matter. From the magnitude of values it is clear that the observed C fluxes are not at all close to what a plant community can do. Methodological approaches analysing biocrust carbon cycling critically must consider that including or excluding sub-biocrust partitions might influence the consideration of biocrust as either as a sink or a source. There is an urgent need for more long-term measurements on different biocrust types and developmental stages in all climatic regions of the world.

Data availability. Currently, data can only be accessed in the form of Excel sheets via the corresponding author.

The Supplement related to this article is available online at <https://doi.org/10.5194/bg-15-491-2018-supplement>.

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue “Biological soil crusts and their role in biogeochemical processes and cycling”. It is a result of the BIOCRUST3 conference, Moab, USA, 26 to 30 September 2016.

Acknowledgements. We acknowledge the Waayni people, the original inhabitants of the Boodjamulla area, and thank the staff of the Boodjamulla National Park and Adels Grove. Special thanks to Tres McKenzie for on-site assistance. We also thank AgForce Qld and Century Mine for their financial and kind support. Burkhard Büdel gratefully acknowledges financial support by the German Research Foundation (DFG).

Edited by: Emilio Rodriguez-Caballero

Reviewed by: Matthew Bowker and two anonymous referees

References

- Belnap, J.: Surface disturbances: their role in accelerating desertification, *Environ. Monit. Assess.*, 37, 39–57, 1995.
- Belnap, J., Weber, B., and Büdel, B.: Biological soil crusts as an organizing principle in drylands, in: *Biological soil crusts: An organizing principle in drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., *Ecol. Stud.*, 226, 3–13, 2016.
- Beraldi-Campesi, H. and Retallack, G. J.: Terrestrial ecosystems in the Precambrian, in: *Biological soil crusts: An organizing principle in drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., *Ecol. Stud.*, 226, 37–54, 2016.
- Biederman, J. A., Scott, R. L., Bell, T. W., Bowling, D. R., Dore, S., Garatuza-Payan, J., Kolb, T. E., Krishan, P., Krofcheck, D. J., Litvak, M. E., Maurer, G. E., Meyers, T. P., Oechel, W. C., Papuga, S. A., Ponce-Campos G. E., Rodriguez, J. C., Smith, W. K., Vargas, R., Watts, C. J., Yopez, E. A., and Goulden, M. L.: CO₂ exchange and evapotranspiration across dryland ecosystems of southwestern North America, *Glob. Change Biol.*, 23, 4204–4221, 2017.
- Billi, D.: Subcellular integrities in *Chroococcidiopsis* sp. CCMEE 029 survivors after prolonged desiccation revealed by molecular probes and genome stability assays, *Extremophiles*, 13, 49–57, 2009.
- Bowling, D. R., Bethers-Marchetti, S., Lunch, C. K., Grote, E. E., and Belnap, J.: Carbon, water, and energy fluxes in a semiarid cold desert grassland during and following multiyear drought, *J. Geophys. Res.*, 115, G04026, <https://doi.org/10.1029/2010JG001322>, 2010.
- Bowling, D. R., Grote, E. E., and Belnap, J.: Rain pulse response of soil CO₂ exchange by biological soil crusts and grasslands of the semiarid Colorado Plateau, United States, *J. Geophys. Res.*, 116, G03028, <https://doi.org/10.1029/2011JG001643> 2011.
- Brostoff, W. N., Sharifi, M. R., and Rundel, P. W.: Photosynthesis of cryptobiotic soil crusts in a seasonally inundated system of pans and dunes in the western Mojave Desert, CA, *Field studies, Flora*, 200, 592–600, 2005.
- Büdel, B.: Synopsis: Comparative Biogeography and Ecology of Soil-Crust Biota, in: *Biological soil crusts: structure, function, and management*, edited by: Belnap, J. and Lange O. L., *Ecol. Stud.*, 150, 141–154, 2001.
- Büdel, B., Vivas, M., and Lange, O. L.: Lichen species dominance and the resulting photosynthetic behavior of Sonoran Desert soil crust types (Baja California, Mexico), *Ecol. Processes*, 2, 6, <https://doi.org/10.1186/2192-1709-2-6>, 2013.
- Büdel, B., Colesie, C., Green, T. A., Grube, M., Suau, R. L., Loewen-Schneider, K., Maier, S., Peer, T., Pintado, A., Raggio, J., Ruprecht, U., Sancho, L. G., Schroeter, B., Türk, R., Weber, B., Wedin, M., Westberg, M., Williams, L., and Zheng, L.: Improved appreciation of the functioning and importance of biological soil crusts in Europe: the Soil Crust International Project (SCIN), *Biodivers. Conserv.*, 23, 1639–1658, 2014.
- Castillo-Monroy, A. P., Maestre, F. T., Rey, A., Soliveres, S., and García-Palacios, P.: Biological soil crust microsites are the main contributor to soil respiration in semiarid ecosystem, *Ecosystems*, 18, 835–847, 2011.
- Colesie, C., Felde, V. J. M. N. L., and Büdel, B.: Composition and Macrostructure of Biological Soil Crusts, in: *Biological soil crusts: An organizing principle in drylands*, *Ecological Studies* 226, edited by: Weber, B., Büdel, B., and Belnap, J., Springer International Publishing Switzerland, 159–172, 2016.
- Darrouzet-Nardi, A., Reed, S. C., Grote, E. E., and Belnap, J.: Observations of net soil exchange of CO₂ in a dryland show experimental warming increases carbon losses in biocrust soils, *Biogeochemistry*, 126, 363–378, 2015.
- Elbert, W., Weber, B., Burrows, S., Steinkamp, J., Büdel, B., Andreae, M. O., and Pöschl, U.: Contribution of cryptogamic covers to the global cycles of carbon and nitrogen, *Nat. Geosci.*, 5, 459–462, 2012.
- Feng, W., Zhang, Y., Wu, B., Qin, S., and Lai, Z.: Influence of environmental factors on carbon dioxide exchange in biological soil crusts in desert areas, *Arid Land Res. Manag.*, 28, 186–196, 2014.
- Fischer, T., Veste, M., Bens, O., and Hüttl, R. F.: Dew formation on the surface of biological soil crusts in central European sand ecosystems, *Biogeosciences*, 9, 4621–4628, <https://doi.org/10.5194/bg-9-4621-2012>, 2012.
- Green, T. G. A., Sancho, L. G., and Pintado, A.: Ecophysiology of desiccation/rehydration cycles in mosses and lichens, in: *Plant desiccation tolerance*, edited by: Lüttge, U., Beck, E., and Bartels, D., *Ecol. Stud.*, 215, 89–120, 2011.
- Grilli-Caiola, M. and Billi, D.: *Chroococcidiopsis* from desert to Mars, in: *Algae and cyanobacteria in extreme environments*, edited by: Seckbach, J., Cellular origin, life in extreme habitats and Astrobiology, 11. Springer, Dordrecht, 553–568, 2007.
- Grilli-Caiola, M., Ocampo-Friedmann, R., and Friedmann, E. I.: Cytology of long-term desiccation in the desert cyanobacterium *Chroococcidiopsis* (Chroococcales), *Phycologia*, 32, 315–322, 1993.

- Housman, D. C., Powers, H. H., Collins, A. D., and Belnap, J.: Carbon and nitrogen fixation differ between successional stages of biological soil crusts in the Colorado Plateau and Chihuahuan Desert, *J. Arid Environ.*, 66, 620–634, 2006.
- Jacobs, A. F. G., Heusinkveld, B. G., and Berkowicz, S. M.: Dew measurements along a longitudinal sand dune transect, Negev Desert, Israel, *Int. J. Biometeorol.*, 43, 184–190, 2000.
- Karsten, U., Herburger, K., and Holzinger, A.: Living in biological soil crust communities of African deserts – physiological traits of green algal *Klebsormidium* species (Streptophyta) to cope with desiccation, light and temperature gradients, *J. Plant Phys.*, 194, 2–12, 2016.
- Lalonde, S. V. and Konhauser, K. O.: Benthic perspective on Earth's oldest evidence for oxygenic photosynthesis, *P. Natl. Acad. Sci. USA*, 112, 995–1000, 2015.
- Lange, O. L.: Photosynthetic productivity of the epilithic lichen *Lecanora muralis*: Long term field monitoring of CO₂ exchange and its physiological interpretation. I. Dependence of photosynthesis on water content, light, temperature, and CO₂ concentration from laboratory measurements, *Flora*, 197, 233–249, 2002.
- Lange, O. L.: Photosynthetic productivity of the epilithic lichen *Lecanora muralis*: Long term field monitoring of CO₂ exchange and its physiological interpretation. II. Diel and seasonal patterns of net photosynthesis and respiration, *Flora*, 198, 55–70, 2003a.
- Lange, O. L.: Photosynthetic productivity of the epilithic lichen *Lecanora muralis*: Long term field monitoring of CO₂ exchange and its physiological interpretation. III. Diel, seasonal, and annual carbon budgets, *Flora*, 198, 277–292, 2003b.
- Lange, O. L. and Belnap, J.: How biological soil crusts became recognized as a functional unit: A selective history, in: *Biological soil crusts: An organizing principle in drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., *Ecol. Stud.*, Springer International Publishing Switzerland, 226, 15–33, 2016.
- Lange, O. L., Kidron, G. J., Büdel, B., Meyer, A., Kilian, E., and Abeliovich, A.: Taxonomic composition and photosynthetic characteristics of the biological soil crusts' covering sand dunes in the western Negev Desert, *Funct. Ecol.*, 6, 519–527, 1992.
- Lange, O. L., Büdel, B., Meyer, A., and Kilian, E.: Further evidence that activation of net photosynthesis by dry cyanobacterial lichens requires liquid water, *Lichenologist*, 25, 175–189, 1993.
- Lange, O. L., Meyer, A., Zellner, H., and Heber, U.: Photosynthesis and water relations of lichen soil crusts: field measurements in the coastal fog zone of the Namib Desert, *Funct. Ecol.*, 8, 253–264, 1994.
- Lange, O. L., Meyer, A., and Büdel, B.: Net photosynthesis activation of a desiccated cyanobacterium without liquid water in high air humidity alone. Experiments with *Microcoleus sociatus* isolated from a desert soil crust, *Funct. Ecol.*, 8, 52–57, 1994.
- Lange, O. L., Reichenberger, H., and Walz, H.: Continuous monitoring of CO₂ exchange of lichens in the field: short term enclosure with an automatically operating cuvette, *Lichenologist*, 29, 259–274, 1997a.
- Lange, O. L., Belnap, J., Reichenberger, H., and Meyer, A.: Photosynthesis of green algal soil crust lichens from arid lands in southern Utah, USA: role of water content on light and temperature responses of CO₂ exchange, *Flora*, 192, 1–15, 1997b.
- Lange, O. L., Belnap, J., and Reichenberger, H.: Photosynthesis of the cyanobacterial soil-crust lichen *Collema tenax* from arid lands in southern Utah, USA: role of water content on light and temperature responses of CO₂ exchange, *Funct. Ecol.*, 12, 195–202, 1998.
- Lehnert, L. W., Thies, B., Trachte, K., Achilles, S., Osses, P., Baumann, K., Schmidt, J., Samolov, E., Jung, P., Leinweber, P., Karsten, U., Büdel, B., and Bendix, J.: A case study on fog/low stratus occurrence at Las Lomitas, Atacama Desert (Chile) as a water source for biological soil crusts, *Aerosol Air Qual. Res.*, 18, 254–269, 2018.
- Lüttge, U.: Cyanobacteria: multiple stresses, desiccation-tolerant photosynthesis and di-nitrogen fixation, in: *Plant desiccation tolerance*, edited by: Lüttge, U., Beck, E., and Bartels, D., *Ecol. Stud.*, 215, 23–43, 2011.
- Maestre, F. T., Escolar, C., De Guevara, M. L., Quero, J. L., Lazaro, R., Delgado-Baquerizo, M., Ochoa, V., Berdugo, M., Gozalo, B., and Gallardo, A.: Changes in biocrust cover drive carbon cycle responses to climate change in drylands, *Glob. Change Biol.*, 19, 3835–3847, 2013.
- Ouyang, H. and Hu, C.: Insight into climate change from the carbon exchange of biocrusts utilizing non-rainfall water, *Sci. Rep.*, 7, 2573, <https://doi.org/10.1038/s41598-017-02812-y>, 2017.
- Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating global carbon uptake by lichens and bryophytes with a process-based model, *Biogeosciences*, 10, 6989–7033, <https://doi.org/10.5194/bg-10-6989-2013>, 2013.
- Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating impacts of lichens and bryophytes on global biogeochemical cycles, *Global Biogeochem. Cy.*, 28, 71–85, 2014.
- Raggio, J., Green, T. G. A., Sancho, L. G., Pintado, A., Colesie, C., Weber, B., and Büdel, B.: Biological Soil Crusts across Europe behave as a single functional type regardless of composition or location, *Geoderma*, 306, 10–17, 2017.
- Rajeev, L., Nunes da Rocha, U., Klitgord, N., Luning, E. G., Fortney, J., Axen, S. D., Shih, P. M., Bouskill, N. J., Bowen, B. P., Kerfeld, C. A., Garcia-Pichel, F., Brodie, E. L., Northen, T. R., and Mukhopadhyay, A.: Dynamic cyanobacterial response to hydration and dehydration in a desert biological soil crust, *ISME J.*, 7, 2178–2191, 2013.
- Ronen, R. and Galun, M.: Pigment extraction from lichens with dimethylsulfoxide (DMSO) and estimation of chlorophyll degradation, *Environ. Exp. Bot.*, 24, 239–245, 1984.
- Sancho, L. G., Belnap, J., Colesie, C., Raggio, J., and Weber, B.: Carbon budgets of biological soil crusts at micro-, meso-, and global scales, in: *Biological soil crusts: an organizing principle in drylands*, edited by: Weber, B., Büdel, B., and Belnap, J., *Ecol. Stud.*, 226, 287–304, 2016.
- Su, Y. G., Wu, L., and Zhang, Y. M.: Characteristics of carbon flux in two biologically crusted soils in the Gurbantunggut Desert, Northwestern China, *Catena*, 96, 41–48, 2012.
- Su, Y. G., Wu, L., Zhou, Z. B., Liu, Y. B., and Zhang, Y. M.: Carbon flux in deserts depends on soil cover type: A case study in the Gurbantunggut desert, North China, *Soil Biol. Biochem.*, 58, 332–340, 2013.
- Watanabe, Y., Martini, J. E. J., and Ohmoto, H.: Geochemical evidence for terrestrial ecosystems 2.6 billion years ago, *Nature*, 408, 574–578, 2000.
- Weber, B., Graf, T., and Bass, M.: Ecophysiological analysis of moss-dominated biological soil crusts and their separate components from the Succulent Karoo, South Africa, *Planta*, 236, 129–139, 2012.

- Williams, W. J. and Büdel, B.: Species diversity, biomass and long-term patterns of biological soil crusts with special focus on Cyanobacteria of the *Acacia aneura* Mulga Lands of Queensland, Australia, *Algol. Stud.*, 140, 23–50, 2012.
- Williams, W. J. and Eldridge, D. J.: Deposition of sand over a cyanobacterial soil crust increases nitrogen bioavailability in a semi-arid woodland, *Appl. Soil. Ecol.*, 49, 26–31, 2011.
- Williams, W. J., Büdel, B., Reichenberger, H., and Rose, N.: Cyanobacteria in the Australian northern savannah detect the difference between intermittent dry season and wet season rain, *Biodivers. Conserv.*, 23, 1827–1844, 2014.
- Wilske, B., Burgheimer, J., Karnieli, A., Zaady, E., Andreae, M. O., Yakir, D., and Kesselmeier, J.: The CO₂ exchange of biological soil crusts in a semiarid grass-shrubland at the northern transition zone of the Negev desert, Israel, *Biogeosciences*, 5, 1411–1423, <https://doi.org/10.5194/bg-5-1411-2008>, 2008.
- Wilske, B., Burgheimer, J., Maseyk, K., Karnieli, A., Zaady, E., Andreae, M. O., Yakir, D., and Kesselmeier, J.: Modeling the variability in annual carbon fluxes related to biological soil crusts in a Mediterranean shrubland, *Biogeosciences Discuss.*, <https://doi.org/10.5194/bgd-6-7295-2009>, 2009.