

# Two equivalent norms for vector-valued holomorphic functions

by Peter VIETEN, Kaiserslautern

## Abstract

The following two norms for holomorphic functions  $F$ , defined on the right complex half-plane  $\{z \in \mathbf{C} : \Re(z) > 0\}$  with values in a Banach space  $X$ , are equivalent:

$$\begin{aligned}\|F\|_{H_p(\mathbf{C}_+)} &= \sup_{a>0} \left( \int_{-\infty}^{\infty} \|F(a+ib)\|^p db \right)^{1/p}, \quad \text{and} \\ \|F\|_{H_p(\Sigma_{\pi/2})} &= \sup_{|\theta|<\pi/2} \left( \int_0^{\infty} \|F(re^{i\theta})\|^p dr \right)^{1/p}.\end{aligned}$$

As a consequence, we derive a description of boundary values of sectorial holomorphic functions, and a theorem of Paley-Wiener type for sectorial holomorphic functions.

## 1 Two equivalent norms for vector-valued holomorphic functions

In this note we study holomorphic functions  $F$  defined on the open right complex half-plane  $\mathbf{C}_+ = \{z \in \mathbf{C} : \Re(z) > 0\}$  with values in a complex Banach space  $X$ . To such functions we assign the following extended real numbers:

$$\begin{aligned}\|F\|_{H_p(\mathbf{C}_+)} &= \sup_{a>0} \left( \int_{-\infty}^{\infty} \|F(a+ib)\|^p db \right)^{1/p}, \quad \text{and} \\ \|F\|_{H_p(\Sigma_{\pi/2})} &= \sup_{|\theta|<\pi/2} \left( \int_0^{\infty} \|F(re^{i\theta})\|^p dr \right)^{1/p}.\end{aligned}$$

Let  $H_p(\mathbf{C}_+, X)$  and  $H_p(\Sigma_{\pi/2}, X)$  be the respective normed spaces of those functions  $F$  for which  $\|F\|_{H_p(\mathbf{C}_+)}$  and  $\|F\|_{H_p(\Sigma_{\pi/2})}$  are finite.

Držbašjan and Martirosjan [1] proved that, for  $p = 2$ ,  $H_p(\mathbf{C}_+, \mathbf{C})$  is isomorphic to  $H_p(\Sigma_{\pi/2}, \mathbf{C})$ , and Sedleckii [10] showed that this result holds for all  $0 < p < \infty$ . Luxemburg [6] gave a new prove of Sedleckii's result. Our main Theorem 1 states that, for all  $1 \leq p < \infty$  and every Banach space  $X$ , the norms  $\|\cdot\|_{H_p(\mathbf{C}_+)}$  and  $\|\cdot\|_{H_p(\Sigma_{\pi/2})}$  are equivalent, and the spaces  $H_p(\mathbf{C}_+, X)$  and  $H_p(\Sigma_{\pi/2}, X)$  are isomorphic. As a consequence, we derive in the second section a description of boundary values of sectorial holomorphic functions, and we give in the third section a theorem of Paley-Wiener type for sectorial holomorphic functions.

Throughout this note we use the following notations. The letter  $X$  stands for a complex Banach space, and we write  $X^*$  for its dual. We denote by  $L_p(\mathbf{R}, X)$  the usual Bochner  $L_p$ -space, and  $L_p(\mathbf{R}, \mathbf{C})$  is abbreviated by  $L_p(\mathbf{R})$ . For  $z = a + ib \in \mathbf{C}_+$  the Poisson kernel  $P_z$  is given by

$$P_z(t) = \frac{1}{\pi} \frac{a}{(b-t)^2 + a^2}, \quad -\infty < t < \infty.$$

We say that  $F : \mathbf{C}_+ \rightarrow X$  is the Poisson integral of some function  $f \in L_p(\mathbf{R}, X)$  if

$$F(z) = \int_{-\infty}^{\infty} P_z(t) f(t) dt, \quad z \in \mathbf{C}_+.$$

**THEOREM 1** *Let  $1 \leq p \leq \infty$  and let  $F : \mathbf{C}_+ \rightarrow X$  be holomorphic. Then  $F \in H_p(\mathbf{C}_+, X)$  if and only if  $F \in H_p(\Sigma_{\pi/2}, X)$ . Moreover,*

$$2^{-1} \|F\|_{H_p(\mathbf{C}_+)} \leq \|F\|_{H_p(\Sigma_{\pi/2})} \leq 2^{-1/p} \|F\|_{H_p(\mathbf{C}_+)}.$$

*Proof.* There is nothing to prove if  $p = \infty$ . Let  $1 \leq p < \infty$ , take  $F \in H_p(\mathbf{C}_+, X)$ , and assume that  $F$  is the Poisson integral of some function  $f \in L_p(\mathbf{R}, X)$ . Let  $-\pi/2 < \theta < \pi/2$ ,  $r > 0$ , and take  $x^* \in X^*$  with  $\|x^*\| \leq 1$ . From the proof of Theorem (3.3) in [6] it follows that

$$|x^* F(re^{i\theta})|^{p/2} \leq \frac{\sin(\theta)}{\pi} \int_{-\infty}^{\infty} P(t, \theta) |x^* f(rt)|^{p/2} dt,$$

where  $P(t, \theta) = (1 + t^2 - 2t \cos(\theta))^{-1}$ . Hence

$$\|F(re^{i\theta})\|^{p/2} \leq \frac{\sin(\theta)}{\pi} \int_{-\infty}^{\infty} P(t, \theta) \|f(rt)\|^{p/2} dt.$$

We conclude from this estimate (see the proof of Theorem (3.3) in [6])

$$\int_0^\infty \|F(re^{i\theta})\|^p dr \leq \left( \frac{\sin(\theta)}{\pi} \int_{-\infty}^\infty \frac{1}{\sqrt{|t|}} P(t, \theta) dt \right)^2 \|f\|_{L_p}^p \leq 2 \|F\|_{H_p(\mathbf{C}_+)}^p.$$

If  $F \in H_p(\mathbf{C}_+, X)$  is not the Poisson integral of a function in  $L_p(\mathbf{R}, X)$  then we proceed as follows. For  $a > 0$  and  $z \in \mathbf{C}_+$  put  $F_a(z) = F(a+z)$ . Then  $F_a$  belongs to  $H_p(\mathbf{C}_+, X)$  with  $\|F_a\|_{H_p(\mathbf{C}_+)} \leq \|F\|_{H_p(\mathbf{C}_+)}$ , and  $F_a$  is the Poisson integral of  $f_a \in L_p(\mathbf{R}, X)$ , where  $f_a(t) = F(a+it)$ . This follows from the corresponding scalar-valued result (see e.g. the proof of Theorem I.3.5. in [2]) by applying linear functionals. Now, the first part of the proof yields for all  $-\pi/2 < \theta < \pi/2$

$$\int_0^\infty \|F_a(re^{i\theta})\|^p dr \leq 2 \|F_a\|_{H_p(\mathbf{C}_+)}^p \leq 2 \|F\|_{H_p(\mathbf{C}_+)}^p.$$

Hence it follows for all  $0 < A < B < \infty$

$$\int_A^B \|F(re^{i\theta})\|^p dr = \lim_{a \rightarrow 0^+} \int_A^B \|F_a(re^{i\theta})\|^p dr \leq 2 \|F\|_{H_p(\mathbf{C}_+)}^p.$$

Now, letting  $A \rightarrow 0$  and  $B \rightarrow \infty$  yields the second inequality.

Conversely, let  $F \in H_p(\Sigma_{\pi/2}, X)$ . For  $\pi/2 < \alpha < \pi$  and  $z \in \mathbf{C}_+$  put

$$F_\alpha(z) = \left( \frac{\pi}{2\alpha} \right)^{1/p} z^{\frac{1}{p}(\frac{\pi}{2\alpha}-1)} F(z^{\frac{\pi}{2\alpha}}).$$

Then  $F_\alpha \in H_p(\Sigma_{\pi/2}, X)$  and

$$\|F_\alpha\|_{H_p(\Sigma_{\pi/2})} \leq \|F\|_{H_p(\Sigma_{\pi/2})}.$$

On the other hand, we know from [6] that  $x^* \circ F_\alpha$  belongs to  $H_p(\mathbf{C}_+)$  for all  $x^* \in X^*$ . Hence  $x^* \circ F_\alpha$  is the Poisson integral of its boundary function  $x^* \circ f_\alpha \in L_p(\mathbf{R})$ , where  $f_\alpha(t) = F_\alpha(it)$  for  $t \neq 0$ . Since  $f_\alpha$  is  $p$ -integrable we conclude

$$F_\alpha(z) = \int_{-\infty}^\infty P_z(t) f_\alpha(t) dt.$$

Put  $g_\alpha(t) = \|f_\alpha(t)\|$ . Then

$$\|g_\alpha\|_{L_p} = \|f_\alpha\|_{L_p} \leq 2 \|F_\alpha\|_{H_p(\Sigma_{\pi/2})} \leq 2 \|F\|_{H_p(\Sigma_{\pi/2})}.$$

We now distinguish the cases  $p > 1$  and  $p = 1$ .

If  $p > 1$  then, by the reflexivity of  $L_p(\mathbf{R})$ , there exists a sequence  $(\alpha_n)$  in  $(\pi/2, \pi)$  with  $\lim_{n \rightarrow \infty} \alpha_n = \pi/2$  and a function  $g \in L_p(\mathbf{R})$  so that  $g$  is the weak limit of  $(g_{\alpha_n})$  in  $L_p(\mathbf{R})$ . This implies for all  $z \in \mathbf{C}_+$

$$\begin{aligned} \|F(z)\| &= \lim_{n \rightarrow \infty} \|F_{\alpha_n}(z)\| \leq \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} P_z(t) \|f_{\alpha_n}(t)\| dt \\ &= \int_{-\infty}^{\infty} P_z(t) g(t) dt =: G(z). \end{aligned}$$

If  $p = 1$  then, by Helly's theorem, there exists a sequence  $(\alpha_n)$  in  $(\pi/2, \pi)$  with  $\lim_{n \rightarrow \infty} \alpha_n = \pi/2$  and a function  $\phi : \mathbf{R} \rightarrow \mathbf{R}$  of bounded variation so that  $\phi$  is the weak\* limit of  $(g_{\alpha_n})$  in  $C_0(\mathbf{R})^*$ . Consequently,

$$\begin{aligned} \|F(z)\| &= \lim_{n \rightarrow \infty} \|F_{\alpha_n}(z)\| \leq \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} P_z(t) \|f_{\alpha_n}(t)\| dt \\ &= \int_{-\infty}^{\infty} P_z(t) d\phi(t) =: G(z). \end{aligned}$$

It follows e.g. from [2], I.3. that, in either case,  $G : \mathbf{C}_+ \rightarrow \mathbf{R}$  is a harmonic function with

$$\sup_{a>0} \int_{-\infty}^{\infty} |G(a+ib)|^p db \leq \sup_n \|g_{\alpha_n}\|_{L_p}^p \leq 2^p \|F\|_{H_p(\Sigma_{\pi/2})}^p.$$

We finally conclude

$$\sup_{a>0} \int_{-\infty}^{\infty} \|F(a+ib)\|^p db \leq \sup_{a>0} \int_{-\infty}^{\infty} \|G(a+ib)\|^p db \leq 2^p \|F\|_{H_p(\Sigma_{\pi/2})}^p. \quad \equiv$$

## 2 Boundary values of vector-valued holomorphic functions

We investigate now the boundary behavior of vector-valued functions being holomorphic in a sector

$$\Sigma_\alpha = \{re^{i\theta} : r > 0, |\theta| < \alpha\}.$$

Similar to  $H_p(\Sigma_{\pi/2}, X)$  we introduce for each  $0 < \alpha < \pi/2$  the space  $H_p(\Sigma_\alpha, X)$  which consists of holomorphic functions  $F : \Sigma_\alpha \rightarrow X$  such that

$$\|F\|_{H_p(\Sigma_\alpha)} = \sup_{|\theta| < \alpha} \left( \int_0^\infty \|F(re^{i\theta})\|^p dr \right)^{1/p} < \infty$$

if  $p < \infty$ , and such that

$$\|F\|_{H_\infty(\Sigma_\alpha)} = \sup_{z \in \Sigma_\alpha} \|F(z)\| < \infty$$

if  $p = \infty$ . We start with a study of the boundary behavior of harmonic functions  $F : \mathbf{C}_+ \rightarrow X$ , and then use this result for a description of the boundary values of  $H_p(\Sigma_\alpha, X)$ -functions. For this reason, let us introduce the following spaces of functions and classes of operators: The space  $h_p(\mathbf{C}_+, X)$  consists of harmonic functions  $F : \mathbf{C}_+ \rightarrow X$  for which the following norm is finite:

$$\|F\|_{h_p} := \sup_{a > 0} \left( \int_{-\infty}^\infty \|F(a + it)\|^p dt \right)^{1/p}$$

if  $p < \infty$ , and  $\|F\|_{h_\infty} := \sup_{z \in \mathbf{C}_+} \|F(z)\|$  if  $p = \infty$ . We say that  $F$  has a boundary function  $f \in L_p(\mathbf{R}, X)$  if  $F$  is the Poisson integral of  $f$ .

Let  $1 \leq p \leq \infty$  and  $1/p + 1/q = 1$ . Put  $Y_q(\mathbf{R}) := L_q(\mathbf{R})$  if  $q < \infty$ , and let  $Y_\infty(\mathbf{R})$  be the space of complex-valued continuous functions on  $\mathbf{R}$  which vanish at infinity. An operator  $T : Y_q(\mathbf{R}) \rightarrow X$  is called  $p$ -bounded,  $p > 1$ , if there exists a function  $f \in L_p(\mathbf{R})$  such that

$$\|Tg\| \leq \int_{-\infty}^\infty |g(t)|f(t) dt, \quad g \in Y_q(\mathbf{R}),$$

and  $T$  is called 1-bounded if there exists a function  $\phi : \mathbf{R} \rightarrow \mathbf{R}$  of bounded variation such that

$$\|Tg\| \leq \int_{-\infty}^\infty |g(t)| d\phi(t), \quad g \in Y_\infty(\mathbf{R}).$$

It follows from [3]. IV.4, Theorem 8, that for  $p > 1$  an operator  $T$  is  $p$ -bounded if and only if it is order summing, and [9], Proposition 1.3.22 states that an operator  $T : Y_\infty(\mathbf{R}) \rightarrow X$  is 1-bounded if and only if it is absolutely summing. We say that  $T$  is represented by  $g \in L_p(\mathbf{R}, X)$  if

$$Th = \int_{-\infty}^\infty h(t)g(t) dt, \quad h \in Y_q(\mathbf{R}).$$

If  $p > 1$  and if  $X$  has the Radon-Nikodym property then it follows from [3], IV.4, Theorem 8, that every  $p$ -bounded operator can be represented by an  $L_p(\mathbf{R}, X)$ -function.

**THEOREM 2** *Let  $1 \leq p \leq \infty$ . A harmonic function  $F : \mathbf{C}_+ \rightarrow X$  belongs to  $h_p(\mathbf{C}_+, X)$  if and only if there exists a  $p$ -bounded operator  $T : L_q(\mathbf{R}) \rightarrow X$  such that*

$$F(z) = T(P_z) = \lim_{a \rightarrow 0^+} \int_{-\infty}^{\infty} P_z(t) F(a + it) dt, \quad z \in \mathbf{C}_+.$$

*If  $1 < p \leq \infty$  and if  $X$  has the Radon-Nikodym property then  $F$  has a boundary function  $f \in L_p(\mathbf{R}, X)$ .*

*Proof.* Let  $F \in H_p(\mathbf{C}_+, X)$ . Then, for every  $a > 0$ , the function  $F_a : \mathbf{C}_+ \rightarrow X$  defined by  $F_a(z) = F(a + z)$  belongs to  $h_p(\mathbf{C}_+, X)$ , and  $\|F_a\|_{h_p} \leq \|F\|_{h_p}$ . Since  $F_a$  is bounded and continuous in the closed right half-plane  $\overline{\mathbf{C}}_+$  (this follows e.g. from [2], I.3.)  $F_a$  is the Poisson integral of its boundary function (see [2], Lemma I.3.4. for the scalar case). It follows that

$$F(z) = \lim_{a \rightarrow 0^+} F_a(z) = \lim_{a \rightarrow 0^+} \int_{-\infty}^{\infty} P_z(t) F_a(it) dt.$$

Collecting these informations yields:

1. For all  $h$  in the total subset  $\{P_z : z \in \mathbf{C}_+\}$  of  $Y_q(\mathbf{R})$  the limit of  $T_a h := \int_{-\infty}^{\infty} h(t) F_a(t) dt$  exists, as  $a \rightarrow 0^+$ .
2. The family  $(T_a)$  of operators from  $Y_q(\mathbf{R})$  into  $X$  is uniformly bounded by  $\|T_a\| \leq \|F\|_{H_p(\mathbf{C}_+)}$ .

Hence, by the Banach-Steinhaus-Theorem, there exists an operator  $T : L_q(\mathbf{R}) \rightarrow X$  with  $\|T\| \leq \|F\|_{H_p(\mathbf{C}_+)}$ , and such that

$$Th = \lim_{a \rightarrow 0^+} \int_{-\infty}^{\infty} h(t) F_a(t) dt = \lim_{a \rightarrow 0^+} \int_{-\infty}^{\infty} h(t) F(a + it) dt. \quad (1)$$

Moreover, the operator  $T$  is  $p$ -bounded. To see this, consider the functions  $g_a \in L_p(\mathbf{R})$  defined by  $g_a(t) = \|F_a(it)\|$ . Then the family  $\{g_a : a > 0\}$  is bounded in  $L_p(\mathbf{R})$ . Consequently, if  $p > 1$ , there exists a positive null-sequence  $(a_n)$  so that  $(g_{a_n})$  has a weak limit  $g \in L_p(\mathbf{R})$ , if  $p < \infty$ , and so

that  $(g_{a_n})$  has a weak\* limit  $g \in L_\infty(\mathbf{R})$ , if  $p = \infty$ . If  $p = 1$  then, by Helly's theorem,  $(a_n)$  can be chosen so that  $(g_{a_n})$  has a weak\* limit  $\psi : \mathbf{R} \rightarrow \mathbf{R}$  of bounded variation. Consequently,

$$\begin{aligned} \|Th\| &\leq \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} |h(t)| \|F_{a_n}(it)\| dt \\ &= \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} |h(t)| g_{a_n}(t) dt = \int_{-\infty}^{\infty} |h(t)| g(t) dt \end{aligned}$$

if  $p > 1$ , and

$$\|Th\| \leq \int_{-\infty}^{\infty} |h(t)| d\psi(t)$$

if  $p = 1$ .

If  $X$  has the Radon-Nikodym property, then, according to [3], IV.4, Theorem 8, for  $1 < p \leq \infty$  every  $p$ -bounded operator can be represented by a function  $f \in L_p(\mathbf{R}, X)$ .  $\equiv$

The above considerations for  $h_p(\mathbf{C}_+, X)$ -functions clearly hold also for  $H_p(\mathbf{C}_+, X)$ -functions. Our next goal is to show that for  $p > 1$  similar assertions are true for functions  $G \in H_p(\Sigma_\alpha, X)$ . Therefore, we exchange the cartesian coordinates “ $a + it$ ” by polar coordinates “ $te^{i\eta}$ ”, and for each  $0 < \eta < \alpha$  we introduce the function  $G_\eta(t) = G(|t|e^{i\text{sgn}(t)\eta})$ ,  $t \in \mathbf{R}$ . We say that  $G$  has a boundary function  $g \in L_p(\mathbf{R}, X)$  if

$$\lim_{\eta \rightarrow \alpha^-} \int_{-\infty}^{\infty} h(t) G_\eta(t) dt = \int_{-\infty}^{\infty} h(t) g(t) dt, \quad h \in L_q(\mathbf{R}).$$

**THEOREM 3** *Let  $1 < p \leq \infty$  and  $0 < \alpha < \pi$ . For every  $G \in H_p(\Sigma_\alpha, X)$  there exists a  $p$ -bounded operator  $T : L_q(\mathbf{R}) \rightarrow X$  such that*

$$Th = \lim_{\eta \rightarrow \alpha^-} \int_{-\infty}^{\infty} h(t) G_\eta(t) dt, \quad h \in L_q(\mathbf{R}).$$

*If  $1 < p \leq \infty$  and if  $X$  has the Radon-Nikodym property then  $G$  has a boundary function  $g \in L_p(\mathbf{R}, X)$ .*

*Proof.* The proof is divided into two steps. In the first step we consider the case  $\alpha = \pi/2$ , and in the second step we prove the assertion for general  $\alpha$ .

Step 1: Let  $G \in H_p(\Sigma_{\pi/2}, X)$  and for  $t \in \mathbf{R}$  define  $\chi_t = \chi_{[0,t]}$  if  $t > 0$ ,  $\chi_t = -\chi_{[t,0]}$  if  $t < 0$  and  $\chi_0 = 0$ . We show first that the limit

$$T\chi_t := \lim_{\eta \rightarrow \alpha^-} \int_0^t G_\eta(u) du \quad (2)$$

exists for every real number  $t$ . We know by Theorem 1 that  $G$  belongs to  $H_p(\mathbf{C}_+, X)$ . Hence, by Theorem 2 there exists a bounded operator  $T : L_q(\mathbf{R}) \rightarrow X$  with

$$Th = \lim_{c \rightarrow 0^+} \int_{-\infty}^{\infty} h(t) G(c + it) dt. \quad (3)$$

Let  $z = a + ib \in \mathbf{C}_+$ . Then there exists  $t \in \mathbf{R}$  and  $0 \leq \eta < \frac{\pi}{2}$  so that  $z = |t|e^{i\operatorname{sgn}(t)\eta}$ . We define three paths  $\Gamma_1(z)$ ,  $\Gamma_2(z)$  and  $\Gamma_3(z)$  in the following way:

$$\begin{aligned} \Gamma_1(z) &= (se^{i\operatorname{sgn}(t)\eta} : 0 \leq s \leq |t|), \\ \Gamma_2(z) &= (s : 0 \leq s \leq a), \\ \Gamma_3(z) &= (a + i\operatorname{sgn}(t)s : 0 \leq s \leq |b|). \end{aligned}$$

Then  $(\Gamma_1(z), \Gamma_2(z), \Gamma_3(z))$  is a closed path in  $\overline{\mathbf{C}}_+$ . Since  $G$  is holomorphic in  $\mathbf{C}_+$ , and since  $\|G(z)\| \leq K\operatorname{Re}(z)^{-1/p}$  (for the scalar case see [4], Lemma on page 149) we infer from Cauchy's theorem

$$\begin{aligned} \int_0^t G_\eta(s) ds &= \operatorname{sgn}(t)e^{-i\operatorname{sgn}(t)\eta} \int_{\Gamma_1(z)} G(\nu) d\nu \\ &= \operatorname{sgn}(t)e^{-i\operatorname{sgn}(t)\eta} \left( \int_{\Gamma_2(z)} G(\nu) d\nu + \int_{\Gamma_3(z)} G(\nu) d\nu \right) \\ &= \operatorname{sgn}(t)e^{-i\operatorname{sgn}(t)\eta} \left( \int_0^a G(s) ds + i \int_0^b G(a + is) ds \right). \quad (4) \end{aligned}$$

Let  $t \in \mathbf{R}$  and put  $z(\eta) = |t|e^{i\operatorname{sgn}(t)\eta} = a(\eta) + ib(\eta)$ . If  $\eta$  converges towards  $\pi/2$  then  $b(\eta)$  converges towards  $t$ . Consequently, equation (3) yields

$$\begin{aligned} &\lim_{\eta \rightarrow \pi/2} i\operatorname{sgn}(t)e^{-i\operatorname{sgn}(t)\eta} \int_0^{b(\eta)} G(a(\eta) + is) ds \\ &= \lim_{\eta \rightarrow \frac{\pi}{2}} \int_0^t G(a(\eta) + is) ds - \lim_{\eta \rightarrow \frac{\pi}{2}} \int_{b(\eta)}^t G(a(\eta) + is) ds \\ &= T\chi_t, \end{aligned}$$



because

$$\left\| \int_{b(\eta)}^t G(a(\eta) + is) ds \right\| \leq |t - b(\eta)|^{1/q} \|G\|_{H_p} \quad \text{and} \quad \lim_{\eta \rightarrow \pi/2} |t - b(\eta)| = 0.$$

The first part  $\int_0^{a(\eta)} G(s) ds$  in the sum of equation (4) converges towards zero for  $\eta \rightarrow \pi/2$ , because  $\|G(s)\| \leq K s^{-1/p}$ . Hence, equation (2) is proved for  $\alpha = \pi/2$ . Since the set  $\{\chi_t : t \in \mathbf{R}\}$  is total in  $L_q(\mathbf{R})$ , and since the family  $\{G_\eta : -\pi/2 < \eta < \pi/2\}$  is bounded in  $L_p(\mathbf{R}, X)$ , there exists, by the Banach-Steinhaus-Theorem, a bounded operator  $T : L_q(\mathbf{R}) \rightarrow X$  such that, for all  $h \in L_q(\mathbf{R})$

$$Th = \lim_{\eta \rightarrow \pi/2^-} \int_{-\infty}^{\infty} h(t) G_\eta(t) dt.$$

Step 2: Now let  $0 < \alpha < \pi$ . Let  $G \in H_p(\Sigma_\alpha, X)$  and put

$$F(z) = \left(\frac{\pi}{2\alpha}\right)^{1/p} z^{\frac{1}{p}(\frac{\pi}{2\alpha}-1)} G\left(z^{\frac{\pi}{2\alpha}}\right).$$

Then, by straightforward calculations,  $F$  belongs to  $H_p(\Sigma_{\pi/2}, X)$ , and for  $0 < \eta < \alpha$  and  $t \in \mathbf{R}$  it follows that

$$\begin{aligned} G_\eta(t) &= \left(\frac{\pi}{2\alpha}\right)^{1/p} (|t| e^{i \operatorname{sgn}(t)\eta})^{\frac{1}{p}(\frac{\pi}{2\alpha}-1)} F\left((|t| e^{i \operatorname{sgn}(t)\eta})^{\frac{\pi}{2\alpha}}\right) \\ &= \left(\frac{\pi}{2\alpha}\right)^{1/p} (|t| e^{i \operatorname{sgn}(t)\eta})^{\frac{1}{p}(\frac{\pi}{2\alpha}-1)} F_{\eta \frac{\pi}{2\alpha}}(\operatorname{sgn}(t) |t|^{\frac{\pi}{2\alpha}}). \end{aligned}$$

If  $h : \mathbf{R} \rightarrow Y$ ,  $Y$  a Banach space, is a function then define  $Bh : \mathbf{R} \rightarrow Y$  by

$$Bh(t) = \left(\frac{\pi}{2\alpha}\right)^{1/p} |t|^{\frac{1}{p}(\frac{\pi}{2\alpha}-1)} h(\operatorname{sgn}(t) |t|^{\frac{\pi}{2\alpha}}),$$

and define  $Ch : \mathbf{R} \rightarrow Y$  by

$$Ch(t) = \left(\frac{2\alpha}{\pi}\right)^{1/q} |t|^{\frac{1}{q}(\frac{2\alpha}{\pi}-1)} h(\operatorname{sgn}(t) |t|^{\frac{2\alpha}{\pi}}).$$

It is easy to see that the operators  $B$  and  $C$  are isometric isomorphisms on  $L_p(\mathbf{R}, Y)$ . Moreover, put  $d_\eta(t) = e^{i \operatorname{sgn}(t) \frac{2\alpha}{p}(\frac{\pi}{2\alpha}-1)}$ . Then for  $h \in L_q(\mathbf{R})$

$$\begin{aligned} \int_{-\infty}^{\infty} h(t) G_\eta(t) dt &= \int_{-\infty}^{\infty} h(t) d_\eta(t) (B F_{\eta \frac{\pi}{2\alpha}})(t) dt \\ &= \int_{-\infty}^{\infty} d_\eta(u) (Ch)(u) F_{\eta \frac{\pi}{2\alpha}}(u) du. \end{aligned}$$

By Step 1,

$$Sh := \lim_{\theta \rightarrow \pi/2^-} \int_{-\infty}^{\infty} h(t) F_{\theta}(t) dt$$

defines a bounded operator  $S : L_q(\mathbf{R}) \rightarrow X$ . Consequently, since  $\eta_{\frac{\pi}{2\alpha}}$  converges towards  $\pi/2$  from below, as  $\eta \rightarrow \alpha^-$ , the result of the Step 1 implies that the following limit exists:

$$\lim_{\eta \rightarrow \alpha^-} \int_{-\infty}^{\infty} h(t) G_{\eta}(t) dt =: Th, \quad h \in L_q(\mathbf{R}).$$

Clearly,  $T : L_q(\mathbf{R}) \rightarrow X$  is a bounded operator. In order to show that  $T$  is  $p$ -bounded, put  $g_{\eta}(t) = \|G_{\eta}(t)\|$ . Then there exists a sequence  $(\eta_n)$  converging towards  $\alpha$  from below, so that  $(g_{\eta_n})$  has a weak limit  $g \in L_p(\mathbf{R})$ , if  $p < \infty$ , and so that  $(g_{\eta_n})$  has a weak\* limit  $g \in L_{\infty}(\mathbf{R})$ , if  $p = \infty$ . Consequently, for  $h \in L_q(\mathbf{R})$

$$\begin{aligned} \|Th\| &\leq \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} |h(t)| \|G_{\eta_n}(it)\| dt \\ &= \limsup_{n \rightarrow \infty} \int_{-\infty}^{\infty} |h(t)| g_{\eta_n}(t) dt = \int_{-\infty}^{\infty} |h(t)| g(t) dt. \end{aligned}$$

If  $X$  has the Radon-Nikodym property it follows from [3], IV.4, Theorem 8, that  $T$  can be represented by a function  $g \in L_p(\mathbf{R}, X)$ .  $\equiv$

### 3 A Paley-Wiener-type theorem for $H_2(\Sigma_{\alpha}, X)$

The classical Paley-Wiener-Theorem [7] states that the complex Laplace transform  $F(z) = \int_0^{\infty} e^{-zt} f(t) dt$  is an isometric isomorphism between the spaces  $L_2([0, \infty))$  and  $H_2(\mathbf{C}_+, \mathbf{C})$  (see also [8], Chapter 19). The proof of this theorem uses heavily the Plancherel theorem for the Fourier transform. By Kwapien's theorem [5] a Banach space valued version of Plancherel's theorem does not exist, in general, but the Fourier transform maps  $L_2(\mathbf{R}, X)$  isomorphically onto  $L_2(\mathbf{R}, X)$  if and only if  $X$  is isomorphic to a Hilbert space. Hence, for Hilbert spaces  $X$ , the following vector-valued version of the Paley-Wiener-Theorem theorem can be proved (see [9], Proposition 2.2.14).

**THEOREM 4** *If  $X$  is isomorphic to a Hilbert space then the  $X$ -valued complex Laplace transform*

$$F(z) = \int_0^\infty e^{-zt} f(t), \quad z \in \mathbf{C}_+,$$

*is an isomorphism from  $L_2([0, \infty), X)$  onto  $H_2(\mathbf{C}_+, X)$ .*

The following corollary is an immediate consequence of this vector-valued version of the Paley-Wiener-Theorem together with Theorem 1.

**COROLLARY 5** *If  $X$  is isomorphic to a Hilbert space then the vector-valued complex Laplace transform*

$$F(z) = \int_0^\infty e^{-zt} f(t), \quad z \in \mathbf{C}_+,$$

*is an isomorphism from  $L_2([0, \infty), X)$  onto  $H_2(\Sigma_{\pi/2}, X)$ .*

We now want give the outline of a proof for a theorem of Paley-Wiener type for the complex Laplace transform of functions belonging to  $H_2(\Sigma_\alpha, X)$ ; more precise:

**THEOREM 6** *Let  $0 < \alpha < \pi/2$ , and let  $X$  be isomorphic to a Hilbert space. Then the complex Laplace transform of any function  $f \in H_2(\Sigma_\alpha, X)$  can be extended holomorphically to a unique function  $F \in H_2(\Sigma_{\alpha+\pi/2}, X)$ , and every  $F \in H_2(\Sigma_{\alpha+\pi/2}, X)$  is the extension of the complex Laplace transform of a unique  $f \in H_2(\Sigma_\alpha, X)$ .*

*Proof.* (an outline) If  $f$  belongs to  $H_2(\Sigma_\alpha, X)$  then one defines a holomorphic function  $F : \Sigma_{\pi/2} \rightarrow X$  in the following way: For  $-\alpha < \theta < \alpha$  define the path  $\Gamma_\theta = (re^{i\theta} : 0 < r < \infty)$ . Now, given  $z \in e^{-i\theta}\mathbf{C}_+$  put

$$F_\theta(z) = \int_{\Gamma_\theta} e^{-z\mu} F(\mu) d\mu.$$

Then  $F_\theta : e^{-i\theta}\mathbf{C}_+ \rightarrow X$  is a holomorphic function, and by Corollary 5

$$\sup_{|\beta| < \pi/2} \int_0^\infty \|F_\theta(re^{i(\theta+\beta)})\|^2 dr \leq M^2 \|f\|_{H_2(\Sigma_\alpha, X)}^2, \quad (5)$$

where  $M$  denotes the norm of the vector-valued complex Laplace transform from  $L_2([0, \infty), X)$  into  $H_2(\Sigma_{\pi/2}, X)$ . Moreover, one can show that  $F_\theta(z) = F_\eta(z)$  whenever  $-\alpha < \theta, \eta < \alpha$  and  $z \in e^{-i\theta}\mathbf{C}_+ \cap e^{-i\eta}\mathbf{C}_+$ . Since  $\bigcup_{|\theta| < \alpha} e^{-i\theta}\mathbf{C}_+ = \Sigma_{\alpha+\pi/2}$  we can define  $F : \Sigma_{\alpha+\pi/2} \rightarrow X$  by  $F(z) = F_\theta(z)$ , where  $\theta$  is chosen so that  $z$  belongs to  $e^{-i\theta}\mathbf{C}_+$ . Then it follows from (5) that  $F$  belongs to  $H_2(\Sigma_{\alpha+\pi/2}, X)$  with  $\|F\|_{H_2(\Sigma_{\alpha+\pi/2})} \leq M \|f\|_{H_2(\Sigma_\alpha)}$ , and  $F|_{\mathbf{C}_+}$  is the complex Laplace transform of  $f$ .

Conversely, assume  $F$  to belong to  $H_2(\Sigma_{\alpha+\pi/2}, X)$ . Then, by [9], Satz 2.3.7,  $F$  is the complex Laplace transform of a holomorphic function  $f : \Sigma_\alpha \rightarrow X$ , and  $F(z)$  can be evaluated by

$$F(z) = \int_{\Gamma_\theta} e^{-z\mu} f(\mu) d\mu, \quad z \in e^{-i\theta}\mathbf{C}_+.$$

It remains to be shown that  $f$  belongs to  $H_2(\Sigma_\alpha, X)$ . Let  $|\theta| < \alpha$ , and, for  $z \in \mathbf{C}_+$ , put  $F_\theta(z) = F(ze^{-i\theta})$ . Then, by Corollary 5,  $F_\theta$  is the complex Laplace transform of a function  $f_\theta \in L_2([0, \infty), X)$ . Hence, for  $z = re^{-i\theta}$  it follows that

$$\begin{aligned} e^{i\theta} \int_0^\infty e^{-rt} f(te^{i\theta}) dt &= \int_{\Gamma_\theta} e^{-z\mu} f(\mu) d\mu \\ &= F(z) = F_\theta(r) = \int_0^\infty e^{-rt} f_\theta(t) dt. \end{aligned}$$

By uniqueness of the Laplace transform it follows that  $e^{i\theta}f(te^{i\theta}) = f_\theta(t)$  almost everywhere in  $[0, \infty)$ . Consequently,

$$\int_0^\infty \|f(re^{i\theta})\|^2 dr = \|f_\theta\|_{L_2}^2 \leq L^2 \|F\|_{H_2(\Sigma_{\alpha+\pi/2})}^2,$$

where  $L$  denotes the norm of the inverse complex Laplace transform from  $H_2(\mathbf{C}_+, X)$  into  $L_2([0, \infty), X)$ .  $\equiv$

## References

- [1] M.M. DRŽBAŠJAN, V.M. MARTIROSIAN, Theorems of Paley-Wiener and Müntz-Szász type, *Soviet Math. Dokl.* **16** (1975), 1559-1563.

- [2] J.B. GARNETT, “Bounded Analytic Functions”, Academic Press, New York, 1981.
- [3] J. DIESTEL, J.J. UHL, “Vector measures”, American Mathematical Society, Providence, 1977.
- [4] P. KOOSIS, “Introduction to  $H_p$  Spaces”, Cambridge University Press, Cambridge, 1980.
- [5] S. KWAPIEN, Isomorphic characterizations of inner product spaces by orthogonal series with vector valued coefficients, *Studia Math.* **44** (1972), 583-595.
- [6] W.A.J. LUXEMBURG, On an inequality of Levinson of the Phragmen-Lindelöf type, *Indagationes Math.* **46**, (1984), 421-427.
- [7] R.E.A.C. PALEY, N. WIENER, “Fourier Transforms in the Complex Domain”, Amer. Math. Soc. Colloquium Publ. 19, New York, 1934.
- [8] W. RUDIN, “Real and Complex Analysis”, McGraw-Hill Book Company, New York, 1987.
- [9] P. VIETEN, “Holomorphie und Laplace Transformation banachraumwertiger Funktionen”, Shaker, Achen, 1995.
- [10] A.M. SEDLECKIĚ, An equivalent definition of  $H^p$  spaces in the half-plane and some applications, *Math. U.S.S.R. Sbornik* **25**, (1975), 69-75.

Peter Vieten  
Fachbereich Mathematik  
Universität Kaiserslautern  
Erwin-Schrödinger-Straße  
67663 Kaiserslautern  
Germany

vieten@mathematik.uni-kl.de